

Outcome of AO External Fixator for Open Tibial Fractures

1. Malik Mohammad Yasin Awan 2. Rashid Hussain 3. Ijaz Ahmed

1. Asstt. Prof. of Orthopedics, Gujranwala Medical College, Gujranwala 2. Medical Officer of Orthopedics, DHQ Teaching Hospital, Gujranwala 3. Asstt. Prof. GTTH, Lahore / LMDC, Lahore

ABSTRACT

Objectives: The purpose of the study was to evaluate the outcome of open tibial shaft fractures (Gustilo IIIA/B) with AO External Fixator & to document complications

Study Design: Perspective study

Place and Duration of Study: This study was conducted at DHQ teaching Hospital, Gujranwala & at Fazil Memorial Hospital, Gujranwala from November 2010 to November 2012

Material & Methods: 50 patients with open tibial shaft fractures were treated with AO External Fixator. Open fractures were classified according to Gustilo Anderson Criteria & wounds with IIIA & IIIB were selected. Outcome was determined by the rate of union, while nonunion, pin tract infection, pin loosening & osteomyelitis were recorded as complications. The follow up period was 08 months.

Results: Out of fifty cases of open tibial shaft fractures, 38 (76%) were men & 12 (24%) were Ladies. Mean age was 35.2 (8-67), 22 (44%) had Gustilo IIIB wound while 28 (56%) had Gustilo IIIA injuries. Pin tract infection & pin loosening rate were 12% & 16% respectively. Nonunion was seen in 10% of the cases. In 41 (82%) of the cases fracture united and average union time was about 26.5 weeks. No case of osteomyelitis seen.

Conclusion: External Fixator is simple & effective treatment for open tibial fractures.

Key Words: Open Tibial Fractures, External Fixator, Gustilo Anderson's Classification, Union

Citation of article: Awan MMY, Hussain R, Ahmed I. Outcome of AO External Fixator for Open Tibial Fractures. Med Forum 2015;26(3):29-32.

INTRODUCTION

Open fractures of the lower limb are potentially devastating. The incidence of open long bone fracture is around 11.5 per 100,000 persons per year in United Kingdom¹. Open fractures can be classified in many ways but the classification devised by Gustilo & Anderson is most common and is used worldwide². Nonunion & infection is quite common in tibial fractures owing to deficient blood supply & soft tissue coverage, so the treatment is controversial^{3, 4}. Open tibial fractures are one of the commonest, complex & unexpected injuries in orthopedic practice⁵. Commonest causes of these fractures are road traffic accidents, firearm injuries, fall from height, fall of heavy objects & industrial injuries⁶. Now the blast injuries are emerging as one of the leading cause in certain areas of our country⁷. Treatments allow anatomical realignment, early mobilization which helps to get early pre-injury status⁸.

Treatment includes bony fixation as well as soft tissue management⁹. Fixation can be external or internal. External fixation is preferred treatment in open fractures, although internal fixation with interlocking

nail can be safely done, if patient is operated within golden period^{10, 11}. External fixator can be applied for comminuted closed tibial fractures. Due to subcutaneous location of tibia, fixator is easy to apply¹². There are certain problems with external fixation which has outweighed its popularity. These are pin tract infections, pin loosening & osteomyelitis¹³.

The aim of the study was to determine the outcome of AO external fixator in the treatment of open tibial shaft fractures in term of union & to document complications i.e. pin tract infection, pin loosening, osteomyelitis & nonunion.

MATERIALS AND METHODS

Study was conducted at DHQ teaching hospital, Gujranwala & at Fazil Memorial Hospital, Gujranwala between November 2010 to November 2012 (Two years). After taking informed consent, patients with Gustilo IIIA/IIIB were selected. Patients having other injuries or Gustilo IIIC fractures were excluded from the study. Demographic information including age, sex mode of injury i.e Road Traffic accident (RTA), fall from height (FH), fall of heavy object (FHO), Firm Arm injury (FAI), Blast (B) & industrial injuries (IND) were recorded.

All the information was recorded on a specified Proforma.

All patients were admitted through emergency where clinical evaluation done, wound washed & limb

Correspondence: Malik Mohammad Yasin Awan,
Asstt. Prof. of Orthopedics, Gujranwala Medical College,
Gujranwala.
Cell No: 03214940269
Email: Orthoandiyasin@yahoo.com

splinted. Necessary laboratory & radiological investigations were carried out. Tetanus prophylaxis, analgesia & I/V antibiotics were given. Patients were operated within 24 hours of presentation. Fractures were fixed with AO external fixator & wound debrided. Post operatively check X-rays were obtained, daily dressing done, range of motion started & IV antibiotics given for three days. Patient discharged at third post operative day. Discharge instructions were given including daily dressing, care of pin site, range of motion at knee and ankle & walking with support. Flow up plan was every 2nd week till two months & then bimonthly. Radiographs were obtained at first day, first month & then on every follow up. On every visit patients were examined for union & complications i.e pin tract infection, pin loosening, nonunion & osteomyelitis. Union was defined as bridging callus crossing three of the four cortices on both AP & lateral radiographs with no pain on pressure or weight bearing. Pin site/tract infection was determined by clinical signs of local erythema, swelling, tenderness, fever, pus discharge or positive bacterial culture. Pin loosening was documented when there was new onset of pain at pin site along with clinical and / or radiographic loosening (radiolucent line along the schanz entry). Nonunion was defined as an absence of bridging callus across a fracture site for expected time (last follow up in our study). Osteomyelitis was defined as clinical signs of fever, swelling, redness, pus discharge along with radiological evidence of sequestrum at fracture site or schanz entry site.

Data was entered & analyzed by SPSS version 10.

RESULTS

Out of fifty patients, 38(76%) were male & 12(24%) were female (figure 2). Youngest patient was of 8 years old & oldest person was 67 years old.




Figure No.1: Types of Fractures

Average age was 35.2 years (8-67). Road traffic accidents were the leading cause in our study with thirty two 32 (64%) Cases. Other injuries include firearm injuries (FAI) 8 (16%), fall from height 5 (10%), Fall of heavy object 2 (4%) while 3 (6%) cases were industrial injuries (figure 3). Pin tract infection

was in 6 cases, 12% & pin loosening was in eight patients, 16%. All pin loosening occurred after three months. Nonunion was seen in 5 subjects (10%) & no case of osteomyelitis seen. 22 patient had Gustilo IIIB injuries while 28 patients had Gustilo IIIA injuries (figure 1). Union achieved in 82% of the cases while 8% of the cases developed malunion.

Table No.1: Gustilo and Anderson classification^{2,14}

Gustilo Grade	Definition
I	Open fracture, clean wound <1 cm in length
II	Open fracture, wound > 1 cm but < 10 cm in length without extensive soft-tissue damage, flaps, avulsions.
III	Open fracture with extensive soft-tissue laceration (>10 cm) damage, or an open segmental fracture. This type also includes open fractures caused by firearm injuries, fractures requiring vascular repair, or fractures that have been open for 8 hours prior to treatment.
IIIA	Type III fracture with adequate periosteal coverage of the fractured bone despite the extensive soft- tissue laceration or damage
IIIB	Type III fracture with extensive soft-tissue loss and periosteal stripping and bone damage, Usually associated with massive contamination. Will often need further soft-tissue coverage procedure (i.e free or rotational flap)
IIIC	Type III fracture associated with an arterial injury requiring repair, irrespective of degree of soft-tissue injury.




Figure No.2: Gender Distribution




Figure No.3: Etiology




Figure No.4: Complications

DISCUSSION

Our study showed 76% of the male involvement which other studies also have proved (fig 2) ^{14,15,16,17}. Operative treatment of the tibial shaft fractures usually leads to healing without any consequences on life & working ability¹². The most common methods used in treating tibial shaft fractures are intramedullary nail, AO plates & external Fixator¹⁸.

The external fixator in open tibial fractures not only solves the problem of managing soft tissue injuries but also provide a reasonable fixation for bone to heal. With AO fixator it is possible to adhere to safe & effective fixation techniques, avoid damage to vital structures, have access to wound & adjust the fixator so that it is biomechanically compatible with fractures ¹⁹.

Mean age in our study was 35.2 years which was quite comparable with other studies. All these studies have shown that these injuries occur in younger age group^{20,21,22,23}. Mean time of fracture healing in our study was 26.5 weeks. Tucker et al ²⁴ and schatzker²⁵ in separate studies reported union time of 25.6 weeks and 21.9 weeks respectively. Similarly Wheelwright, Court-Brown ²⁶ and Adrover et al ²⁷ reported a union time of 27.5 weeks & 26 weeks respectively. The union & nonunion rate in our study was 82% & 10% respectively. Kaftandziev²⁸ in his study produced union in 71.1% while Bratislav stojkovic ²⁹ reported a union rate of 83.68% in his 49 patients.

In our study pin site/ tract infection rate was 12% & pin loosening was 16% (fig4). In a study by Parameswarma AD et al infection rate was 11.2% in 285 fractures. The incidence of Pin tract infection for the unilateral fixator group and the hybrid fixator group were not significant different^{29, 30}. In another study average age of union was 21 weeks, Infection was 8%, nonunion was 11%, these results were produced in 1438 subjects by song k et al³¹. Shalamon J et al in his study proved that most of the Pin tract infection are mild and can be treated by

local or systemic antibiotics. Loosening did not require a change of method of stabilization³².

CONCLUSION

External fixator is simple & effective mean of treating open tibial fractures.

REFERENCES

1. Soni A, Tzafetta K, Knight S. Gustilo III C fractures in lower limb. *J Bone Joint Surg Br* 2012; 94-B:698-703.
2. George W, Wood II. General Principles of Fracture Treatment. *Campbell's Operative Orthopaedics* 2006; 10(3):2670-71.
3. Esterhai JL, Queenan J. Management of soft tissue wounds associated with type III open fractures. *Orthop Clin North Am* 1991; 22:427-32.
4. Gustilo RB, Mendoza RM, Williams DN. Problems in the management of type III (severe) open fractures: a new classification of type III open fractures. *J Trauma* 1984;24:742-6.
5. Tosic A, Ebraheim NA, Abou Chakra I, Emara K. External fixator in the management of proximal tibial fractures in adults. *Orthopedics* 2001;24: 581-4.
6. Dougherty PJ, Vaidya R, Silverton CD, Bartlett C, Najibi S. Joint and long-bone gunshot injuries. *J Bone Joint Surg* 2009;91:980-97.
7. Johnson EN, Burns TC, Hayda RA, Hospenthal DR, Murray CK. Infectious complications of open type III tibial fractures among combat casualties. *Clin Infect Dis* 2007;45:409-15.
8. Boyton DM, Schmeling JG. Nonreamed Intramedullary Nailing of Open Tibial Fractures. *J Am Acad Orthop Surg* 1994;2(2):107-114.
9. Ruedi T, Murhy W. AO principles of fracture management, Thieme, Stuttgart-New York 2000.
10. Emami A, Mjoberg B, Karlstrom G, Larsson S. Treatment of closed tibial shaft fractures with unilateral external fixation. *Injury* 1995;26(5): 299-303.
11. Oh CW, Park BC, Kyung HS, Kim SJ, Kim HS, Lee SM, Ihn JC. Percutaneous plating for unstable tibial fractures. *J Orthop Sci* 2003;8(2):166-9.
12. Mitkovic M. New concepts in external fixation. *Prosveta Nis* 1993.
13. Velazco A, Fleming LL. Open fractures of the tibia treated by the Hoffmann external fixator. *Clin Orthop* 1983;180:125-32.
14. Gustilo RB. Current concepts in the management of open fractures. *AAOS Instr Course Lect* 1987; 36:359-66.
15. Trafton PG. Closed unstable fractures of the tibia. *Clin Orthop* 1988; 230: 58-67.
16. Jensen JS, Hansen FW, Johansen J. Tibial shaft fractures. A comparison of conservative treatment

- and internal fixation with conventional plates or AO compression plates. *Acta Orthop Scand* 1977; 48(2):204-12.
17. Khan MA, Khan SW, Qadir RI. Role of external fixator in the management of type II & III open tibial fracture. *J Postgrad Med Inst* 2004;18:12-7.
 18. Holbrook JL, Swiontkowski MF, Sanders R. Treatment of open fractures of the tibial shaft: Ender nailing versus external fixation: a randomised, prospective comparison. *J Bone Joint Surg Am* 1989;71-A:1231-8.
 19. Henley MB, Chapman JR, Agel J, et al. Treatment of II, IIIA and IIIB open fractures of the tibial shaft: a prospective comparison of undreamed interlocking intramedullary nails and half-pin external fixators. *J Orthop Trauma* 1998;12:1-7.
 20. Ornetta P III, Bergman M, Watnik N, Berkowitz G, Steuer J. Treatment of grade IIIB open tibial fractures: a prospective randomised comparison of external rotation and non-reamed locked nailing. *J Bone Joint Surg Br* 1994;76-B:13-9.
 21. Tu YK, Lin CH, Su JI, Hsu DT, Chen RJ. Unreamed interlocking nail versus external fixator for open type III tibia fractures. *J Trauma* 1995; 39: 361-7.
 22. Finkemeier CG, Schmidt AH, Kyle RF, Templeman DC, Varecka TF. A prospective, randomized study of intramedullary nails inserted with and without reaming for the treatment of open and closed fractures of the tibial shaft. *J Orthop Trauma* 2000;14:187-93.
 23. Keating JF, O'Brien PJ, Blachut PA, Meek RN, Broekhuysen HM. Locking intramedullary nailing with and without reaming for open fractures of the tibial shaft: a prospective, randomized study. *J Bone Joint Surg Am* 1997;79-A:334-41.
 24. Tucker HL, Kendra JC, Kinnebrew TE. Management of unstable open and closed fractures using the Ilizarov Method. *Clin Orthop* 1992; 280: 125-35.
 25. Schatzker H, David R, Stolerio J, Grimberg B, Soundry M. Treatment of open tibial fractures with primary suture and Ilizarov fixation. *Clin Orthop* 1997; 335:268-74.
 26. Wheelwright EF, Court-Brown CM. Primary external fixation and secondary intramedullary nailing in the treatment of tibial fractures. *Injury* 1992;23:373-6.
 27. Adrover PA, Garin DM, Alvarez JM, Alonso CP. External fixation and secondary intramedullary nailing of open tibial fractures. *J Bone Joint Surg* 1997;79(3):433-37.
 28. Kaftandziev I, Pejnova S, Saveski J. Operative treatment of iii grade open fractures of the tibial diaphysis. *Contribution. Sec Biol Med Sci* 2006; XXVII(I):121-131.
 29. Bratislav S, Saša M, Mile R, Miodrag S, Igor K. Tibial shaft fractures treated by the external fixation method. *Med Biol* 2006;13(3):145-47.
 30. Parameswarma AD, Roberts CS, Seligson D, Voor M. Pin tract infection with Contemporary External fixation: Nonunion how much of a problem? *J Orthop Trama* 2003;17(7):503-7.
 31. Song k, Sangeorzan, Bruce MD, et al. union with external fixator. *J Pediatric Orthop* 1996;16(5): 635-9.
 32. Schalamon J, Petnehazyt, et al. Pin tract infection with External fixation of Pediatric fractures. *J Pediatric Surg* 2007;42(9):1584-7.