

Frequency of Apertognathia in Preadolescent Orthodontic Patients

Apertognathia in Preadolescent Orthodontic Patients

Muhammad Azeem¹, Muhammad Ammar Riaz², Adeela Riaz³, Shumaila Haq¹ and Muhammad Imran Khan¹

ABSTRACT

Objective: The purpose of this study was to calculate the frequency of an apertognathia in preadolescents.

Study Design: Descriptive / Cross Sectional study

Place and Duration of Study: This study was conducted at the Department of Orthodontics, de'Montmorency College of dentistry, Lahore from 1.1.2014 to 31.12.2016.

Materials and Methods: Preadolescents with apertognathia were selected by purposive sampling technique. Plaster models of 100 patients were taken and assessed for the presence of apertognathia. The frequency of apertognathia and its percentage with respect to gender and severity grade was calculated.

Results: The frequency of apertognathia was found to be 10%. Male to female ratio was 1:2.

Conclusion: It was concluded that females are more affected by apertognathia than the males in preadolescents. The high frequency of apertognathia i.e. 10% in preadolescents suggests that there is need to focus on preventive and interceptive orthodontics.

Key Words: Apertognathia, Open bite, Preadolescent

Citation of article: Azeem M, Riaz MA, Riaz A, Haq S, Khan MI. Frequency of Apertognathia in Preadolescent Orthodontic Patients. Med Forum 2017;28(5):74-76.

INTRODUCTION

Apertognathia is a absence of vertical overlapping between upper and lower incisors when the jaws are brought together. It is multifactorial in nature; different etiological factors reported in literature include sucking habits, tongue thrust, TMJ involvement, and amelogenesis imperfecta. Association of cervical vertebral column morphology and head posture issues in preorthodontic patients with apertognathia is already proven. Combining traditional techniques to correct apertognathia have been proposed. Traditional techniques of apertognathia management reported in literature are occlusal bite blocks, modified protraction headgear, anterior vertical elastics, multiloop edgewise archwire therapy, and at present the use of skeletal anchorage devices.

Another term used in describing apertognathia is anterior open bite (AOB). Apertognathia is classified as dental or skeletal, anterior or posterior and unilateral or bilateral.

It may classify as simple or complex. The implication of an apertognathia irrespective of its cause include aesthetic issues, speech and phonetic issues, lingual interposition during deglutition and difficulty in biting from front teeth. Severity grades of apertognathia are: Moderate (0-2 mm), Severe (3-4 mm) and Extreme (more than 4 mm).

The frequency of Apertognathia ranges from 2% to 12% and varies between ethnic groups and by age and sex. In view of the fact that the frequency in different populations is different; this study was design to find out the frequency of apertognathia in preadolescents and its gender distribution.

MATERIALS AND METHODS

This study was conducted at the Department of Orthodontics, de'Montmorency College of Dentistry, Lahore in which orthodontic plaster models of 100 untreated patients, between the chronological ages of 6 and 12 years and irrespective of gender, were included to determine the frequency of apertognathia. Duration of this study was January 2014 to December 2016. The inclusion criteria were, patients having no previous orthodontic treatment and presence of AOB, whereas patients with history of tooth extraction and having craniofacial syndromes were excluded. The amount of AOB was confirmed by measuring interincisal distance with digital vernier callipers on plaster models. The data was analyzed in Statistical Package for the Social Sciences software package (SPSS) 20. The mean age and gender distribution among the selected sample was calculated

¹. Department of Orthodontics, de'Montmorency College of Dentistry, Lahore

². Department of Orthodontics, RHC Machiwal, District Vehari.

³. Department of Orthodontics, BHU, District Vehari

Correspondence: Muhammad Azeem, Assistant Professor, Department of Orthodontics, de'Montmorency College of Dentistry, Lahore.

Contact No: 0345-8409007

Email: dental.concepts@hotmail.com

Received: March 27, 2017;

Accepted: April 29, 2017

RESULTS

The mean age of the patients was 9 years. Out of total sample of 100, 10 (10%) patients had open bite malocclusion. Out of 10, 4 (40%) were males while 6 (60%) were females. 80 % had moderate open bite of <1mm and 20% had >1mm anterior open bite. The male to female ratio was found out to be 1:2. (Table No.1)

Table No. 1: Frequency of Apertognathia among patients (N=100)

Parameter	Frequency
Apertognathia patients	10 (10 %)
Males having Apertognathia	4 (40 %)
Females having Apertognathia	6 (60 %)
<1mm Apertognathia	80%
>1mm Apertognathia	20%

DISCUSSION

The incidence of apertognathia in our study was found to be 10%. The results of this study revealed that moderate type was more prevalent than severe type of Apertognathia. The male to female ratio in our study was 1:2. This is close to the Belgian ratio of 2:3 but in contrast to the findings of certain studies, where no gender differences were found.^{17,18}

Although several reported studies concluded the frequency of apertognathia in 6-12 year old patients, the findings are difficult to compare and contrast; in part, because of different methodologies, age differences of sample, interexaminer variations, and the different sample sizes. Frequency of apertognathia in the present study was found out to be lower than that reported in Argentinean, Uganda, Hawaii and Kuwaitis where the studies revealed the percentage range of 2-4%.¹⁹⁻²² However, results are similar to certain studies where frequency of apertognathia was found out to be in range of 8-10%.²³⁻²⁶

No study has been reported specifically in our Punjab population regarding frequency of apertognathia in preadolescent subjects. Results are similar to those of Ahmed et al. where they found frequency of apertognathia in preadolescent subjects of Hyderabad population at 8% with male to female ratio of 1:3.²⁷

The limitation of this study is small sample size; further large scale studies are suggested. We did not investigate etiological causes of apertognathia, nor did we observe whether the apertognathia was skeletal or dental in origin. We will focus on these aspects in the future studies.

CONCLUSION

It was concluded that Apertognathia is more common among females than males in preadolescents. The high frequency of Apertognathia i.e. 10% in preadolescents

suggests that there is need to focus on preventive and interceptive orthodontics.

Conflict of Interest: The study has no conflict of interest to declare by any author.

REFERENCES

1. Ng CS, Wong WK, Hagg U. Orthodontic treatment of anterior open bite. *Int J Paediatr Dent* 2008; 18(2):78-83.
2. Huang GJ. Giving back to our specialty: Participate in the national anterior open-bite study. *American journal of orthodontics and dentofacial orthopedics: official publication of the Ame Associ Orthod, its constituent societies, and the American Board of Orthodontics* 2016;149(1):4.
3. Ngan P, Fields HW. Open bite: a review of etiology and management. *Pediatr Dent* 1996; 19(2):91-8.
4. Kim P, Sarauw MT, Sonnesen L. Cervical vertebral column morphology and head posture in preorthodontic patients with anterior open bite. *Am Orthod and Dentofacial Orthopedics* 2014; 145(3):359-66.
5. de Menezes LM, Ritter DE, Locks A. Combining traditional techniques to correct anterior open bite and posterior crossbite. *Am Orthodontics and Dentofacial Orthopedics* 2013;143(3):412-20.
6. Woodside DG, Linder-Aronson S. Progressive increase in lower anterior face height and the use of posterior occlusal bite-block in its management. *Orthodontics: State of art, essence of the science* ed. St. Louis: Mosby Co. 1986:209.
7. Alcan T, Keles A, Erverdi N. The effects of a modified protraction headgear on maxilla. *American Journal of Ortho and Dentofacial Orthopedics* 2000;117(1):27-38.
8. Küçükkeles N, Acar A, Demirkaya AA, Evrenol B, Enacar A. Cephalometric evaluation of open bite treatment with NiTi arch wires and anterior elastics. *Am ortho and dentofacial orthopedics* 1999;116(5):555-62.
9. Kim YH. Anterior openbite and its treatment with multiloop edgewise archwire. *The Angle Orthod* 1987;57(4):290-321.
10. Park YC, Lee HA, Choi NC, Kim DH. Open bite correction by intrusion of posterior teeth with miniscrews. *The Angle Ortho* 2008;78(4):699-710.
11. Ize-Iyamu IN, Isiekwe MC. Prevalence and factors associated with anterior open bite in 2 to 5 year old children in Benin city, Nigeria. *Afri Health Sci* 2012;12(4):446-51.
12. Ireland AJ, Mc Donald F. *Treatment and biomechanics* oxford, Great Britain: Oxford university press; 2003.p. 52.
13. Maciel CT, Leite IC. Etiological aspects of anterior open bite and its implications to the oral functions.

- Pro-Fono Revista de Atualizacao Cientifica 2005; 17(3):293-302.
14. Chung CJ, Hwang S, Choi YJ, Kim KH. Treatment of skeletal open-bite malocclusion with lymphangioma of the tongue. *Am Ortho and Dentofacial Orthopedics* 2012;141(5):627-40.
 15. Mizrahie E.A review of anterior open bite. *Br J Orthod* 1978; 5:21-27.
 16. Choi YJ, Kim DJ, Nam J, Chung CJ, Kim KH. Cephalometric configuration of the occlusal plane in patients with anterior open bite. *Am J Orthodont and Dentofacial Orthoped* 2016;149(3):391-400.
 17. Burford DA, Noar JH. The causes, diagnosis and treatment of anterior open bite. *Dental update* 2003;30(5):235-41.
 18. Farsi NM, Salama FS, Pedro C. Sucking habits in Saudi children: prevalence, contributing factors and effects on the primary dentition. *Pediatr Dent* 1997;19:28-33.
 19. Muviz BR. Epidemiology of malocclusion in Argentine children. *Commun dent and oral Epidemiol* 1986;14(4):221-4.
 20. Batarangaya A. Asurvey of Occlusal traits in an adolescent population in Uganda [Thesis]. Cape town university of western cape South Africa. 2004.
 21. Lauc T. Orofacial analysis on the Adriatic islands: an epidemiological study of malocclusions on Hvar Island. *The Eur Orthod* 2003;25(3):273-8.
 22. Behbehani F, Årtun J, Al-Jame B, Kerosuo H. Prevalence and severity of malocclusion in adolescent Kuwaitis. *Medical Principles and Practice* 2005;14(6):390-5.
 23. Jones ML, Mourino AP, Bowden TA. Evaluation of occlusion, trauma, and dental anomalies in African-American children of metropolitan Headstart programs. *J Clin Pediatr Dent* 1992; 18(1):51-4.
 24. Carvalho AC, Paiva SM, Scarpelli AC, Viegas CM, Ferreira FM, Pordeus IA. Prevalence of malocclusion in primary dentition in a population-based sample of Brazilian preschool children. *Eur Paediatr Denti* 2011;12(2):107-11.
 25. Hassanali J, Pokhariyal GP. Anterior tooth relations in Kenyan Africans. *Arch Oral Biol* 1993 1;38(4):337-42.
 26. Larsson E. The prevalence and aetiology of prolonged dummy and finger-sucking habits. *The Eur J Orthodont* 1985 Aug 1;7(3):172-6.
 27. Ahmed K, Shah R, un Nissa Q. frequency of anterior open bite in preadolescent patients reporting for orthodontic treatment. *Pak Orthodony* 2010;2(1).