

# Analyzing the Reduction of Symptoms in Dacryocystorhinostomy Patients With and Without Silicon Intubation A Study Using Randomized Controlled Trials

Silicon Intubation VS  
Dacryocystorhinostomy  
in Treating  
Nasolacrimal Duct  
Blockage

Nazullah<sup>1</sup>, Nuzat<sup>1</sup>, Samina Karim<sup>2</sup>, Mohammad Israr<sup>2</sup> and Romaisa<sup>2</sup>

## ABSTRACT

**Objective:** To assess the efficacy of silicon intubation vs dacryocystorhinostomy in treating nasolacrimal duct blockage

**Study Design:** A Randomized Controlled Trial Study

**Place and Duration of Study:** This study was conducted at the Department of Clinical Ophthalmology, Khyber Girls's medical College, Hayatabad Medical Complex (HMC), Peshawar from 09<sup>th</sup> October 2022 to 9<sup>th</sup> April 2023.

**Methods:** 446 individuals with nasolacrimal duct obstruction (NLDO) were included in the research. Every patient was divided into two groups. Dacryocystorhinostomy (DCR) with silicon intubation was performed on Group A, whereas DCR without intubation was performed on Group B.

**Results:** The sample as a whole was 35.1 + 9.2 years old on average. Group A's mean age was 34.9 + 9.3 years, whereas Group B's mean age was 35.2 + 9.1 years (p 0.730). Males made up 61.9% of group A and 56.5% of group B, respectively (p 0.248). In group A, the mean duration of symptoms was 11 + 3.2 days, whereas in group B, it was 10.4 + 2.9 days (p 0.082). Upon follow-up, group A's effectiveness (measured by the total remission of symptoms) was 79.4%, whereas group B's was 69.5% (p 0.017).

**Conclusion:** When compared to DCR without intubation, silicon intubation greatly increases the effectiveness of DCR in individuals with NLDO.

**Key Words:** Nasolacrimal duct obstruction, dacryocystorhinostomy, intubation, silicon, and efficacy.

**Citation of article:** Nazullah, Nuzat, Karim S, Israr M, Romaisa. Analyzing the Reduction of Symptoms in Dacryocystorhinostomy Patients With and Without Silicon Intubation A Study Using Randomized Controlled Trials. Med Forum 2024;35(3):75-78.doi:10.60110/medforum.350317.

## INTRODUCTION

Total resistance to lacrimal irrigation with 100% regurgitation from the same or opposite punctum or a lacrimal sac mucocele without later reasons was characterized as primary acquired nasolacrimal duct obstruction (PANDO)<sup>1</sup>. There have been many surgical techniques published since Cadweli's proposal of endonasal dacryocystorhinostomy (DCR) in 1893, including MMED and laser endoscopic DCR<sup>2</sup>. Regardless of surgical approach, common reasons for DCR failure include synechial adhesion with middle

turbinate and/or nasal septum and expanding cicatricial closure for secondary healing with/without granuloma development<sup>3</sup>. There were several operational aides for endonasal endoscopic mechanical DCR. Examples of medical interventions included canalicular stenting, intraoperative or postoperative mitomycin C, and absorbable or non-absorbable materials packed with or without medication, such as topical steroids. Ophthalmologists began to favor DCR with silicone intubation in the 1970s. They recommended it because ostium's opening preservation improved surgical patency. Previous investigations have connected silicone stent failure to granulomatous inflammation. Recent research discusses DCR surgery with 4 silicone intubation<sup>4</sup>. The most popular technique for preventing rhinostomy closure is the use of silicone stents. By preserving fistula patency and postponing fibrous closure after healing, silicone intubation may enhance the results of endoscopic DCR<sup>5</sup>. Silicone stenting during endoscopic DCR is still up for discussion, however. According to some studies, the silicone stent may induce tissue granulation, which raises the possibility of adhesions, postoperative infections, punctal lacerations, and surgical failure<sup>6</sup>. There were conflicting findings from two meta-analyses on silicone intubation during endoscopic DCR<sup>5</sup>. Research has

<sup>1</sup>. Department of Ophthalmology, Khyber Institute of Ophthalmic Medical Sciences, Peshawar.

<sup>2</sup>. Department of Ophthalmology, Hayatabad Medical Complex, Peshawar.

Correspondence: Samina Karim, TMO Department of Ophthalmology, Khyber Institute of Ophthalmic Medical Sciences, Peshawar.

Contact No: 033351616042

Email: drsaminakarim@yahoo.com

Received: July, 2023

Accepted: September, 2023

Printed: March, 2024

contrasted endoscopic DCR using silicone intubation with not using it<sup>7-12</sup>. A previous research found DCR with silicon intubation efficient in 93.3% and failed in 6.7%. In another study, 90.3% of endoscopic endonasal DCR treatments were effective<sup>9</sup>. Silicone intubation increased success to 93.7% from 86.7% without it. The present study compares DCR success rates in our community's NDO patients with and without silicon intubation. We were prompted to perform this study after experiencing patient attrition with NDO and DCR failures, whether intubated or not<sup>9</sup>. The literature is vast, however, several studies were undertaken with small sample numbers and had inconsistent and equivocal findings. If silicon intubation is equally or more successful than not using it, we will discuss the study's results and urge local ophthalmologists to further research and regular use of it during DCR for NDO. This research will show local DCR success rates with and without intubation for NDO<sup>10</sup>.

## METHODS

Hospital ethics and scientific committee authorized the study. The OPD department included all NDO patients (per operational criteria) in the study. Every patient supplied written informed permission after being informed of the research's goals and advantages. All patients had medical histories and ophthalmologic examinations. Block randomization divided patients into two groups. DCR patients in Group A received canalicular silicone stenting or intubation. Because Group B patients had DCR without silicon intubation, the silicone stent was inserted by both puncta and knotted in the nasal cavity to relieve canthal strain. One skilled CPSP fellow ophthalmologist performed all surgeries. After four weeks, all patients were evaluated for symptom relief and saline injection-confirmed duct patency. 58 Premade proformas documented all the aforementioned. Research bias and confounders were eliminated using a stringent exclusion procedure. The following are inclusion criteria: All 18–50-year-olds with main acquired nasolacrimal duct blockage, either gender. Congenital dacryocystitis and presacral occlusion such as canalicular blockage and punctual stenosis are excluded. • Atrophic rhinitis, chronic granulomatous disorders, and nasal tumors may affect surgery outcomes. • Previous lacrimal surgery failures. Epiphora after radiation/trauma. Confounders like this may affect research outcomes.

## RESULTS

The study comprised 446 NLDO patients. Split all patients in two. Group A had silicon-intubated DCR, but Group B did not. Each group had 223 patients. The age distribution is in Table 1. The sample averaged 35.1 + 9.2 years. Group A averaged 34.9 + 9.3 years, whereas Group B averaged 35.2 + 9.1 years (p 0.730). Gender distribution is in Table 2. Group A had more

men than B, but B had more women. The mean symptom duration was 11 + 3.2 days in group A and 10.4 + 2.9 days in group B (p 0.082). See Table 3. Table 4 shows follow-up efficacy (symptom resolution). Group A had 79.4% DCR symptom remission following intubation, whereas group B had 69.5% (p 0.017).

**Table No. 1: Comparison Of Age Between Both Groups (N=223 Each)**

Age	DCR with silicon intubation No %		DCR without silicon intubation No %	
20-30 years	76	34.1%	74	33.2%
30-40 years	88	39.1%	86	39.0%
40-50 years	59	26.1%	62	27.8%
Total	223		223	

NO= number, % percentage, p value= 0.948

**Table No. 2: Comparison Of Gender Between Both Groups (N=223 Each)**

Gender	DCR with silicon intubation No %		DCR without silicon intubation No %	
Male	138	61.9%	126	56.5%
Female	85	38.1%	97	43.5%
Total	223	100%	223	100%

NO= number, % percentage, p value=0.248

**Table No. 3: Comparison Of Duration Of Nldo Between Both Groups (N=223 Each)**

Duration of NLDO	DCR with silicon intubation No %		DCR without silicon intubation No %	
5-10 days	77	34.5%	134	60.1%
10-15 days	146	65.5%	89	39.1%
Total	223	100%	223	100%

NO= number, % percentage, p value=0.01

**Table No. 4: Comparison Of Efficacy Between Both Groups (N=223 Each)**

Efficacy	DCR with silicon intubation No %		DCR without silicon intubation No %	
Yes	177	79.4%	155	69.5%
NO	46	20.6%	68	30.5%
Total	223	100%	223	100%

NO= number, % percentage, p value=0.017

## DISCUSSION

Cryocystorhinostomy is the most frequent treatment for persistent dacryostenosis or nasolacrimal duct obstruction. Surgical DCR creates drainage between the nasal cavity and lacrimal sac<sup>11</sup>. LA-DCR, EN-DCR, and EX-DCR are the three DCR methods. In the 1970s, ophthalmologists preferred silicone intubated DCR<sup>11</sup>. They recommended its use and noticed that ostium opening preservation increased postoperative patency. Granulomatous inflammation increases silicone stent failure risk, according to prior studies<sup>12</sup>. Recent work

discusses silicone intubation during DCR surgery from different views. This study examined DCR success with and without stents and compared it to previous studies<sup>13</sup>. In a 2011 meta-analysis of DCR for nasolacrimal duct blockage with and without silicone tubes, DCR had equal success rates<sup>14</sup>. The meta-analysis found no benefit to silicone stent intubation for major DCR. Since 2010, more prospective comparison studies have shown that silicone intubation in primary DCR increased the success rate of DCR without intubation by 68 percent, even if these benefits were not statistically significant. In a large randomized controlled trial,<sup>15</sup> found that silicone intubation prevented the ostium from sealing, increasing DCR success. For effect size (population proportions 0.892 versus 0.943), sample size (111 and 105), and alpha (0.05, 2-tailed), the previous meta-analysis had four RCTs with a power of 0.274<sup>16</sup>. How silicone intubation would work during DCR surgery remained unknown. In the EX-DCR subgroup, DCR with intubation had a considerably greater success rate after surgery than DCR without intubation, according to this cumulative meta-analysis. Statistically significant change [RR, 1.06; 95%CI (1.02–1.11),  $p = 0.006$ ]<sup>17</sup>. The finding varied considerably from the previous meta-analysis. The preceding meta-analysis's low statistical power and few trials may explain the discrepancies. If the research has a negative outcome, consider its power<sup>18</sup>. If not, researchers risk type II errors and abandoning promising medicines. "Meta-analysis" combines the data of numerous "combinable." independent studies. Weak included studies enhance statistical power, decrease random error, and increase sample size<sup>19</sup>. Intranasal tissue granulation, adhesion, infection, hemorrhage, punctural or canalicular laceration, tube displacement or loss, and conjunctival irritation were common after surgery<sup>20</sup>. These concerns were linked to silicone tubes. The silicone tube may induce tissue granulation, a topic of controversy. Silicone intubation, an inorganic foreign material, may cause granulation and rhinostomy closure. Longari et al. found that stents decreased ostial size more. Turbinoseptal synaechia<sup>21</sup>, peristomal granuloma, and scar tissue produced most of this. The success rate for silicon stenting was 79.4% in 223 cases. Due to the endoscope, EnDCR has replaced the external DCR. Understanding the lateral nasal wall's anatomy and changes is crucial<sup>22</sup>. The orbicularis oculi muscle's pumping action may explain EnDCR's strong functional results. Silicon tubing prevents stoma fibrous closure, keeping the fistula open after surgical recovery. After EnDCR<sup>23</sup>, patients who retained silicon tubes outperformed those who extruded them. In a recent study, silicon stent recipients had a 79.4% success rate, whereas non-stent recipients had 69.5%. According to another study, the stent group had a greater Epiphora resolution rate, although it was not statistically significant. Kakkar et al.<sup>19</sup> and Unlu et al.<sup>20</sup> found that

silicon tubes in children cause complications, but they found no significant differences between silicon stent DCR and standard DCR. Silicon stents greatly increase primary DCR failure, according to a retrospective study<sup>24</sup>. They advised against silicon stent placement unless there is a particular cerebral obstruction. Due to granulation tissue formation, silicon stents in DCR prevent osteotomy and common canalicular occlusion. The 91.3% effectiveness rate of this strategy matches Elmorsy et al. Study<sup>23</sup>.

## CONCLUSION

DCR with silicon intubation is significantly effective in patients with NLDO compared to DCR without intubation.

**Recommendations:** More randomized controlled studies with bigger sample sizes and multicenters are needed to establish the best local data to justify silicon intubation during DCR in NLDO patients.

**Acknowledgment:** We would like to thank the hospital administration and everyone who helped us complete this study.

### Author's Contribution:

Concept & Design of Study:	Nazullah
Drafting:	Nuzat, Samina Karim
Data Analysis:	Mohammad Israr, Romaisha
Revisiting Critically:	Nazullah, Nuzat
Final Approval of version:	Nazullah

**Conflict of Interest:** The study has no conflict of interest to declare by any author.

**Source of Funding:** None

**Ethical Approval:** No.HMC-CARD-F-1 and approval No.0051 Dated 25.11.2021

## REFERENCES

1. Massoud OM, Elbakary MA, Shalaby OE, Hashish AA. Evaluation of the role of silicone intubation in non-complicated external dacryocystorhinostomy. *The Open Ophthalmol J* 2020;14(1).
2. Nitin T, Uddin S, Paul G. Endonasal Endoscopic Dacryocystorhinostomy with and Without Stents—A Comparative Study. *Ind J Otolaryngol Head Neck Surg* 2022;74(Suppl 2):1433-41.
3. Yu B, Xia Y, Sun JY, Ye Q, Tu YH, Zhou GM, Wu WC. Surgical outcomes in acute dacryocystitis patients undergoing endonasal endoscopic dacryocystorhinostomy with or without silicone tube intubation. *Int J Ophthalmol* 2021;14(6):844.
4. Matoušek P, Lubojacký J, Masárová M, Čábalová L, Červenka S, Komínek P. Does Bicanalicular Intubation Improve the Outcome of Endoscopic Dacryocystorhinostomy? *J Clin Med* 2022;11(18):5387.

5. Orlova EI, Davydov DV. Lacrimal stents in the lacrimal pathways' surgery. *Ophthalmol J* 2017;10(4):48-55.
6. Lapusneanu L, Neacsu A. Method of stenting the lacrimal sac after performing dacryocystorhinostomy. *Romanian J Rhinol* 2019;9(34):101-4.
7. Buttanri IB, Serin D. Silicone intubation indications in external dacryocystorhinostomy. *Med Hypothesis Discov Innov Ophthalmol* 2014;3(4):101-2.
8. Feng YF, Cai JQ, Zhang JY, Han XH. A meta-analysis of primary dacryocystorhinostomy with and without silicone intubation. *Can J Ophthalmol* 2011;46(6):521-7.
9. Gu Z, Cao Z. Silicone intubation and endoscopic dacryocystorhinostomy: a meta-analysis. *J Otolaryngol Head Neck Surg* 2010;39(6):710-3.
10. Syed MI, Head EJ, Madurska M, Hendry J, Eriktila OC, Cain AJ. Endoscopic primary dacryocystorhinostomy: are silicone tubes needed? Our experience in sixty-three patients. *Clin Otolaryngol* 2013;38(5):406-10.
11. Knisely A, Harvey R, Sacks R. Long-term outcomes in endoscopic dacryocystorhinostomy. *Curr Opin Otolaryngol Head Neck Surg* 2015;23(1):53-8.
12. Saeed BM. Endoscopic DCR without stents: clinical guidelines and procedure. *Eur Arch Otorhinolaryngol* 2012;269(2):545-9.
13. Al-Qahtani AS. Primary endoscopic dacryocystorhinostomy with or without silicone tubing: a prospective randomized study. *Am J Rhinol Allergy* 2012;26(4):332-4.
14. Shashidhar K, Nagalotimath U, Dixit D. Endoscopic dacryocystorhinostomy with and without silicone stenting: a comparative study. *Al Ameen J Med Sci* 2014;7(3):244-247.
15. Kim SI, Jin HJ, Kim S, Hwang SH. The clinical efficacy of silicone stents for endoscopic dacryocystorhinostomy: a meta-analysis. *Clin Experimental Otorhinolaryngol* 2018;11(3):151.
16. Gurdeep S. Powered endoscopic dacryocystorhinostomy with mucosal flaps without stenting. *Med J Malaysia* 2008;63(3):237.
17. Acharya K, Pradhan B, Thapa N, Khanal S. Comparison of outcome following endoscopic dacryocystorhinostomy with external dacryocystorhinostomy. *Nepalese J ENT Head Neck Surg* 2012;2(2):2-3.
18. Feng YF, Cai JQ, Zhang JY, Han XH. A meta-analysis of primary dacryocystorhinostomy with and without silicone intubation. *Canadian J Ophthalmol / J Canadien d'Ophthalmologie* 2011;46(6):521-7.
19. Kakar V, Chugh J, Sachdeva S, Sharma N, Ramesh. Endoscopic dacryocystorhinostomy with and without silicone stent: A comparative study. *Int J Otorhinolaryngol* 2009; 9:1.
20. Unlu HH, Aslan A, Toprak B, Guler C. Comparison of surgical outcomes in primary endoscopic dacryocystorhinostomy with and without silicone 85 intubation. *Annals of Otology, Rhinol Laryngol* 2002;111(8):704-9.
21. Cavaliere M, De Luca P, Scarpa A, De Bonis E, Troisi D, Cassandro C, Concilio et al. Longitudinal randomized study to evaluate the long-term outcome of endoscopic primary dacryocystorhinostomy with or without silicone tube. *Eur Archives Oto-Rhino-Laryngol* 2022:1-5.
22. Orsolini MJ, Schellini SA, Souza Meneguim RL, Catâneo AJ. Success of endoscopic dacryocystorhinostomy with or without stents: systematic review and meta-analysis. *Orbit* 2020;39(4):258-65.
23. Elmorsy S, Fayek HM. Rubber tube versus silicone tube at the osteotomy site in external dacryocystorhinostomy. *Orbit* 2010;29(2):76-82.
24. Penttilä E, Smirnov G, Tuomilehto H, Kaarniranta K, Seppä J. Endoscopic dacryocystorhinostomy as a treatment for lower lacrimal pathway obstructions in adults. *Allergy Rhinol* 2015;6(1):ar-2015.