

Original Article

# To Study the Role of Serum Biliurbin and Lipoproteins in Prediction of Ischemic Heart Disease

1. Muhammad Binyameen 2. Murad Khan 3. Shariq Suhail Jafar 4. Gul-e-Raana 5. Asima Malik 6. Rukhshan Khurshid

1. Asstt. Prof. of Physiology, AIMC, Lahore 2. Assoc. Prof. of Biochemistry, CMH Lahore 3. Asstt. Prof. of Cardiology, AIMC, Lahore 4. Assoc. Prof. of Biochemistry, GMC, Gujranwala 5. Prof. of Biochemistry, FJMC, Lahore 6. Asstt. Prof. of Biochemistry, FJMC, Lahore

## ABSTRACT

**Background:** An inverse relationship between the bilirubin level and ischaemic heart disease was accounted.

**Study Design:** Cross Sectional Study.

**Place and Duration of Study:** This study was conducted at Dept. of Biochemistry, Fatima Jinnah Medical College, Lahore from Feb. 2007 to July 2007.

**Materials and Methods:** Study was performed to determine if serum bilirubin, when combined with various risk factors like lipid and lipoprotein predict ischemic heart disease (IHD).

**Results:** It is observed that the bilirubin is negatively correlated with hemoglobin, albumin, HDL and LDL-cholesterol. On the other hand a direct correlation with total lipid, cholesterol and ratio of cholesterol/HDL-cholesterol is also observed. High level of serum creatinine level was also found in the patients as compared to normal subject.

**Conclusion:** Our study observed not a definite relationship between serum bilirubin and IHD. Hence the relationships between bilirubin and lipoproteins (risk factors of IHD) require further clarification, although abnormal intermediary metabolism and antioxidant deficiency may be possible linking factors.

**Key Words:** Bilirubin, Lipoproteins and IHD.

## INTRODUCTION

Bilirubin and albumin may act as antioxidants. Their circulating levels are lower in those patients with ischemic heart disease (IHD) and could be further reduced by more extensive atherosclerosis, i.e. peripheral vascular disease<sup>1</sup>.

Circulating bilirubin is considered to protect human tissues from peroxidation of organic compounds, lipids and thus inhibiting foam cell formation in the arterial wall.<sup>2</sup> Low bilirubin was associated with several cardiovascular risk factors, in particular vascular aging, smoking, low concentrations of high-density lipoprotein cholesterol, low forced expiratory volume and low serum albumin<sup>3</sup>.

Low bilirubin level in blood may prove to be a significant marker for the evaluation of the general antioxidant status of the human organism. Mechanisms that give rise to this phenomenon are probably diverse and not well-studied yet. Processes of the formation of free oxygen and peroxide radicals are known to take place in numerous pathological conditions<sup>4</sup>.

The effects of serum bilirubin on blood lipids and lipoproteins in a number of patients were investigated by a group of workers<sup>5</sup>. The findings were that total cholesterol (TC) and low-density lipoprotein-cholesterol (LDL-C) were negatively correlated with total bilirubin and direct bilirubin. Their results indicate that bilirubin affects metabolism of lipoproteins and the low level of serum bilirubin is a new risk factor for coronary heart disease. This parameter may, alone or in

combination with other factors, make it possible to distinguish individuals with a risk of CAD<sup>6</sup>.

It was found<sup>7</sup> that elevated serum creatinine has been associated with increased mortality in hypertensive persons, the elderly, and patients with myocardial infarction or stroke in whom cardiovascular disease is the major cause of death.

Present study tried to find out the role of serum bilirubin in prediction of ischemic heart disease when combined with risk factors like lipid and lipoprotein.

## MATERIALS AND METHODS

Hundred patients with IHD were included in the study. Patients were taken from in and out department of Medicine. The traditional risk factors like cholesterol, high-density lipoprotein cholesterol (HDL-C), cholesterol/HDL-C ratios, triglycerides as well as serum bilirubin, albumin and creatinine were determined by standard methods. Besides, age, cigarette smoking, and systolic blood pressure were also recorded.

## RESULTS

Table 1 shows that the mean age of patients was 46 years. Among these 50% smokers and their mean blood pressure were 100/65mmHg.

Assessment of biochemical parameters was tabulated (Table 2). Level of bilirubin was increased in patients as compared to normal subjects but this shows no significant difference. In case of creatinine and albumin, level of creatinine was significantly ( $P<0.01$ )

increased in patients whereas the level of albumin was significantly decreased ( $P<0.01$ ) in patients as compared to the normal subjects. Among lipid profile, It was observed that the levels of serum cholesterol and total lipid were increased in patients. A highly significant difference ( $P<0.01$ ) was observed in case of serum cholesterol. Level of HDL-chol, LDL-chol and triglyceride were decreased in patients as compared to control subjects but significant difference ( $P<0.01$ ) was only observed in case of HDL-cholesterol. On the other hand ratio of cholesterol/HDL-cholesterol was increased in patients as compared to control subjects.

**Table No.1: Variation in physical Parameters**

Parameters	
Mean age	46.7 $\pm$ 2.17
Smoking	50% smokers
Mean Blood Pressure(mm Hg)	100/65 $\pm$ 3.4/1.6

**Table No.2: Variation in bilirubin and other biochemical parameters in serum of Patients and their controls (No of cases in parenthesis and values are in mean $\pm$ s.e.m)**

Parameters	Patients (100)	Control subjects (25)
Bilirubin (mg/dl)	0.94 $\pm$ 0.06	0.42 $\pm$ 0.04
Creatinine (mg/dl)	1.9 $\pm$ 0.81**	0.78 $\pm$ 0.03
Albumin (gm/dl)	3.3 $\pm$ 0.08**	4.65 $\pm$ 0.02
Cholesterol (mg/dl)	254.6 $\pm$ 14.8**	185.0 $\pm$ 5.9
HDL-chol (mg/dl)	37.56 $\pm$ 0.53**	53.56 $\pm$ 0.59
LDL-chol (mg/dl)	130.56 $\pm$ 7.31	141.87 $\pm$ 5.52
Ratio of chol/HDL-chol	6.77	5.38
Triglyceride (mg/dl)	113.5 $\pm$ 4.6	125.25 $\pm$ 12.83

\*\* $P<0.01$ =Highly significant difference

## DISCUSSION

Many risk factors for IHD have been identified. Recently an association between low concentration of serum bilirubin and increased risk of IHD has been reported. It was also observed that bilirubin may affect on the lipoproteins. However, information on this topic remains scarce.

Results of present study indicate that the bilirubin is negatively correlated with hemoglobin, albumin, HDL and LDL-cholesterol. A study observed significantly higher total cholesterol and triglyceride levels and lower high-density lipoprotein cholesterol (HDL-C) levels in the AMI patients. The bilirubin and albumin were lower in AMI patients than controls<sup>8</sup>. Another study provide substantial evidence that serum bilirubin may play a protective role in peripheral arterial disease as well as ischaemic heart disease<sup>9</sup>. It was reported<sup>10</sup> that bilirubin is a potent antioxidant generated intracellularly during the degradation of heme by enzyme heme-oxygenase. Increased heme oxygenase

activity was associated with enhanced tissue bilirubin tissue bilirubin content land can increased rate of bilirubin release. Bilirubin may provide cardio protection against reperfusion injury.

On the other hand a direct correlation with total lipid, cholesterol and ratio of cholesterol/HDL-cholesterol is also observed. It was suggested<sup>11</sup> that serum bilirubin may be combined with LDL-C/HDL-C ratios, cholesterol/HDL-C ratios, cholesterol, or with various apolipoproteins to improve the prediction of CAD. Another group<sup>12</sup> found that bilirubin affects metabolism of lipoproteins and the low level of serum bilirubin is a new risk factor for coronary heart disease.

High level of serum creatinine level was found in patients as compared to normal subject. Same was reported by a group of workers<sup>7</sup>. They suggested a direct relationship of serum creatinine with IHD and it shows a role of renal impairment in the disease. Higher levels of HDL cholesterol were associated with a significant decrease in risk of nonfatal stroke. In contrast, elevated total cholesterol showed a weak positive association with nonfatal strokes. The marked inverse association between HDL cholesterol and stroke seen in hypertensives emphasizes the importance of those modifiable risk factors for stroke known to lower the concentrations of HDL cholesterol<sup>13</sup>.

## CONCLUSION

Present study found no relationship between serum bilirubin and IHD. Hence the relationships between bilirubin require further clarification, although abnormal intermediary metabolism and antioxidant deficiency may be possible linking factors.

## REFERENCE

1. Krijgsman B, Papadakis JA, Ganotakis ES, Mikhailidis DP, Hamilton G. The effect of peripheral vascular disease on the serum levels of natural anti-oxidants: bilirubin and albumin. *Int Angiol* 2002;21(1):44-52.
2. Icsó J, Nanova D. IHD and bilirubin level in a group of hospitalized patients. *Vnitr Lek* 1997, 43: 801-3
3. Breimer LH, Wannamethee G, Ebrahim S, Shaper AG. Serum bilirubin and risk of IHD in middle aged British men. *Clin Chem* 1995, 28:557-63.
4. Lingenhel A, Kollerits B, Schwaiger JP, Hunt SC, Gress R, Hopkins PN, et al. Serum bilirubin levels, UGT1A1 polymorphisms and risk for coronary artery disease. *Exp Gerontol* 2008;43(12):1102-7.
5. Li Y, Zaho S, Effect of serum bilirubin on lipoproteins. *Human I Ko To Hsueh Hsueh Pao* 1998; 23:578-80.
6. Shcherbinina MB[Low blood bilirubin level: possible diagnostic and prognostic importance] *Klin Med (Mosk)* 2007;85(10):10-4.
7. Wannamethee SG, Shaper AG, Penny IJ. Serum

creatinine concentration and risk of cardiovascular disease: a possible marker for increased risk of stroke. *Stroke* 1997;28:557-63.

8. Kumar A, Nagtilak S, Sivakanesan R, Gunasekera S. Cardiovascular risk factors in elderly normolipidemic acute myocardial infarct patients--a case controlled study from India. *Southeast Asian J Trop Med Public Health* 2009;40(3):581-92.

9. Breimer LH, Mikhailidis DP. Bilirubin and peripheral arterial disease: 15 years later. *Expert Opin Ther Targets* 2009;13(2):139-40.

10. Clark JE, Foresti R, Sarathchandra P, Kaur H, Green CJ, Motterlini R. Heme oxygenase-derived bilirubin ameliorates post-Ischemic myocardial dysfunction. *Am J Physiol Heart Circ Physiol* 2000; 278(2):H643-51

11. Schwertner HA, Fischer JR. Comparison of various lipids and bilirubin concentration as risk factors for predicting CAD. *Atherosclerosis* 2000;150:381-7.

12. Ko GT, Chan JC, Woo, J Lau E. Serum bilirubin and cardiovascular risk factors in a Chinese population. *J Cardiovascualr Risk* 1996;3:459-63.

13. Wannamethee SG, Shaper AG, Ebrahim S. HDL-Cholesterol, total cholesterol, and the risk of stroke in middle-aged British men. *Stroke* 2000; 31(8): 1882-8.

**Address for Corresponding Author:**

**Dr. Murad Khan,**  
Assoc. Prof. of Biochemistry,  
CMH Lahore  
Cell No.0300-4682235