

Vol. 35, No. 1 January, 2024

ISSN 1029 - 385 X (Print)

ISSN 2519 - 7134 (Online)



MEDICAL FORUM MONTHLY

RECOGNISED BY
PMC & HEC

APNS
Member

CPNE
Member

ABC
Certified

Open Access Journal

Journal of all Specialities


“Medical Forum” Monthly Recognised and Index by

- ☞ PMDC with Index Pakistan No.48 since 1998
- ☞ HEC since 2009
- ☞ Pakmedinet Since 2011
- ☞ Medlip (CPSP) Since 2000
- ☞ PASTIC & PSA Since 2000
- ☞ NLP Since 2000
- ☞ WHO, Index Medicus (IMEMR) Since 1997
- ☞ EXCERPTA MEDICA, Netherlands Since 2000
- ☞ EMBASE SCOPUS Database Since 2008
- ☞ Registered with International Standard Serial Number of France bearing ISSN 1029-385X (Print), ISSN 2519-7134 (Online) Since 1992
- ☞ Registered with Press Registrar Govt. of Pak bearing No.1221-B Copr. Since 2009
- ☞ ABC Certification Since 1992
- ☞ On Central Media List Since 1995
- ☞ Med. Forum Published under Medical Academic Foundation (MAF) from Lahore Since 1989
- ☞ Open Access, Peer Review & Online Journal
- ☞ Email: med_forum@hotmail.com, medicalforum@gmail.com
- ☞ website: www.medforum.pk

<h1>MEDICAL FORUM MONTHLY</h1>	ISSN 1029 - 385 X (Print)	ISSN 2519 - 7134 (Online)	
	APNS Member	CPNE Member	ABC Certified
	Peer Review Journal	Online Journal	Published Since 1989
	e-journal available on: www.medforum.pk		

Medical Forum Recognized and Indexed by

PMDC-IP-0048 (1998), HEC-Y-Category (2009), Pastic and PSA, Isd (2000), Medlip, Karachi (2000), NLP, Isd (2000), Pakmedinet, Isd (2011), Excerpta Medica, Netherlands (2000), EMBASE Scopus Database (2008), Index Medicus (IMEMR) WHO (1997), ABC Certification, Govt. of Pak. (1992), Central Media list, Govt. of Pak (1995), Press Reg. No.1221-B Copr (2009)

Published under Reg.No.RP/11256/L/S/18
Medical Academic Foundation


Editorial Executives

Patron-in-Chief Prof. Mahmood Ali Malik Medicine	Editor-in-Chief Prof. Azhar Masud Bhatti Public Health Specialist & Nutritionist	Managing Editor Prof. Nasreen Azhar Consultant Gynaecologist
Co-Editors Tahir Masud Jan (Canada) Dr. Meshaal Azhar (Pak) Dr. Faryal Azhar (Pak)	Editor Dr. Mohsin Masud Jan	Associate Editors Prof. Syed Mudassar Hussain (Pak) Prof. M. Mohsin Khan (Pak) Dr. Iftikhar A. Zahid (Pak)

National Editorial Advisory Board

Prof. Abdul Hamid	Forensic Medicine	Sialkot	03239824782	drabdulhamid12345@hotmail.com
Prof. Abdul Khaliq Naveed	Biochemistry	Rawalpindi	03215051950	khaliqnaveed2001@yahoo.com
Prof. AftabMohsin	Medicine	Lahore	03314101516	aftabmohsin@yahoo.com
Prof. Anjum Habib Vohra	Neurosurgery	Lahore	03008443218	omer@brain.net.pk
Prof. Asad Aslam Khan	Ophthalmology	Lahore	03008456377	drasad@lhr.comsats.net.pk
Prof. Haroon Khurshid Pasha	Paed. Surgery	Multan	03008633433	haroonkpasha@hotmail.com
Prof. Haroon Nabi	Dermatology	Lahore	03004000216	haroonnabi@hotmail.com
Prof. Javed Akram	Medicine	Lahore	03008450505	vc@uhs.edu.pk
Prof. Kh. M. Azeem	Surgery	Lahore	03334242122	khawaja.azeem@sihs.org.pk
Prof. Khalid Masood Gondal	Surgery	Lahore	03328483823	rc_lahore@cpsp.edu.pk
Prof. M. Amjad	ENT	Lahore	03334254695	professoramjad@yahoo.com
Prof. M. Amjad Amin	Surgery	Multan	03336103262	dramjadamin@gmail.com
Prof. M. Sabir	Anatomy	Sialkot	03005183021	raosabirdr62@gmail.com
Prof. Mahmood Nasir Malik	Medicine	Lahore	03009487434	nasirphysician@yahoo.com
Prof. Majeed Ahmad Ch.	Surgery	Lahore	03008440415	prof_abdulmajeed@hotmail.com
Prof. Mian Rasheed	Forensic Medicine	Rawalpindi	03025033559	drmian1000@hotmail.com
Prof. Pervez Akhtar Rana	Forensic Medicine	Lahore	03009422511	pzrana@gmail.com
Prof. Rukhsana Majeed	Community Medicine	Quetta	03337808138	majidrukhsana@hotmail.com

Prof. Safdar Ali Shah	Urology	Lahore	03334391474	drsafdar-ali@hotmail.com
Prof. SardarFakhar Imam	Medicine	Lahore	03008451843	drfakhar@lhr.paknet.com.pk
Prof. Shahid Mehmood	Surgery	Rawalpindi	03215001120	shahid63@gmail.com
Prof. Syed M. Awais	Orthopaedics	Lahore	03334348716	awais@kemu.edu.pk
Prof. Syed Nazim Hussain Bukhari	Medicine & Chest	Lahore	03009460515	nhbokhari@yahoo.com
Prof. Zafarullah Ch.	Surgery	Lahore	03072222533	administrator@csp.edu.pk

International Editorial Advisory Board

Dr. Amjad Shad	Neurosurgery	UK	447963442419	amjad.shad@uhcw.nhs.uk
Dr. Ghazanfar Ali	Gastroenterology	UK	447800760008	ghazanfarali@hotmail.com
Dr. Haider Abbas	Urology	UK	447816149374	haidersyed@hotmail.com
Dr. Khalid Rashid	Cardiology	UK	447740477756	khalid.rashid@cht.nhs.uk
Dr. M. Shoaib Khan	Medicine	UAE	00971503111420	msksd2000@yahoo.com
Dr. Basil Nouman Hashmi	Surgery	UK	00447806611517	basilhashmi@doctor.net.uk
Dr. Sohail Saied	Surgery	UK	00441923285114	sohailsaied@gmail.com
Dr. Safdar Ali	Cardiology	USA	0016307816668	safdarali@sbcglobal.net
Dr. Parashu Ram Mishra	Surgery & Gastroenterology	Nepal	+9779841233450	drparashuram.mishra@gmail.com
Dr. Mansoor M. Mian	Psychiatry	USA	+1 (972)375 7821	mmian2000@yahoo.com
Dr. Sohail Qureshi	Orthopaedic	UK	00447734329666	quraishisohail@yahoo.com
Dr. Mushtaq Ahmad Mughal	Orthopaedics	UK	00447971886006	mahmed01@blueyonder.co.uk
Dr. Mansoor Tahir	Radiology	UK	00447921838093	drmansoortahir@yahoo.com

Business Manager: Nayyar Zia Ch.

Legal Advisors: Kh. EjazFeroz (Barrister),
Kh. Mazhar Hassan &Firdos Ayub Ch. (Advocates)

Published under: Medial Academic Foundation (MAF) Reg. No.RP/11256/L/S/18

Published By: Prof. Nasreen Azhar, Gohawa Road, Link Defence / New Airport Road,
Opposite Toyota Motors, Lahore Cantt. Lahore.
Mobile Nos. 0331-6361436, 0300-4879016, 0345-4221303, 0345-4221323.
E-mail: med_forum@hotmail.com, medicalforum@gmail.com
Website: www.medforum.pk

Printed By: Naqvi Brothers Printing Press, Darbar Market, Lahore.

Rate per Copy: Rs.3000.00

Subscription Rates : Pakistan (Rs.30000.00), USA & Canada (US\$ 500.00),
(annually) China, Japan, UK &Middle East (US\$ 450.00)

CONTENTS

Editorial

Migraine in Children and Adults _____	1-2
Moshin Masud Jan	

Original Articles

1. Frequency of Cardiac Arrest and Arrhythmias Associated with Admitted Patients in the Cardiac Care Unit in Hazara Division _____	3-7
1. Sardar Fawad Gul 2. Mohsin Khan 3. Mohammad Imran Khan 4. Zia Qamar 5. Adnan Haider 6. Zulqarnain Dilawar	
2. Dentists' and Dental Students' Perspectives on Amalgam Restoration In Saudi Arabia _____	8-13
1. Abdulaziz Alhumaid 2. Ahmed Alharbi 3. Abdulaziz Alrebdi 4. Ebrahim Alshawy 5. Nawaf Almutairi 6. Yasir Alyahya	
3. Effects of Intra-Cameral Dexamethasone after Uncomplicated Phacoemulsification _____	14-17
1. Muhammad Saad Ullah 2. Muhammad Usama Rahim 3. Hassan Shoaib	
4. Anatomical Variation in the Location of Mandibular Foramen with Age Using Cone Beam Computed Tomography _____	18-22
1. Asma Sattar 2. Naheed Imran 3. Muhammad Ishfaq 4. Sana Arbab 5. Munawar Aziz Khattak 6. Imran Khattak	
5. Evaluation of Obstetric Anaesthesia and its Association with Maternal Outcomes in Women with Placenta Previa: A Cross-Sectional Study _____	23-27
1. Shandana Bawar 2. Qudsia Qazi 3. Syeda Sitwat Fatima	
6. The Importance of Immunohistochemical Evaluation of Ki67 in Detecting Early Malignant Changes in Colorectal Adenomatous Polyps _____	28-32
1. Sabika Batool 2. Talat Mirza 3. Fouzia Lateef 4. Sobia Hassan	
7. Analysis of Fingerprint Patterns in Relation to ABO Blood Groups: A Comparative Study _____	33-36
1. Zulfiqar Ali Buzdar 2. Mansoor Mirza 3. Ambreen Serwer 4. Muhammad Anwar Sibtain Fazli 5. Faiza Munir	
8. Gender-Based Analysis of the Lambdoid Cranial Suture among Human Cadavers Presented for Postmortem Examination at King Edward Medical University, Lahore _____	37-41
1. Mansoor Mirza 2. Zulfiqar Ali Buzdar 3. Ambreen Serwer 4. Muhammad Anwar Sibtain Fazli 5. Faiza Munir	
9. Assessment of the Labial Alveolar Bone Thickness Overlying Maxillary Anterior Teeth in Different Age Groups, Genders, and Sides of the Arch: A Cone Beam Computed Tomographic Study _____	42-46
1. Naheed Imran 2. Asma Sattar 3. Imran Khattak 4. Sana Arbab 5. Munawar Aziz 6. Syed Amjad Shah	
10. Delayed Presentation of Large Goitre, A Cross Sectional Study at a Tertiary Care Hospital in Karachi _____	47-49
1. Mariam Imran 2. Saad Abdul Razzaq 3. Zahid Mehmood 4. Ghansham Rawtani 4. Hazrat Bilal Burki	
11. Percutaneous Nephrolithotomy: A Single Center Experience of 162 Cases of Standard Percutaneous Nephrolithotomy _____	50-53
1. Zohair 2. Akhtar Nawaz 3. Waqas 4. Siddique Akbar 5. Sardar Alam	
12. Complications of Ultrasound Guided Percutaneous Nephrostomy in Adults: A retrospective Study _____	54-57
1. Akhtar Nawaz 2. Waqas 3. Zohair 4. Siddique Akbar 5. Sardar Alam	
13. Comparison of Treatment Response of Different Drugs in Common Migraine _____	58-62
1. Salman Khan 2. Imran Ahmad 3. Muhammad Jalil 4. Ibrar Saleem 5. Fawad Ahmad 6. Khuram Haq Nawaz	
14. Effect of Informed Consent on Patient's Anxiety Regarding Third Molar Surgery _____	63-67
1. Kiran Bashir 2. Sadia Paiker 3. Erum Riaz 4. Wajiha Walayat 5. Syed Muhammad Zaki Mehdi 6. Fatima Khattak	

15. The Influence of Surgical Volume on Outcomes in Radical Cystectomy: A Population-Based Analysis _____	68-72
1. Shahjehan 2. Syed Ameroon Shah 3. Maham Arooj Zaidi 4. Khalid Islam 5. Muhammad Zohair 6. Naseeb Dad	
16. Frequency of Peptic Ulcer Disease in Patients with Chronic Use of Nonsteroidal Anti-Inflammatory Drugs (NSAIDs) _____	73-75
1. Shakeel Ahmad 2. Muhammad Naeem 3. Hamid Ullah 4. Liaqat Ali	
17. Comparison of the Skin Closures Using Staples Versus Prolene Sutures in Patients Undergoing Clean Elective Abdominal Surgeries _____	76-79
1. Shumaila Naseer 2. Tariq Hayat Khan 3. Ayaz Gul 4. Nida Mumtaz	
18. The Status of Hepatitis B Vaccination Among Hemodialysis Patients at a Tertiary Care Hospital _____	80-82
1. Shad Muhammad 2. Arbab Muhammad Ali 3. Muhammad Ikram	
19. Frequency of Inflammatory Bowel Disease in Patients who Underwent Colonoscopy for Lower Gastrointestinal Bleeding _____	83-86
1. Dilaram Khan 2. Inayat Ullah 3. Mohammad Sohail 4. Aamir Ghaffoor	
20. Developmental Profile as a Predictor of Behavior phenotype in Down Syndrome Children _____	87-91
1. M Bilal Abid 2. Syeda Wajeeha Zahra 3. Muneeba Kamran 4. Sultan Badar Munir 5. Ehsan Ullah 6. Munawar Ghous	
21. Chest Pain Management Using Prehospital Point-of-Care Troponin and Paramedic Risk Assessment _____	92-95
1. Feras Almarshad 2. Ghulam Mustafa	
22. Risk Factors for Epidural Anesthesia Blockade Failure in Cesarean Section _____	96-99
1. Abid Haleem Khattak 2. Muhammad Sheharyar Ashraf 3. Amjid Ali 4. Jawad Hameed 5. Samar Naeem 6. Kashaf Noor	
23. Antiemetic Prophylaxis with Droperidol in Morphine-Based Intravenous Patient Controlled Analgesia _____	100-103
1. Muhammad Sheharyar Ashraf 2. Abid Haleem Khattak 3. Jawad Hameed 4. Amjid Ali 5. Kashaf Noor 6. Samar Naeem	
24. Micro and Macrocytic Anemia – A Population Based Cross-Sectional Study _____	104-108
1. Ayesha Zaib-Un-Nisa 2. Iyad Naeem Muhammad 3. Sheikh Abdul Khaliq 4. Agha Umer Draz Khan	

Narrative Review

25. Interdisciplinary Collaboration in Pediatric Dentistry: Challenges and Opportunities – A Narrative Review _____	109-115
Mohammed Ali Habibullah	

Case Report

26. C1Q Nephropathy, An Unusual Occurrence in a Middle-Aged South Asian Woman _____	116-118
1. Zohaib Ramzan 2. Shahid Anwar 3. Syed Ali Raza 4. Zoha Majeed	
27. Author Index January to December 2022 _____	119-125
Azhar Masud Bhatti	
28. Subject Index January to December 2022 _____	126-147
Azhar Masud Bhatti	

Editorial

Migraine in Children and Adults

Mohsin Masud Jan

Editor

Migraine is a common problem worldwide with significant morbidity and economic impact.¹ The direct costs of migraine are directly related to the severity of migraine pain and disability, and rise dramatically with prescription medication usage.^{2,3} The indirect costs exceed the costs of medical care, however, and work-related disability is the most important determinant of the economic impact of migraine.⁴ Migraineurs often miss work (absenteeism) or have reduced productivity at work (presenteeism).

Migraine is a type of headache characterized by recurrent attacks of moderate to severe throbbing and pulsating pain on one side of the head. The pain is caused by the activation of nerve fibers within the wall of brain blood vessels traveling inside the meninges (three layers of membranes protecting the brain and spinal cord).

Headaches are very common in children and teens. In fact, more than half will suffer from headaches at some point, and by 18 years the majority of adolescents have had them. And while most headaches are part of a viral illness, some are migraines. In fact, recurring migraines affect as many as one in 10 children and teens overall.

Migraines sometimes occur even earlier. Before puberty, boys and girls are equally likely to have them.

After puberty, migraines are more common in girls.

Migraines are often one-sided in adults. In children they are more likely to be felt on both sides of the head, either in both temples or both sides of the forehead.

While it's not always easy to tell a migraine from another kind of headache, children often report throbbing pain may experience nausea and sensitivity to light and noise.

The flashing lights and other vision changes people often see as a migraine begins are less common in children. However, parents may notice that their child is more tired, irritable, or pale before a migraine begins and takes a while to get back to normal after it ends.

A number of different factors can increase your risk of having a migraine. These factors, which trigger the headache process, vary from person to person and include sudden changes in weather or environment, too much or not enough sleep, strong odors or fumes, emotion, stress, overexertion, loud or sudden noises, motion sickness, low blood sugar, skipped meals, tobacco, depression, anxiety, head trauma, hangover, some medications, hormonal changes and bright or flashing lights.

Epidemiologic data suggest that successful therapy of the most severely affected migraineurs may significantly impact the overall economic burden of migraine.⁵ Migraine therapy employs preventive and symptomatic measures with pharmacologic and non-pharmacologic treatments are often used in both strategies. With careful examination of headache diaries and lifestyle influences, approximately 50-75% of migraineurs are able to identify factors that provoke their headaches.^{6,7} Awareness and avoidance of specific migraine triggers are incorporated into the treatment strategy to decrease the frequency of migraine in a given individual. Triggers for migraine include various foods and beverages, stress or relief of stress, and hormonal factors (such as menstruation and pregnancy).^{6,8,9}

Migraine is divided into four phases, all of which may be present during the attack. First phase is premonitory symptoms occur up to 24 hours prior to developing a migraine. These include food cravings, unexplained mood changes (depression or euphoria), uncontrollable yawning, fluid retention, or increased urination.

Second phase is Aura. Some people will see flashing or bright lights or what looks like heat waves immediately prior to or during the migraine, while others may experience muscle weakness or the sensation of being touched or grabbed.

Third phase is Headache. A migraine usually starts gradually and builds in intensity. It is possible to have migraine without a headache.

Fourth phase is Postdrome. Individuals are often exhausted or confused following a migraine. The postdrome periods may last up to a day before people feel healthy again.

It appears that migraines are caused by the nerves being more sensitive, and more reactive to stimulation. That stimulation could be stress, fatigue, hunger, almost anything. Migraines run in families. In fact, most migraine sufferers have someone in the family who gets migraines too.

The scientific reasons of migraine are those trigger chemicals, such as serotonin to narrow the blood vessels. Serotonin is a chemical necessary for communication between nerve cells. It can cause narrowing the blood vessels throughout the body. When serotonin or estrogen levels change, the result for some is a migraine.

The best way to prevent migraines is to identify and avoid triggers. The triggers are different in each person, which is why it's a good idea to keep a headache diary.

When a child gets a headache, write down what was happening before the headache, how badly it hurt and where, what helped, and anything else about it one can

think of. This helps to see patterns that can help to understand child's particular triggers. It's a good idea to make sure a child gets enough sleep, eats regularly and healthfully, drinks water regularly, gets exercise, and manages stress. Doing this not only helps prevent migraines, but is also good for overall health.

When a migraine strikes, sometimes just lying down in a dark, quiet room with a cool cloth on the forehead is enough. If it's not, ibuprofen or acetaminophen can be helpful. It's important not to give the child these medications more than about 14 days a month, as giving them more often can lead to rebound headaches and make everything worse. If those approaches aren't enough, a class of medications called triptans can be helpful in stopping migraines in children ages 6 and up. If a child experiences frequent or severe migraine, leading to missed days of school or otherwise interfering with life, doctors often use medications to prevent migraines. There are a number of different kinds, and your doctor can advise you on what would be best for your child. Some girls get migraines around the time of their period. If that happens frequently, sometimes taking a prevention medicine around the time of menses each month can be helpful.

REFERENCES

1. Stovner LJ, Hagen K, Jensen R, et al. The global burden of headache: A documentation of headache prevalence and disability worldwide. *Cephalalgia* 2007;27:193-210.
2. Hu XL, Markson RB, Lipton RB. Disability and economic costs of migraine in the United States. *Ann Intern Med* 1999; 159: 813-818.
3. Hawkins K, Wang S, Rupnow M. Direct cost burden among insured US employees with migraine. *Headache: J Head Face Pain* 2008; 48:553-563.
4. Stewart WF, Lipton RB, Simon D. Work-related disability: Results from the American migraine study. *Cephalalgia* 1996; 16: 231-238.
5. Etemad LR, Yang W, Globe D, Barlev A, Johnson KA. Costs and utilization of triptan users who receive drug prophylaxis for migraine versus triptan users who do not receive drug prophylaxis. *J Manag Care Pharm* 2005;11: 137-144.
6. Henry P, Auray JP, Gaudin AF, et al. Prevalence and clinical characteristics of migraine in France. *Neurol* 2002;59:232-237.
7. Kelman L. The triggers or precipitants of the acute migraine attack. *Cephalalgia* 2007;27: 394-402.
8. Bener A, Uduman SA, Qassimi EMA, et al. Genetic and environmental factors associated with migraine in schoolchildren. *Headache: J Head Face Pain* 2000; 40: 152-157.
9. Spierings ELH, Ranke AH, Honkoop PC. Precipitating and aggravating factors of migraine versus tension-type headache. *Headache: J Head Face Pain* 2001; 41: 559-564.

Frequency of Cardiac Arrest and Arrhythmias Associated with Admitted Patients in the Cardiac Care Unit in Hazara Division

Sardar Fawad Gul¹, Mohsin Khan², Mohammad Imran Khan³, Zia Qamar², Adnan Haider² and Zulqarnain Dilawar¹

ABSTRACT

Objective: The present study will determine the arrhythmias proportion associated with sudden cardiac arrest in patients admitted to cardiac care unit.

Study Design: Cross-sectional study

Place and Duration of Study: This study was conducted at the Cardiology Department of Ayub Teaching hospital from April, 2023 to November, 2023.

Methods: The patients who needed cardiac resuscitation were accessed and were included in current study. We recorded initial cardiac rhythm, features of cardiac arrhythmia, demographic variables, comorbidities, and in-hospital complications till the patient's discharge. The categorical variables were presented in terms of percentages while continuous variables were presented as mean and standard deviation. Analysis of variance and chi-square was used to measure the significance which is kept less than 0.005. The data was analyzed using SPSS version 21.

Results: A total of 220 patients met the inclusive criteria and had SCA. The mean age of the studied sample was 65±9 years. The most common rhythm analyzed initially was VT in 32% (70) of individuals followed by PEA in 24% (53). Myocardial ischemia was found to be the most common immediate precipitating cause of arrhythmias (38%). ROSC was attained in 47% of patients, among which 58 patients survived to discharge (STD). PEA carried the worst mortality overall among cardiac arrhythmias while VT had the most favorable outcome with a proportion of 48.5% (34/70).

Conclusion: We concluded that shockable rhythm i.e. VT is still prevalent in developing countries like Pakistan while globally the behavior of arrhythmias has been changed to non-shockable rhythms of unknown cause. The patients among which ROSC was restored within 5-10 minutes survived to discharge with good neurological status. However, it was observed that as the time for ROSC is prolonged, the neurological outcome and survival rate declines. Overall, PEA carries the worst mortality.

Key Words: Cardiac care unit (CCU), shockable rhythm, cardiac arrhythmias, resuscitation, restoration of spontaneous circulation (ROSC).

Citation of article: Gul SF, Khan M, Khan MI, Qamar Z, Haider A, Dilawar Z. Frequency of Cardiac Arrest and Arrhythmias Associated with Admitted Patients in the Cardiac Care Unit in Hazara Division. Med Forum 2024;35(1):3-7. doi:10.60110/medforum.350101.

INTRODUCTION

Rapid recognition of peri-infarction malignant arrhythmias in modern cardiac care units evolved into different terrains, where patients suffering from heterogeneous concomitant comorbidities may now be treated successfully. Globally, cardiovascular disease (CVD) is the major cause of morbidity and death¹.

¹. Department of Cardiology / Physician² / Interventional Cardiology³, Ayub Teaching Hospital Abbottabad.

Correspondence: Dr. Mohsin Khan, Resident Physician, Ayub Teaching Hospital Abbottabad.

Contact No: 03065531722

Email: mohsinkhan1464@yahoo.com

Received: December, 2023

Accepted: December, 2023

Printed: January, 2024

From 1990 to 2019, there were approximately twice as many CVD prevalence cases (271 million) as there were fatalities due to CVD (12.1 million to 18.6 million)². Current literature suggests that 80% of CVD cases occur in low- and middle-income countries, because of urbanization and ageing^{3,4}. According to the study done in Khyber Pakhtunkhwa, the prevalence of cardiovascular disease (CVD) is 17.5%⁵.

In a developing country like Pakistan, advanced health care like Primary percutaneous coronary intervention (PCI) is a big challenge to be available in every tertiary care hospital. For this reason, pharmacological thrombolysis is the most common method used in tertiary care hospitals⁶. In our country, due to low socioeconomic status, the presentation of cardiac events to hospitals is delayed. According to one institutional study, the attributed symptoms of other diseases is the most common reason for delayed presentation to hospital⁵. The delayed presentation and underlying comorbidities subject the patient to life-threatening

cardiac arrhythmia. Delayed diagnosis and thrombolysis lead to late blood flow storage, which increases infarct size, and subjects the patient to cardiac arrhythmia. So, the majority of CVD patients—about a third died upon arrival at the hospital, so timely diagnosis and immediate interventions are necessary⁷. Missing thrombolytic treatment also creates a group of individuals at high risk for developing later cardiac events, such as mortality, heart attack, stroke failure, and potentially fatal arrhythmias⁸.

Underlying ischemic heart disease is the great victim of sudden cardiac arrest (SCA) in adults. Based on electrophysiology, the SCA can be categorized into ventricular tachycardia (VT), ventricular fibrillation (VF), asystole (ASY), and pulseless electrical activity (PEA). Studies have shown that in-hospital cardiac arrest associated with VT/VF can be survived to discharge because of shockable rhythm⁹. Previously it was thought that VT and VF were the most common arrhythmias associated with SCD but currently, the incidence has declined contrary to PEA/ASY— whose incidence increased for unknown reasons. Over time this change in the behavior of arrhythmias also necessitates to study of the nature and prognostic implications of cardiac arrhythmias. PEA/ASY is more prevalent and accounts for 30% of cardiac mortality and 20% of total mortality in adults¹⁰. Globally the overall survival rate from cardiac arrest is <10%¹¹.

It's important to assess the characteristics of different type of arrhythmias and their response to intervention. With an extensive review of the literature, we have found that the objective data on acute cardiac care and arrhythmias in CCUs are still scarce, and producing evidence-based institutional guidelines remains an issue. Starting from this background we have decided to undertake this study to know the major types of arrhythmias and response to intervention. Moreover, the present study also determined the demographic characteristics, comorbidities, and underlying causes of arrhythmias. This will create reliable data that help treat CCU physicians in better understanding and prevention of arrhythmias in cardiac patients.

METHODS

This was a prospective cross-sectional study that recruited all consecutive admissions to the cardiac care unit (CCU) of Ayub Teaching Hospital from 1st April 2023 to 30th November 2023. The institutional ethical committee approved the study with the safety of human subjects. A non-probability purposive sampling technique was applied for sample collection. The source of data collection was an online predesigned Google form, which was filled out by the healthcare professional and submitted with a special ID code.


The CCU at this facility is a single 16-bed, in which all admissions are triaged, accepted, and cured by a certified cardiologist of the institution. Information

regarding cardiac arrhythmia, treatment, interventional procedure, and end event was explored from patient's files and electronic data from the Hospital management system (HMS). We categorized primary arrhythmias into seven different groups, namely ventricular fibrillation, ventricular tachycardia, pulseless electrical activity, asystole, torsade de point, supraventricular tachycardia, and unstable atrial fibrillation. We recorded the frequency and type of arrhythmias during admission. Moreover, the treatment, procedure, stay and outcome were also recorded for each patient. The primary endpoint of the study was to determine the frequency and type of arrhythmia in patients admitted to CCU. The patients with asymptomatic arrhythmia, hemodynamically stable atrial fibrillation, asymptomatic bradyarrhythmia, and diagnosed arrhythmia patients who don't need CCU care were excluded from the study.

Interquartile ranges (IQRs) and medians were used to display continuous data, while absolute numbers and percentage values were used to show categorical variables. The chi-square test was used to compare categorical variables, and the one-way analysis of variance was used to compare continuous variables. A p-value less than 0.05 was considered significant. The data was analyzed using SPSS version 21.

RESULTS

About 350 patients were presented with arrhythmias in current study but only 220 patients met the inclusive criteria (n=220) subjects of cardiac arrest were recruited with known initial cardiac rhythm. The mean age of the sample was 65±9years. 77% sampled population was obese with a mean BMI of 34kg/m². The relation of an age with type of arrhythmia is shown in graphs 01.



Graph No. 1: Relation age and type of arrhythmia (p=0.000)

Over 92% sampled population was diabetic (204/220) and only 9% (20/220) had controlled diabetes. Eighty-five percent of patients (188/220) were hypertensive and 65% had a duration greater than 5 years. The most common rhythm recorded initially on ECG was VT in 70 patients (32%) followed by, 53 PEA patients (24%), ASY49(22%), VF 40(18%), and SVT 8(4%) patients. No cases of torsade de points and unstable atrial fibrillation were recorded during the study period. The

characteristics are shown in table 1. Myocardial infarction was the most common immediate factor found for arrhythmia with frequency of 84/220(38%), followed by CCF 76(34.5%), arrhythmia with undetermined cause in 53(24%), pulmonary embolism 06(3%) and hypotension in 1 (0.4%) patient. Dilated cardiomyopathy was the most common underlying cardiac comorbidity with frequency of 74(33.6%), Ischemic cardiomyopathy 56(25.4%), valvular 50(22.7%), hypertrophic cardiomyopathy 24(11%) and

no underlying structural heart disease was found in 16(7%) of individuals. About 162 (73.6%) individuals died because of arrhythmia, 44(20%) survived with a good neurological state at hospital discharge while 14(6.3%) had poor neurological sequelae. All patients with PEA died in the hospital while 26 VT patients survived in a good conscious state at the hospital discharge detailed in Table 1. The factor which affects mortality is listed in Table 2.

Table No. 1: Different features of cardiac arrhythmia in patients of CCU

Characteristics	Asystole (ASY) n=49 (22%)	Pulseless Electrical activity (PEA) n=53(24%)	Supra ventricular Tachycardia (SVT)n=08(4%)	Ventricular Fibrillation (VF) n=40 (18%)	Ventricular Tachycardia (VT) n=70 (32%)	P value	
Age(years)	71.9	64.5	62	70	65	0.000	
BMI (Kg/m ²)	34	25	38	31	34	0.003	
Hypertension	45	47	09	44	47	0.005	
Diabetes	Poor control	42	45	20	34	0.026	
	Controlled	06	04	02	02		
	Non-diabetic	04	03	04	02		
Event location	CCU	44	52	10	38	0.003	
	Ward	05	01	00	05		
	Public	00	00	03	00		00
Pharmacological agent given for arrhythmia	Amiodarone	12	07	05	23	0.000	
	Atropine	01	02	01	01		03
	Digoxin	09	01	01	05		02
	Epinephrine	02	03	01	01		01
	lidocaine	01	01	01	01		00
	Nor epinephrine	16	28	00	19		19
Preexisting cardiac comorbidities	DCM	28	18	03	13	0.000	
	HCM	01	03	01	09		10
	ICMP	04	22	03	11		16
	Valvular	16	10	01	11		12
	unknown	04	02	05	00		05
Electrical cardioversion	12	06	06	25	58	0.000	
Discharge status	Survived with good conscious	05	00	07	05	0.000	
	Survived with poor conscious	00	00	02	05		07
	Died in hospital	44	53	01	35		29

Table No.2: Factors significantly affecting hospital discharge status in patients experiencing cardiac arrest

Factors	Died in hospital 162(73%)	Survived with poor conscious status 14(6.3%)	Survived with good conscious status44 (20%)	Total	Significance level
Diabetes	Poor controlled	140	14	30	0.001
	Good controlled	10	00	10	
	nondiabetic	12	00	04	
Hypertension	yes	138	12	38	0.017
	no	24	02	06	
Event location	CCU	148	13	40	0.03
	ward	12	1	3	

	Public places	02	00	01	03	
	ASY without subsequent VT/VF	44	06	10	60	0.000
	PEA without subsequent VT/VF	42	16	10	68	0.001
Underlying comorbidities	DCM	54	05	15	74	0.000
	HCM	18	1	5	24	
	ICMP	41	4	11	56	
	Valvular	37	3	10	50	
	No cause	12	1	3	16	
Return of spontaneous circulation	<5min	01	01	00	02	0.000
	5-10min	19	04	13	36	
	10-20min	19	04	19	42	
	>20min	05	05	12	22	
	Not return	118	00	00	118	
Pharmacological agent	Amiodarone	57	05	16	78	0.001
	Atropine	06	00	02	08	0.001
	Digoxin	13	01	04	18	
	Epinephrine	06	01	01	08	
	Lidocaine	03	00	01	04	
	Nor-epinephrine	61	05	16	82	
	DNR	16	02	04	22	

DISCUSSION

We comprehensively evaluated 220 patients for cardiac arrhythmias who experienced sudden cardiac arrest either in public places, hospital wards, or CCU. Besides the high prevalence of CVD, the national literature regarding arrhythmia in CCU is still scarce.

In the present study, the mean age of sampled data who experienced sudden cardiac arrest is older than reported by Faiza Ahmed et al¹². — the recent study conducted in Karachi. The reason could be the small sample size, demography, and period of the COVID-19 pandemic. Overpopulation is older than reported by Moosajee, U.S. et al¹³ and Khan NU et al¹⁴ but interestingly the mean age is near to that reported by Tseng ZH et al¹⁵. The sampled population who experienced cardiac arrest due to VF, ASY, and PEA are older than the population with VT and SVT. We observed that the frequency of VT/VF (shockable rhythm) is still higher than reported previously by Faiza Ahmed et al¹². Although the global incidence of VT/VF is declining because of unknown cause¹¹. The reason for the difference could be the hospital setting and sample size, we observed arrhythmia in CCU while Faiza Ahmed et al observed it in Medical ICU, CCU, COVID-ICU, and HDU¹². Cardiac ischemia is the most common immediate factor responsible for arrhythmias and dilated cardiomyopathy is the most common preexisting cardiac comorbidity in patients who experience sudden cardiac arrest¹¹. Contrary to Faiza Ahmed et al. respiratory insufficiency is the most common immediate factor responsible for arrhythmia and ASY is the most common initial rhythm observed in sudden cardiac arrest¹². Unfortunately, over 85% of all patients who experienced SCA are diabetic, hypertensive with obesity.

ROSC was achieved in 47% of patients, and survived to discharged (STD) proportion is 24%. In comparison

with Moosajee, U.S. et al 27% ROSC was achieved and 7.5% STD. The difference in proportion is likely due to event location. In our study, 99% of patients experienced hospital SCD while in Moosajee, U.S. et al only 33% of patients experienced SCD in the hospital. It is predetermined that hospital cardiac arrest has more survival than hospital SCD¹². All patients in whom the ROSC achieved less than 5 min survived, and greater than 50% in whom ROSC achieved less than 20 min survived while mortality increased after 20 min ROSC. In terms of initial rhythm vs STD, 55% VT and 18% VF (shockable rhythm) STD as compare to 12% ASY and 00% PEA (non-shockable rhythm). These findings are almost consistent as reported by Moosajee, U.S. et al¹³. A negligible mortality difference is noted in patients who experienced ASY vs patients who experienced ASA with subsequent VT/VF while a significant difference is noted in patients with good neurological outcomes in survived patients. A significant difference is noted in the mortality of patients who experienced PEA with subsequent VT/VF Vs only PEA. Mortality is higher in patients with only the PEA group vs the PEA with the VT/VF group which is 80% vs 61% respectively. 45% of patients STD who initially received amiodarone as initial antiarrhythmic while epinephrine was commonly antiarrhythmic reported by Moosajee, U.S. et al¹³. The variance could be due to a relative difference in the proportion of initial cardiac rhythm.

CONCLUSION

The present study concluded that the patients with cardiac arrhythmia are older than previously reported at global and national levels. Over 85% sampled population have poorly controlled hypertension and diabetes. All patients encountered in CCU with arrhythmia are overweight. The incidence of VT is still

high in developing countries like Pakistan despite global decline. The most common immediate cause responsible for arrhythmia is myocardial ischemia. The proportion of ROSC was high in patients who experienced cardiac arrest inside the hospital. The survival to discharge with a good conscious state is high in patients whose ROSC is 5-10min, while mortality and poor neurological outcome drastically increased in patients with whom ROSC is \geq 20min. PEA carries the worst in-hospital mortality, all patients died during management, while shockable rhythm—VT carries favorable discharge status among all arrhythmias..

Author's Contribution:

Concept & Design of Study: Sardar Fawad Gul
 Drafting: Mohsin Khan,
 Mohammad Imran Khan
 Data Analysis: Zia Qamar, Adnan
 Haider, Zulqarnain
 Dilawar
 Revisiting Critically: Sardar Fawad Gul,
 Mohsin Khan
 Final Approval of version: Sardar Fawad Gul

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.RC-EA-2023/055 dated 02.04.2023

REFERENCES

- Roth GA, Mensah GA, Johnson CO, Addolorato G, Ammirati E, Baddour LM, et al. Global burden of cardiovascular diseases and risk factors, 1990–2019: update from the GBD 2019 study. *J Am Coll Cardiol* 2020;76(25):2982-3021.
- Li Y, Cao GY, Jing WZ, Liu J, Liu M. Global trends and regional differences in incidence and mortality of cardiovascular disease, 1990–2019: findings from 2019 global burden of disease study. *Eur J Preventive Cardiol* 2023;30(3):276-86.
- Sharif Z, Ptaszek LM. Global disparities in arrhythmia care: mind the gap. *Heart Rhythm* 2022;3(6):783-92.
- Schwalm JD, McKee M, Huffman MD, Yusuf S. Resource effective strategies to prevent and treat cardiovascular disease. *Circulation* 2016;133(8):742-55.
- Haq I. Epidemiological Assessment and Frequency of Cardiovascular Diseases in Peshawar KP, Pakistan: A Cross-Sectional study. *Annals Romanian Society Cell Biol* 2021;25(7):1377-88.
- Haq I. Epidemiological Assessment and Frequency of Cardiovascular Diseases in Peshawar KP, Pakistan: A Cross-Sectional study. *Annals Romanian Society Cell Biol* 2021;25(7):1377-88.
- Hasan A, Tawab S, Arif M, Muzamil M, Javed A, Aslam M. Effect of Gender on Efficacy of Streptokinase in Acute ST Segment Elevation Myocardial Infarction. *Pak Heart J* 2019;52(3).
- Stefanini GG, Alfonso F, Barbato E, Byrne RA, Capodanno D, Colleran R, et al. Management of Myocardial Revascularization Failure: An Expert Consensus. *Trials* 2018;39:2192-07.
- Woolcott OO, Reinier K, Uy-Evanado A, Nichols GA, Stecker EC, Jui J, et al. Sudden cardiac arrest with shockable rhythm in patients with heart failure. *Heart Rhythm* 2020;17(10):1672-8.
- Rodríguez-Reyes H, Muñoz-Gutiérrez M, Salas-Pacheco JL. Current behavior of sudden cardiac arrest and sudden death. *Archivos de cardiología de México* 2020;90(2):183-9.
- Kaupila JP, Hantula A, Kortelainen ML, Pakanen L, Perkiömäki J, Martikainen M, et al. Association of initial recorded rhythm and underlying cardiac disease in sudden cardiac arrest. *Resuscitation* 2018;122:76-8.
- Ahmed F, Abbasi L, Ghouri N, Patel MJ. Epidemiology of in-hospital cardiac arrest in a Pakistani tertiary care hospital pre-and during COVID-19 pandemic. *Pak J Med Sci* 2022;38(2):387.
- Moosajee US, Saleem SG, Iftikhar S, Samad L. Outcomes following cardiopulmonary resuscitation in an emergency department of a low-and middle-income country. *Int J Emerg Med* 2018;11:1-9.
- Khan NU, Razzak JA, Ahmed H, Furqan M, Saleem AF, Alam H, et al. Cardiopulmonary resuscitation: outcome and its predictors among hospitalized adult patients in Pakistan. *Int J Emerg Med* 2008;1:27-34.
- Tseng ZH, Olgin JE, Vittinghoff E, Ursell PC, Kim AS, Sporer K, et al. Prospective countywide surveillance and autopsy characterization of sudden cardiac death: POST SCD study. *Circulation* 2018;137(25):2689-700.

Dentists' and Dental Students' Perspectives on Amalgam Restoration in Saudi Arabia

Abdulaziz Alhumaid¹, Abdulaziz Alrebdi², Ebrahim Alshawy³, Nawaf Almutairi² and Yasir Alyahya² and Ahmed Alharbi¹

ABSTRACT

Objective: The study aims to assess dentists and dental students prospectively regarding amalgam and their opinions when dealing with it in the dental practice in Saudi Arabia.

Study Design: A cross-sectional study.

Place and Duration of Study: This study was conducted at the Qassim University's College of Dentistry from April 2021 to May 2023.

Methods: A cross-sectional study among dentists and dental students in the Kingdom of Saudi Arabia was conducted. The study included dentists from both public and private health care facilities. Data was collected using online google form, which consisted of demographic information and knowledge, attitude and practices of participants toward the use of amalgam material. Data was analyzed using chi-square test.

Results: Out of 122 participants, the majority were students (n=62), followed by specialists (n=25) and general practitioners (GP) dentists (n=22). A significantly higher percentage of participants revealed that the amalgam is not dangerous in the workplace (n=60, 49.2%). Moreover, a number of 81 participants (66.4%) said that amalgam restorations cannot be replaced by resin restoration. The specialist reported that most of their patients (n=22, 88.0%) do not prefer amalgam restorations due to the dark colour of the amalgam.

Conclusion: According to the results of this study, dental professionals in Saudi Arabia believe that amalgam is a risk-free material. Alternatives to dental amalgam are generally preferred by dental professionals mainly for esthetic concerns.

Key Words: Amalgam, Dentist, Dental Amalgam, Dental restoration, filling, KAP, Mercury

Citation of article: Alhumaid A, Alrebdi A, Alshawy E, Almutairi N, Alyahya Y, Alharbi A. Dentists' and Dental Students' Perspectives on Amalgam Restoration in Saudi Arabia. Med Forum 2024;35(1):8-13. doi:10.60110/medforum.350102.

INTRODUCTION

Dental amalgam has been commonly used as a dental restorative material for over 150 years and has provided patients with a valuable and comparatively affordable treatment ever since. The data that is currently available demonstrates that dental amalgams are beneficial and risk-free; however, some of these concerns have been brought to light.¹

Even though there is ongoing debate regarding the impact that waste containing mercury its physical properties, longevity, and less manual dexterity, this substance holds high importance in public health services.^{2,3} Likewise national health authorities advised in previous years that amalgam should not be the first choice when placing restorations.⁴ In addition, there is a discernible effort in the direction of dental educational institutions placing a greater emphasis on teaching dental students how to use mercury-free substitutes, in accordance with the guidelines outlined in the Minamata Convention on Mercury.

According to reports, public and media discourse influences dentists' views on amalgam.⁵ In their survey, Khairuldean and Sadig reported that 75% of Saudi Arabian dentists were aware of the controversy surrounding the safety of amalgam.⁵ Approximately, 85% of these dentists believed amalgam to be safe, while only 41% have been cognizant of all of the clinical symptoms of amalgam toxicity. Patients' perspective of dental aesthetics indicates that part of the population is unsatisfied with the metallic colour of the restorations in their teeth. Even so, there was an indication of a transition away from using silver

¹. College of Dentistry, Qassim University, Qassim, Saudi Arabia.

². Department of Conservative Dental Sciences, College of Dentistry, Qassim University, Qassim, Saudi Arabia.

³. Department of Orthodontics and Pediatric Dentistry, College of Dentistry, Qassim University, Qassim, Saudi Arabia.

Correspondence: Abdulaziz Alrebdi, Department of Conservative Dental Sciences, College of Dentistry, Qassim University, Qassim, Saudi Arabia.
Contact No: +1 9296726710
Email: AB.ALREBDI@qu.edu.sa

Received: October, 2023
Accepted: December, 2023
Printed: January, 2024

amalgam to much more aesthetic tooth-coloured restorations over the last decade.⁶ This could explain why dentists' and patients' preferences have changed in recent years. We tried to find out the answers to these issues in this report. This research looks at how a subset of dentists and students thought about amalgam and how they felt about dealing with it in Saudi Arabia.

METHODS

A cross-sectional study among dentists in the Kingdom of Saudi Arabia was conducted. This study was carried out at Qassim University's College of Dentistry from April 2021 to May 2023, with approval from the College of Dentistry Research Center (Reg. EA/6110/2021). A sample of 10 dentists and dental students just beginning clinical practice in Saudi Arabia was used for the pilot test of the questionnaire to evaluate its clarity in terms of both its structure and its content. The inclusion criteria were dental students beginning clinical practice and dentists. The study included dentists from both public and private healthcare facilities. Junior students who had not begun their clinical practice were among the exclusion criteria. Reminder emails were sent in two stages to enhance the response rates: one in February 2022 and another in November 2022. The questionnaire has 18 closed-ended questions and it was divided into two sections. The first section was regarding the demographic data and the characteristics of the practice (private practice, government hospitals). The second part focused on

measuring the perspective and knowledge of dentists regarding amalgam use and hazards, case selection, and safety concerns. The last question that was posed to the participants inquired as to whether or not they routinely employed amalgam restorations in their professional work. The collected data were analyzed using Statistical Package for Social Science (SPSS Inc). Descriptive statistics and frequency tables were compiled in order to provide an overview of the responses. For the statistical data analysis, the Chi-square test was utilized for categorical variables, and Fisher's exact test was used to assess the variations among groups.

RESULTS

A cross-sectional study was conducted among dental professionals between April 2021 to May 2023. Before the start of the study, Institutional ethical clearance was obtained from the College of Dentistry, Qassim University. The present study questionnaire was sent to 150 participants. Only 122 participants completed the questionnaire (response rate = 81.3%). Out of 122 participants, a number of 80 participants (65.6%) were males and 42 participants (34.4%) were female. It was observed that, out of the 122 participants, the majority of them were students (n=62), followed by specialists (n=25) and GP dentists (n=22). Most of the specialists and GP dentists were working in the government sector (n=108, 88.5%), while 14 participants (11.5%) were working in the private sector (Table 1).

Table No.1: Demographic Profile

Variables	Student	Intern	GP dentist	Specialist	Consultant	Total	p-Value
Gender							
Male	37 (59.7%)	3 (27.3%)	17 (77.3%)	21 (84.0%)	2 (100.0%)	80 (65.6%)	0.007
Female	25 (40.3%)	8 (72.7%)	5 (22.7%)	4 (16.0%)	0 (0.0%)	42 (34.4%)	
Age							
20-30	62 (100.0%)	11(100.0%)	22(100.0%)	25(100.0%)	2 (100.0%)	122(100.0%)	
Service sector							
Private	1 (1.6%)	0 (0.0%)	8 (36.4%)	5 (20.0%)	0 (0.0%)	14 (11.5%)	0.000
Government	61 (98.4%)	11(100.0%)	14(63.6%)	20 (80.0%)	2 (100.0%)	108(88.5%)	

Table No. 2: Knowledge about Amalgam

Variables	Student	Intern	GP dentist	Specialist	Consultant	Total	p-Value
5. Patients perception in the amalgam toxicity comes from?							
Social media	49 (79.0%)	10 (90.9%)	16 (72.7%)	19 (76.0%)	1 (50.0%)	95 (77.9%)	0.358
Research	6 (9.7%)	1 (9.1%)	3 (13.6%)	1 (4.0%)	0 (0.0%)	11 (9.0%)	
Banned	5 (8.1%)	0 (0.0%)	2 (9.1%)	3 (12.0%)	0 (0.0%)	10 (8.2%)	
The long contravery	2 (3.2%)	0 (0.0%)	1 (4.5%)	2 (8.0%)	1 (50.0%)	6 (4.9%)	
6.What is your patient's view on the amalgam fillings?							
Safe	13 (21.0%)	3 (27.3%)	1 (4.5%)	11 (44.0%)	1 (50.0%)	29 (23.8%)	0.026
Unsafe	5 (8.1%)	2 (18.2%)	6 (27.3%)	5 (20.0%)	0 (0.0%)	18 (14.8%)	
Uncertain	44	6 (54.5%)	15	9 (36.0%)	1 (50.0%)	75 (61.5%)	

	(71.0%)		(68.2%)				
9- What is your perception in using dental amalgam							
Higher strength and longevity	39 (62.9%)	4 (36.4%)	8 (36.4%)	13 (52.0%)	1 (50.0%)	65 (53.3%)	0.128
Less technique sensitive	9 (14.5%)	2 (18.2%)	2 (9.1%)	2 (8.0%)	1 (50.0%)	16 (13.1%)	
Unconservative restorations	14 (22.6%)	5 (45.5%)	12 (54.5%)	10 (40.0%)	0 (0.0%)	41 (33.6%)	
15- Is dental amalgam an occupation hazard at your place of work?							
Yes	28 (45.2%)	10 (90.9%)	13 (59.1%)	10 (40.0%)	1 (50.0%)	62 (50.8%)	0.047
No	34 (54.8%)	1 (9.1%)	9 (40.9%)	15 (60.0%)	1 (50.0%)	60 (49.2%)	
Total	62 (100.0%)	11 (100.0%)	22 (100.0%)	25 (100.0%)	2 (100.0%)	122 (100.0%)	

Table 3: Attitude about Amalgam use

Variables	Student	Intern	GP dentist	Specialist	Consultant	Total	P- Value
12- How often do you use amalgam for restorations in your daily clinical practice?							
Always	1 (1.6%)	1 (9.1%)	0 (0.0%)	2 (8.0%)	0 (0.0%)	4 (3.3%)	0.003
Sometime	4 (6.5%)	2 (18.2%)	2 (9.1%)	10 (40.0%)	1 (50.0%)	19 (15.6%)	
Rarely	19 (30.6%)	4 (36.4%)	3 (13.6%)	7 (28.0%)	0 (0.0%)	33 (27.0%)	
Never	38 (61.3%)	4 (36.4%)	17 (77.3%)	6 (24.0%)	1 (50.0%)	66 (54.1%)	
14- What is your reasons not to use amalgam filling?A							
Mercury toxicity	27 (43.5%)	3 (27.3%)	5 (22.7%)	3 (12.0%)	0 (0.0%)	38 (31.1%)	0.001
Unesthetics	7 (11.3%)	3 (27.3%)	7 (31.8%)	12 (48.0%)	1 (50.0%)	30 (24.6%)	
Patient's desire	25 (40.3%)	2 (18.2%)	6 (27.3%)	3 (12.0%)	0 (0.0%)	36 (29.5%)	
Unconservative	3 (4.8%)	3 (27.3%)	4 (18.2%)	7 (28.0%)	1 (50.0%)	18 (14.8%)	
16- Do you agree or disagree that successful amalgam restorations can be replaced with composite resin?							
Agree	25 (40.3%)	2 (18.2%)	8 (36.4%)	5 (20.0%)	1 (50.0%)	41 (33.6%)	0.307
Disagree	37 (59.7%)	9 (81.8%)	14 (63.6%)	20 (80.0%)	1 (50.0%)	81 (66.4%)	
18- Do you agree or disagree with the established ban on amalgam use in some countries?							
Agree	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	-
Disagree	62(100.0%)	11 (100.0%)	22(100.0%)	25(100.0%)	2 (100.0%)	122(100.0%)	
Total	62(100.0%)	11 (100.0%)	22 (100.0%)	25(100.0%)	2 (100.0%)	122(100.0%)	

Table No.4: Practice of amalgam restoration

Variables	Student	Intern	GP dentist	Specialist	Consultant	Total	p-Value
7- What is your patient concerned regarding amalgam filling?							
Color	53 (85.5%)	8 (72.7%)	16 (72.7%)	22 (88.0%)*	1 (50.0%)	100 (82.0%)	0.000
Toxicity	8 (12.9%)	3 (27.3%)	4 (18.2%)	2 (8.0%)	0 (0.0%)	17 (13.9%)	
Health problem	0 (0.0%)	0 (0.0%)	2 (9.1%)	1 (4.0%)	0 (0.0%)	3 (2.5%)	
Environment effects	1 (1.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (50.0%)	2 (1.6%)	
8- Would you recommend an alternative to amalgam?							
Yes	47 (75.8%)	8 (72.7%)	18 (81.8%)	19 (76.0%)	1 (50.0%)	93 (76.2%)	0.761
No	8 (12.9%)	2 (18.2%)	3 (13.6%)	4 (16.0%)	0 (0.0%)	17 (13.9%)	

Uncertain	7 (11.3%)	1 (9.1%)	1 (4.5%)	2 (8.0%)	1 (50.0%)	12 (9.8%)	
10- What criterion will you use to decide whether or not to replace amalgam restoration? p							
Patient wishes	22 (35.5%)	4 (36.4%)	4 (18.2%)	4 (16.0%)	0 (0.0%)	34 (27.9%)	0.100
Aesthetic	15 (24.2%)	2 (18.2%)	1 (4.5%)	5 (20.0%)	1 (50.0%)	24 (19.7%)	
Criteria of defective restoration	25 (40.3%)	5 (45.5%)	17 (77.3%)	16 (64.0%)	1 (50.0%)	64 (52.5%)	
11- Which of the following amalgam substitutes do you prefer? P							
Glass ionomer and resin modified glass ionomer	11 (17.7%)	1 (9.1%)	3 (13.6%)	1 (4.0%)	0 (0.0%)	16 (13.1%)	0.218
Indirect restoration inlay and onlay	23 (37.1%)	7 (63.6%)	5 (22.7%)	7 (28.0%)	1 (50.0%)	43 (35.2%)	
Resin composite	28 (45.2%)	3 (27.3%)	14 (63.6%)	17 (68.0%)	1 (50.0%)	63 (51.6%)	
13- In which of the following do you use dental amalgam?P							
Simple cavity	7 (11.3%)	0 (0.0%)	0 (0.0%)	1 (4.0%)	0 (0.0%)	8 (6.6%)	0.141
Complex cavity	18 (29.0%)	5 (45.5%)	6 (27.3%)	14 (56.0%)	0 (0.0%)	43 (35.2%)	
Core material	8 (12.9%)	1 (9.1%)	1 (4.5%)	2 (8.0%)	1 (50.0%)	13 (10.7%)	
Not used	29 (46.8%)	5 (45.5%)	15 (68.2%)	8 (32.0%)	1 (50.0%)	58 (47.5%)	
17- If your patient had faulty amalgam restoration, what is your retreatment plan?							
Amalgam	3 (4.8%)	0 (0.0%)	2 (9.1%)	2 (8.0%)	0 (0.0%)	7 (5.7%)	0.888
Composite	26 (41.9%)	4 (36.4%)	12 (54.5%)	11 (44.0%)	1 (50.0%)	54 (44.3%)	
Indirect restoration inlay and onlay	33 (53.2%)	7 (63.6%)	8 (36.4%)	12 (48.0%)	1 (50.0%)	61 (50.0%)	
Total	62 (100.0%)	11 (100.0%)	22 (100.0%)	25 (100.0%)	2 (100.0%)	122 (100.0%)	

Comparative analysis of different dental care professionals and their knowledge, attitude and practices are shown in Tables 2,3 and 4. It was observed that the majority of the participants reported that Social media 95 (77.9%) was responsible for spreading myths about amalgam toxicity. However, a number of 75 participants (61.5%) were uncertain about their view on amalgam safety. On the other hand, most of the participants reported that amalgam is not hazardous at the workplace (n=60, 49.2%) (Table 2). Table 3 illustrates the attitude of study participants toward amalgam use. It was surprising to know that even though they said it was not hazardous. Most of the participants (n=66 (54.1%)) do not use amalgam restoration in their daily routine. Moreover, a number of 81 participants (66.4%) disagree that amalgam restorations can be replaced by resin restoration. A significant number of respondents (31.1%) chose "Mercury toxicity" as their reason for not using

amalgam fillings, followed by "Patient's desire" (29.5%) and then "Unesthetics" (24.6%). Lastly, participants' perception of their amalgam practice was recorded. Specialists reported that most of the time, their patients do not prefer amalgam restorations due to colour (n=22 (88.0%)) with a p-value of <0.05. The majority of GPs (n=18 (81.8%)) and consultants (n=19 (76.0%)) recommend an alternative to amalgam. The resin composite was found to be the most recommended material by GPs (n=14 (63.6%)) and consultants (n=17 (68.0%)). Most of the consultants (n=14 (56.0%)) would use amalgam only for complex cavities. However, in the case of faulty amalgam restoration, a number of 61 respondents (50.0%) plan to use indirect (inlay or onlay) restorations (Table 4).

DISCUSSION

The utilization of dental amalgam on a global scale has experienced a sizeable decrease over the past two

decades.^{7,8} A practical assessment of the uses of amalgam in the past and an amalgam ban in the future requires certain presumptions.⁹

In view of this, we asked the patients about their source of information related to amalgam and it was found that Social media (n=95 (77.9%)) was responsible for spreading myths about amalgam toxicity. These results found similar published by other studies.⁹⁻¹¹ When compared with GPs, the current study found that 11% of specialists thought amalgam was safe to use. Nevertheless, a study carried out by Al-Nahedh HN et al. reported that 60.2% of GPs and specialists proclaimed it to be safe for both the dentist and the patient, whereas only 14.9% proclaimed it to be unsafe for both. Al-Nahedh HN et al. and Yaseen et al. reported contradictory results in terms of the percentage of GDPs (53%) and specialists (48%) who consider amalgam to be safe.^{7, 12, 13}

In the current study, the vast majority of intern 10 dentists (90.9 %) and approximately 50% of general practitioner dentists (GP) found amalgam to be an occupationally safe material. The research conducted by Bamise et al¹⁴ revealed that 26% of participants held the belief that mercury could be harmful to the health of humans.¹³

Longevity was cited as the primary reason for considering dental amalgam by the vast majority of respondents findings that were reported by Faraj et al. were consistent with the findings of the current study.^{7,9} It was surprising to know that even though they said it was not hazardous but a number of 66 participants (54.1%) did not use amalgam restoration in their daily routine, which was found to be similar to a study done by another author, who reported that 80% of the participants do not frequently use amalgam restorations in their clinical practice. This could indicate an optimistic influence of the Minamata Convention attributes on Mercury on the dental curriculum.

Moreover, a number of 81 participants (66.4%) in the present study disagreed that amalgam restorations can be replaced by resin restoration. Similarly, Alkhudhairi F discovered that 72% of the people who participated in this study had different opinions. On the other hand, the results of another survey revealed that 21% of dentists removed amalgam restoration at the request of their patients. In the present study, specialists reported that due to colour (n=22 (88.0%)) most of the patients do not prefer amalgam restorations with $p < 0.05$ and the majority of GPs (81.8%) and consultants (76.0%) recommend an alternative to amalgam. This is consistent with the findings of Yaseen.¹² Glass ionomer was the most preferred option as a restorative material in the study conducted by Faraj and coworkers.^{7,9} In terms of colour preferences, similar results were reported by an author and Vidnes- Kopperud et al.¹⁵ favoured esthetics (77.1%) as the main reason to limit the use of amalgam, followed by patients' desire

(58.6%). In addition, Espelid et al. found that regardless of gender, patients were more concerned about the aesthetics of their restorations than the longevity of the restorations.¹⁶

In the current study, to replace faulty amalgam restoration, a number of 61 participants (50.0%) planned to use indirect (inlay or onlay) restorations. While another reported Large restorations (49.4%) and crown build-up (31.5%) were the most common restorative. One of the limitations of the current study is that only a small percentage of dentists participated in the survey, which makes it hard to generalize the results.

CONCLUSION

The results of this study suggested that dentists in Saudi Arabia believe it is safe to use amalgam. Based on these findings, we can draw the conclusion that dental amalgam is well approved by both dentists and patients in Saudi Arabia. Furthermore, the majority of dentists believe that amalgam is safe for both dentists and patients. In general, dentist favor alternatives to dental amalgam for esthetic reasons, and for the most part, dentists do not use dental amalgam routinely as per the requests of their patients.

Acknowledgment: The author(s) gratefully acknowledge Qassim University, represented by the Deanship of Scientific Research, on the financial support for this research under the number (20015-dent-2021-1-2-w) during the academic year 1440 AH / 2019 AD.

Author's Contribution:

Concept & Design of Study:	Abdulaziz Alhumaid, Ahmed Alharbi
Drafting:	Abdulaziz Alrebdi, Ebrahim Alshawy
Data Analysis:	Nawaf Almutairi, Yasir Alyahya
Revisiting Critically:	Abdulaziz Alhumaid, Ahmed Alharbi
Final Approval of version:	Abdulaziz Alhumaid, Ahmed Alharbi

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No. Reg. EA/6110/2021 dated 17.04.2021

REFERENCES

1. Cristina Taut. Dental amalgam: is this the end? J Irish Dental Association 2013;59(4):311-17.
2. Mitchell R JKM, Koike M, Okabe T. Posterior amalgam restorations—usage, regulation, and

- longevity. *Dent Clin North Am* 2007;51(03): 573–589.
3. Rathore M, Singh A, Pant V A. The dental amalgam toxicity fear: a myth or actuality. *Toxicol Int* 2012;19(02):81–88
 4. Vidnes-Kopperud S, Tveit AB, Gaarden T, Sandvik L, Espelid I. Factors influencing dentists' choice of amalgam and tooth-colored restorative materials for Class II preparations in younger patients. *Acta Odontol Scand* 2009;67:74–9.
 5. Alkhudairy F. Attitudes of dentists and interns in Riyadh to the use of dental amalgam. *BMC Res Notes* 2016;9(01):488.
 6. Burke FJT, Wilson NHF, Cheung SW, Mjor IA. Influence of patient factors on age of restorations at failure and reasons for their placement and replacement. *J Dent* 2001;29:317–24
 7. Al-Nahedh HN, El-Hejazi AA, Habib SR. Knowledge and Attitude of Dentists and Patients Toward Use and Health Safety of Dental Amalgam in Saudi Arabia. *Eur J Dent* 2020;14(2):233–238.
 8. Bengtsson UG, Hylander LD. Increased mercury emissions from modern dental amalgams. *Biometals* 2017;30(2):277–283
 9. Faraj BM, Mohammad HM, Mohammad KM. The Changes in Dentists' Perception and Patient's Acceptance on Amalgam Restoration in Kurdistan-Iraq: A Questionnaire-based Cross-Sectional Study. *J Clin Diagn Res* 2015;9(4):ZC22-5.
 10. Harorli O, Harorli H. Evaluation of internet search trends of some common oral problems, 2004 to 2014. *Community Dent Health* 2014;31:188–92.
 11. Udoye C, Aguwa E. Amalgam safety and dentists' attitude: a survey among a Subpopulation of Nigerian dentists. *Oper Dent* 2008;33(4):467–471.
 12. Yaseen SM, AlShahrani I, Daghriri OH, Mufarreh M, Amer A, Tobaiqy S. Dentists perspective on silver amalgam restorations and associated hazards- a cross sectional study. *Br J Med Health Res* 2015;2(6):1–8.
 13. Erçin Ö, Berkmen B, Durukan E, Arhun N. Awareness about dental amalgam among Turkish dentists and patients: a questionnaire and search engine based cross-sectional study. *Int Dent J* 2020;71(2):113–21.
 14. Bamise C, Oginni AO, Adedigba MA, et al. Perception of patients with amalgam fillings about toxicity of mercury in dental amalgam. *J Contemp Dent Pract* 2012;13:289–93.
 15. Vidnes-Kopperud S, Tveit AB, Gaarden T, Sandvik L, Espelid I. Factors influencing dentists' choice of amalgam and tooth-colored restorative materials for Class II preparations in younger patients. *Acta Odontol Scand* 2009;67:74–9.
 16. Kopperud S, Staxrud F, Espelid I, et al. The post-amalgam era: Norwegian dentists' experiences with composite resins and repair of defective amalgam restorations. *Int J Environ Res Public Health* 2016;13:441.

Effects of Intra-Cameral Dexamethasone after Uncomplicated Phacoemulsification

Intra-Cameral
Dexamethasone
after
Phacoemulsification

Muhammad Saad Ullah, Muhammad Usama Rahim and Hassan Shoaib

ABSTRACT

Objective: To assess the efficacy of administering dexamethasone through intra-cameral injection in comparison to the conventional application of topical steroids for individuals undergoing phacoemulsification.

Study Design: Quasi experimental trial study

Place and Duration of Study: This study was conducted at the department of Ophthalmology Ghazi Hospital Dera Ghazi Khan from September 2022 to August 2023.

Methods: Eighty patients of age 55-80 years were enrolled in the study and divided into two groups 1 and 2. In group 1 patient were given intra-cameral dexamethasone and in group 2 equal amount of 0.01 ml normal saline was administered.

Results: At 1st post-operative day mean IOP in group 1 was 14.75 ± 2.55 and in group 2 was 17.85 ± 2.22 . Similarly, at day 7 mean IOP in group 1 was 13.18 ± 2.14 and in group 2 it was 12.44 ± 1.74 . In comparison pre operative IOP was in group 1 and 2 was 16.19 ± 2.54 and 12.44 ± 1.74 respectively.

Conclusion: Intraoperative inflammation following phacoemulsification surgery can be effectively managed by administering intra-cameral dexamethasone injection, resulting in decreased flare and anterior chamber cells, with comparable impacts on IOP and visual acuity when compared to the use of topical steroids.

Key Words: Dexamethasone, Effectiveness, Intra ocular pressure, Intra-cameral injection, Phacoemulsification.

Citation of article: Saad Ullah M, Rahim MS, Shoaib H. Effects of Intra-Cameral Dexamethasone after Uncomplicated Phacoemulsification. Med Forum 2024;35(1):14-17. doi:10.60110/medforum.350103.

INTRODUCTION

Phacoemulsification, a contemporary and widely adopted surgical technique in ophthalmology for cataract removal, emerged as an advancement over the mid-20th-century standard method, extracapsular cataract extraction (ECCE)¹, wherein the entire cloudy lens was removed intact, preserving the outer capsule. Credited to American ophthalmologist Dr. Charles Kelman in the 1960s², phacoemulsification innovatively utilizes ultrasound to fragment the lens material, allowing for aspiration through a small incision, revolutionizing cataract surgery^{3,4}. Ongoing technological advancements in phacoemulsification machines, marked by superior ultrasound technology, advanced fluidics, and sophisticated control systems, have significantly enhanced the safety and precision of cataract surgery^{5,6}.

However, a prevalent complication associated with this technique is the potential for patients to undergo prolonged inflammation, resulting in elevated intraocular pressure, ocular irritation, and the development of cystoid macular edema, often necessitating additional medications or interventions⁷. Corticosteroids, widely employed in ophthalmology for years, exhibit anti-inflammatory and immunosuppressive attributes⁸, proving effective in managing various inflammatory eye conditions⁹. Typically administered as topical eye drops or injections, these steroids are prescribed postoperatively to control inflammation, with topical drops used for a specified duration. Intraocular steroid injections may also be employed intra-operatively or post-operatively, particularly in cases with elevated inflammation risk or pre-existing inflammatory conditions^{10,11}.

The findings of this study may contribute to the optimization of postoperative management strategies in cataract surgery, providing evidence-based guidance for the use of intra-cameral dexamethasone in routine clinical practice. Understanding the specific benefits and potential risks associated with this approach could lead to improved patient outcomes, enhanced surgical recovery, and a more tailored and effective postoperative care protocol for individuals undergoing uncomplicated phacoemulsification.

Department of Ophthalmology, D. G. Khan Medical College, D. G. Khan.

Correspondence: Dr. Muhammad Saad Ullah, Assistant Professor of Ophthalmology, D. G. Khan Medical College.

Contact No: 0333-6052117

Email: drmsaad@gmail.com

Received: September, 2023

Accepted: November, 2023

Printed: January, 2024

METHODS

Study involved 80 patients having age between 55-80 years and admitted at hospital for phacoemulsification under topical anesthesia. Study was started after approval from chairman authorized committee and consent was obtained from patients after detailed description of procedure and study. Demographic, pre-operative examination findings and post-operative findings at day 1 and day 7 were recorded on predesigned performa. Furthermore, before 1 week of procedure biometry was performed. Patients diagnosed as grade II and IV cataract from nuclear sclerosis, intraocular pressure in normal range and 6/12 or above visual acuity were enrolled in the study. Patients with history of previous ocular surgery, recently using non-steroidal anti-inflammatory drugs, having additional disease of eye, steroid susceptibility and use of immunosuppressant drugs were excluded from the study.

One hour before intervention 0.5% Proparacaine Hydrochloride solution was administered to the pointed eye. After establishment of standard anesthesia phacoemulsification was done with soft intraocular acrylic lens implantation. At the end of surgery, injection of intracameral dexamethasone sodium phosphate was given in group 1 patients at 0.4mg/0.1ml concentration. In group 2, equal volume of normal saline was injected as control. All procedures were performed by same ophthalmic surgeon and uneventful surgeries were projected. Post operative steroids and antibiotics were given as per ward protocol.

Post-operative examination was done at day 1 and 7 and outcomes were assessed which include intraocular pressure (IOP) that was measured on Goldmann Applanation Tonometer, visual acuity was measured on Snellen chart. Anterior segment was examined on slit lamp. Anterior chamber flare grading was assessed by using scoring system of SUN Working Group Slit Lamp Grading Scheme. SPSS version 27 was used for data analysis. After application of significance test p value below 0.05 was considered significant.

RESULTS

In group 1, age of patients was 62.73±6.28 years and in group 2 64.37±6.09 years. Regarding gender group 1 comprised 11 (27.5%) females and 29 (72.5%) males and in group 2, 13 (32.5%) females and 27 (67.5%) males. The mean pre-operative visual acuity was in group 1 was 0.68±0.21, at 1st post-operative day 0.29±0.09 and it was 0.22±0.07 at 7th day. Similarly visual acuity in group 2 was 0.76±0.18, 0.35±0.08, and 0.23±0.10 in pre-operative, 1st post-operative day and 7th post-operative respectively. Notably, the visual acuity at these time points was comparable between the two groups and showed no statistically significant differences (Table I).

The anterior chamber flare scores in group 1 at day was not found in any patient in 3 (7.5%) patients, in 16 (40.0%) patients it was faint, in 16 (40.0%) patients it was moderate, intensive in 3 (7.5%) patients and marked in 2 (5.0%) patients. In Group II, the corresponding scores showing zero results were 2.5% or 1 patient, faint results were found in 27.5% or 11 patients and moderate results in 57.5% or 23 patients, marked in 10.0% or 4 patients, and intensive in 2.5% or 1 patient. The difference in anterior chamber flare scores at day 1 between the two groups was (p>0.050). At day 7, the anterior chamber flare scores in Group I included none in 28 (70.0%) patients, faint in 9 (22.5%) patients, moderate in 2 (5.0%) patients, marked in 1 (2.5%) patient, with no patients exhibiting intensive scores. In Group II, the corresponding scores were none in 12 (30.0%) patients, faint in 23 (57.5%) patients, and moderate in 5 (12.5%) patients. The difference in anterior chamber flare scores at day 7 between the two groups was found to be p<0.050 according to Table II. At 1st post-operative day mean IOP in group 1 was 14.75±2.55 and in group 2 was 17.85±2.22. Similarly, 7th post-operative day mean IOP in group 1 was 13.18±2.14 and in group 2 it was 12.44±1.74. In comparison pre operative IOP was in group 1 and 2 was 16.19±2.54 and 12.44±1.74 respectively. However, the observed differences in mean IOP between the two groups showing p>0.050, as indicated in Table III.

Table No. 1: Baseline characteristics and demographics

Characteristic	Group 1	Group 2	p-value
Age (years)	62.73±6.28	64.37±6.09	0.401
Gender			
Male	29 (72.5)	27 (67.5)	0.626
Female	11 (27.5)	13 (32.5)	
Visual Acuity			
Pre-operative	0.68±0.21	0.76±0.18	0.077
1 st Post-operative day	0.29±0.09	0.35±0.08	0.010
7 th Post-operative day	0.22±0.07	0.23±0.10	0.572

Table No. 2: Anterior chamber cells and flare score

Grading	Group 1	Group 2	p-value
	At Day 1		
None	3 (7.5)	1 (2.5)	0.303
Faint	16 (40.0)	11 (27.5)	
Moderate	16 (40.0)	23 (57.5)	
Marked	2 (5.0)	4 (10.0)	
Intensive	3 (7.5)	1 (2.5)	
At Day 7			
None	28 (70.0)	12 (30.0)	0.002
Faint	9 (22.5)	23 (57.5)	
Moderate	2 (5.0)	5 (12.5)	
Marked	1 (2.5)	0 (0.0)	
Intensive	0 (0.0)	0 (0.0)	

Table No. 3: Intraocular Pressure parameters of the study groups

Intra Ocular Pressure (mmHg)	Group 1	Group 2	p-value
Pre-operative	16.19±2.54	17.85±2.22	0.250
1 st Post-operative day	14.75±2.55	14.54±3.81	0.815
7 th Post-operative day	13.18±2.14	12.44±1.74	0.098

DISCUSSION

Due to the historically linked undesirable side effects of cataract formation and elevated intraocular pressure associated with intraocular triamcinolone in phakic eyes, we have chosen to employ dexamethasone as an alternative for treating eye inflammation, addressing relevance about triamcinolone crystalline nature and its potential impact on intraocular pressure¹².

The mean age of patients was 62.73 ± 6.28 years in Group 1, while in Group 2, it was 64.37 ± 6.09 years. A previous study by Jan et al¹³ found a comparable mean age of 71 ± 9.4 years in Group 1 and 69.8 ± 10.5 years in Group 2. Additionally, El-Haddad et al¹⁴ found non-significant effect of intracameral triamcinolone on IOP. But, anti-inflammatory outcomes were highly effective. Contrast observations were reported by Shaheen et al¹⁵, that topical dexamethasone and intracameral triamcinolone have similar effectiveness when used following phacoemulsification procedure. Another study was conducted by Elkhodary et al¹⁶ and reported that intracameral triamcinolone utilization have much better outcomes on post-operative outcomes following phacoemulsification.

In the present study a statistically significant difference was observed between anterior chamber cells and flare at 7th post-operative day as p value <0.05 . But regarding intraocular pressure and visual acuity this difference was not statistically significant p value >0.05 . These findings are in concordance with findings of study conducted by Albialy et al¹⁷ reporting significant impact of anterior chamber cells and flare but insignificant regarding IOP. Manzoor et al¹⁸ highlighted anterior chamber reaction as a predominant factor in their findings. Conversely, Gungor et al¹⁹ observed no significant difference in anterior chamber cells and flare between intra-cameral dexamethasone and topical steroid formulation. Interestingly, our study concurs with findings of Gungor et al regarding IOP and visual acuity. Tan et al²⁰ conducted a study and on comparison of topical drops of dexamethasone and intracameral dexamethasone but no difference was observed

regarding outcomes of visual acuity and post-operative inflammation.

Another study was conducted on pediatric population by Khan et al²¹, in that study 50% of patients were administered dexamethasone intracameral and other half were administered sub-conjunctival dexamethasone. At the end of study variation was observed regarding ocular inflammation as higher frequency 26.7% was observed in group of sub-conjunctival patients and intracameral administration observed 6.7% inflammation.

Limitations: If the study is conducted at a single center, it might lack external validity. Different healthcare settings, patient populations, and surgical practices in other centers may influence the generalizability of the findings.

Recommendations: Stimulate further research on the long-term effects and cost-effectiveness of intracameral dexamethasone. Provide evidence for the integration of intra-cameral dexamethasone in routine phacoemulsification procedures to enhance postoperative outcomes.

CONCLUSION

Intraoperative inflammation following phacoemulsification surgery can be effectively managed by administering intra-cameral dexamethasone injection, resulting in decreased flare and anterior chamber cells, with comparable impacts on IOP and visual acuity when compared to the use of topical steroids.

Author's Contribution:

Concept & Design of Study: Muhammad Saad Ullah
Drafting: Muhammad Usama

Rahim, Hassan Shoaib

Data Analysis: Hassan Shoaib

Revisiting Critically: Muhammad Saad Ullah,
Muhammad Usama
Rahim

Final Approval of version: Muhammad Saad Ullah

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.49 DGKMC dated 03.03.2022

REFERENCES

- Moussa G, Pandey P, Panthagani J, Kutubi M, Masood I. Phacoemulsification with Trabecular micro-bypass stents in complex and moderate to advanced glaucoma: 3 year outcomes. *J Clin Ophthalmol* 2020;4(1):222.
- Yagev R, Khatib N, Barrett C, Lior Y, Lifshitz T, Tsumi E. Intraocular lens implantation as an isolated risk factor for secondary glaucoma in

- pediatric patients. *Can J Ophthalmol* 2019;54(5):621-625.
3. Elgazzar AF, Ahmed REH. Anti-inflammatory effects of intracameral triamcinolone acetonide injection versus topical dexamethasone after cataract surgery: a prospective comparative study. *Glob J Curr Res* 2019;6(4):149-153.
 4. Chang DT, Herceg MC, Bilonick RA, Camejo L, Schuman JS, Noecker RJ. Intracameral dexamethasone reduces inflammation on the first postoperative day after cataract surgery in eyes with and without glaucoma. *Clin Ophthalmol* 2009;3:345-55.
 5. Li A, He Q, Wei L, Chen Y, He S, Zhang Q, et al. Comparison of visual acuity between phacoemulsification and extracapsular cataract extraction: a systematic review and meta-analysis. *Ann Palliat Med* 2022;11(2):551-559.
 6. Ali F. Surgically induced astigmatism after phacoemulsification with clear corneal 3.2 mm incision while using superior versus temporal approach. *Med J South Punjab* 2022;3(1):21-3.
 7. Imdad T, Tanweer FS, Raza A. Effect of Dexamethasone versus Diclofenac Sodium after phacoemulsification with intraocular lens implantation. *JRMC* 2013;17: 254-256.
 8. Jamil AZ, Ahmed A, Mirza KA. Effect of intracameral use of dexamethasone on corneal endothelial cells. *J Coll Physicians Surg Pak* 2014;24(4):245-8.
 9. Nasir J, Javed A, Arshad O, Chatni MH. Spectral domain - optical coherence tomography findings in triple-a syndrome - a case series from Pakistan. *Pak J Med Sci* 2021;37(1):267-271.
 10. Sonmez K, Ozturk F. Complications of intravitreal triamcinolone acetonide for macular edema and predictive factors for intraocular pressure elevation. *Int J Ophthalmol* 2012;5(6):719-25.
 11. Aslan F, Öktem Ç. Does adjuvant intracameral triamcinolone acetonide increase the effectiveness of phacotrabeculectomy? A case-control study. *Cesk Slov Oftalmol* 2020;76(2):68-76.
 12. Daas A, Sherman T, Danieliute L, Goyal S, Amon A, Rodrigues I, et al. Phacoemulsification combined with micropulse cyclodiode laser in glaucoma patients: efficacy and safety. *Eye* 2022;36(11):2188-91.
 13. Jan S, Tahir MA, Faisal SM, Arain AU, Cheema A. effects of intra-cameral dexamethasone after uncomplicated phacoemulsification. *Pak J Ophthalmol* 2023;39(4):279-83.
 14. El-Haddad NS. A single dose intracameral triamcinolone injection following phacoemulsification. *Egypt J Hosp Med* 2021; 84(1):2667-70.
 15. Shaheen KH, Ullah MS, Hussain SA, Furqan A. Intracameral triamcinolone acetonide versus topical dexamethasone: a comparison of anti-inflammatory effects after phacoemulsification. *Cureus* 2020;12(4):e7592.
 16. Elkhodary NM, Hasan ZS, El-Din Mohamed NS. Comparison of anti-inflammatory effect between intracameral triamcinolone acetonide and topical dexamethasone after phacoemulsification. *Egypt J Hosp Med* 2023;90(1):601-6.
 17. Albialy HA, Wagih MM, Gouda DM. Safety and efficacy of intracameral injection of dexamethasone and moxifloxacin at the end of cataract surgery. *Zagazig University Med J* 2022;28(6.2):74-80.
 18. Manzoor A, Moin M. Comparison of antiinflammatory effect between intracameral triamcinolone acetonide and topical dexamethasone after phacoemulsification. *Pak J Ophthalmol* 2018;34(1):19-24.
 19. Gungor SG, Bulam B, Akman A, Colak M. Comparison of intracameral dexamethasone and intracameral triamcinolone acetonide injection at the end of phacoemulsification surgery. *Ind J Ophthalmol* 2014;62(8):861-64.
 20. Tan DT, Chee SP, Lim L, Theng J, Van Ede M. Randomized clinical trial of Surodex steroid drug delivery system for cataract surgery: anterior versus posterior placement of two Surodex in the eye. *Ophthalmol* 2001;108(12):2172-2181.
 21. Khan MA, Ahmad F, Ahsen M, Anwar F, Shahzad MA, Bhatti SA. Examine the comparison of efficacy between intracameral and sub conjunctival dexamethasone for the prevention of postoperative inflammation in patients with cataract surgery. *Pak J Med Health Sci* 2022;16(05):1435.

Anatomical Variation in the Location of Mandibular Foramen with Age Using Cone Beam Computed Tomography

Anatomical Variation of Mandibular Foramen Using Cone Beam CT

Asma Sattar¹, Naheed Imran¹, Muhammad Ishfaq², Sana Arbab¹, Munawar Aziz Khattak¹ and Imran Khattak¹

ABSTRACT

Objective: To use cone-beam computed tomography images to assess the mandibular foramen's location in relation to age.

Study Design: Cross-sectional retrospective study examined the hospital records.

Place and Duration of Study: This study was conducted at the Radiology Department of Khyber College of Dentistry (KCD), in Peshawar, Pakistan 4th November 2021 to 3rd May 2022.

Methods: 1000 CBCT radiographs from patients treated over a two-year period were examined in the initial radio-anatomical investigation. The shortest distance between the mandibular foramen (MF) and Point A, Point P, Point MI, Point MN, and Point O were measured. Ratios were also computed to ascertain the MF's location in relation to these anatomical landmarks. For all data statistical analysis, a significance level of $P \leq 0.05$ was used.

Results: 134 mandibular foramens are associated with people between the ages of 15 and 70, with an average age \pm (SD) of 39.81 ± 14.71 years. The measured mean distances were 17.29, 12.54, 18.70, and 32.43 from the mandibular foramen to Point A, Point P, Point MI, Point MN, and Point O respectively. The MF was found about 3.65 mm above point O. The average measurement between point A and point P was 49.36 mm, whereas the average measurement between point MI and point MN was 50.60 mm. The computed ratios for AMF/AP and MIMF/MIMN were 0.58 and 0.37 mm. The investigation's findings demonstrated that the location of the mandibular foramen varied statistically significantly among age groups.

Conclusion: The mandibular foramen's location varied dramatically with age, according to the study's findings.

Key Words: Mandibular Foramen, Orthognathic, Cone Beam Computed Tomography

Citation of article: Sattar A, Imran N, Ishfaq M, Arbab S, Khattak MA, Khattak I. Anatomical Variation in the Location of Mandibular Foramen with Age Using Cone Beam Computed Tomography. Med Forum 2024;35(1):18-22. doi:10.60110/medforum.350104.

INTRODUCTION

On the inside surface of the mandibular ramus, there is an uneven opening known as the mandibular foramen.¹ The key area for the inferior dental nerve block, which offers regional anaesthetic for various surgical operations in the lower jaw, is where the IAN invades the MF². The location of the MF varies greatly depending on the population and can change with age even in the same person on both sides³.

The identification of the mandibular foramen and IAN is required prior to the osteotomy of the ramus of the

mandible, and these procedures should be performed with caution to avoid injury². It is very common for IAN block to fail or for ramus to fracture in such orthognathic procedures and they are most commonly caused through being uncertain regarding the precise location of the MF in different age groups, races, or ethnicities. The IAN block's predicted failure rate is between 5 & 15%⁴ and 15 to 20%⁵. According to⁶ this failure rate could be as high as 45%. Failure of IAN block might occur due to lack of definite anatomic landmarks, anatomical differences such as mandibular foramen located superior or inferior to its normal position; and improper anesthesia method that may be due to limited mouth opening, a needle placed too anterior or posterior to the normal location⁷.

The MF location have been determined by different authors using different methods such as, dried human mandibles⁸, panoramic radiographs^{9,10}, CT scan⁷ and CBCT to locate mandibular foramen^{2,6,11-13}. CBCT offers better accurate localization of many anatomical features and there is less distortion of image as compared to plain radiograph. Moreover, it has higher accuracy, more resolution, less scan time, and decreased radiation dose in comparison with typical CT imaging².

¹. Department of Oral Biology, Peshawar Dental College, Peshawar, KP, Pakistan.

². Department of Oral & Maxillofacial Surgery, Peshawar Dental College, Peshawar, KP, Pakistan.

Correspondence: Dr. Asma Sattar, Department of Oral Biology, Peshawar Dental College, Peshawar, KP, Pakistan.
Contact No: 0318-9555917
Email: dr.asmasattar1@gmail.com

Received: July, 2023

Accepted: September, 2023

Printed: January, 2024

To our knowledge till date no studies have been carried out among the local population on this subject. The current investigation aims to pinpoint the exact position of the MF. Dental surgeons may use the findings of this study to detect the mandibular foramen, which will enable them to select an easy-to-reach target area for the IANB, lowering the risk that it will fail in the majority of patients. This study's findings will also define a safe zone for extra-oral mandibular ramus osteotomy treatments, reducing the possibility of harm to the inferior alveolar nerve in individuals having these procedures.

METHODS

The Institutional Review Board (IRB) of the prime foundation gave its approval to this cross-sectional retrospective study, on 10 Sep. 2021 (Approval no: Prime/IRB/2021-358). Ethical approval for data collection was granted by RRB-KCD vide notification No. 3065/RRB/KCD dated 3rd November 2021. The study was carried out in Peshawar. CBCT images were collected from Khyber College of Dentistry's radiology department (KCD). Khyber College of Dentistry (KCD) is a referral hospital in Peshawar, and its health services are available to all patients from KPK of varying socioeconomic backgrounds. In the supervision of an oral and maxillofacial surgeon, the observer interpreted CBCT images. The Radiology Department of Khyber College of Dentistry (KCD) presently has the CBCT radiographs studied in this study. Planmeca Romexis software was used to import CBCT images into the computer CBCT pictures of individuals between the ages of 15 and 70 had been included in this investigation, however radiographs of patients with asymmetrical faces, pathological lesions in the mandibular ramus, or those who had bilaterally absent or malpositioned mandibular first molars were not included. The senior radiology technician obtained the CBCT images at KCD in accordance with the manufacturer guidelines and a stringent, standardised scanning methodology. These radiographs were produced using a cone beam computed tomography scanner with an exposure period of 9 seconds and a voxel size of 400 μ m approx. depending on the FOV. To prevent inter-observer variations, the same examiner evaluated all the cone beam computed tomography images. Additionally, they were assessed under standard viewing conditions, which included enhancing radiograph clarity by modifying the brightness and opacity settings. The investigator got training to identify the MF and other mandibular landmarks prior to doing the radiographic examination, using a set of cone beam computed tomographic images that were not a part of the research. The calibration training procedure included hands-on discussion sessions and demonstrations of mandibular CBCT landmark identification techniques as well as demonstrations of

the steps involved in determining the distances between mandibular landmarks using the CBCT software. Axial, sagittal, cross-sectional and panoramic views were used to locate MF. The mandibular landmarks used in past studies and the location of the MF were measured using the software ruler to estimate their distances (in millimeters)^{2,6,11-13}.

Among the landmarks were the anterior border's deepest point on the ramus (A), the deepest point of ramus' posterior border (P), the mandibular first molar's occlusal plane (O), the most superior point of curvature of the mandibular notch (MN), the most inferior point of mandibular incisura (MI),

The subsequent calculations involved the following measurements and ratios:

- The shortest path AP: To ascertain the ramus's horizontal dimension between positions A and P.
- To find the height of the ramus, take the shortest MIMN distance (between points MI and MN).
- The MF's horizontal placement is described by the AMF/AP ratio.
- MIMF/MIMN ratio describes the vertical position of the MF.

Each participant's mandibular foramen was evaluated bilaterally, and its location as well as the patient's MRN number and age was noted on the proforma. Based on their ages, the participants were then separated into the following four groups (Table 1).

Statistical Assessment:

- The statistical analysis was performed using the Statistical Package for Social Sciences (SPSS), version 20.
- To analyse data, descriptive statistics were employed.
- In order to ascertain whether the age groups differed noticeably from one another, a one-way ANOVA test was employed.
- $P \leq 0.05$ was designated as the statistical significance level for the test.

RESULTS

All Cone beam computed tomographic images of patients treated over a 2-year period were investigated in this radioanatomical investigation. A total of 1000 cone beam computed tomographic images were initially evaluated, and of them, 100 CBCTs met the requirements for inclusion in the research. The mandibular foramen of each CBCT image was evaluated mean age \pm (SD) of patients were 39.81 ± 14.71 years (range =15-70 years).

Table 1: Age groups included in this study.

Groups	Age
Young adults	From 15 to 25 years
Adults	From 26 to 40 years
Middle age	From 41 to 60 years
Elderly	From 61 to 70 years

To determine if age-related changes were statistically significant, a one-way ANOVA test was used. (P ≤ 0.05).

Young adults (15-25 years old) had the lowest MIMF, MNMF, OMF, and MIMN values (Table: 2).

PMF and AP values were lowest in adults (26-40 years) while this group showed highest values of MIMF/MIMN ratio (Table: 2).

Middle age group (41-60 years) showed lowest values of AMF and AMF/AP ratio (Table: 2).

AMF, PMF, MIMF, MNMF, OMF, AP, MIMN AND AMF/AP ratio were higher in elderly patients (61-70 years) while MIMF/MIMN ratio decreased in this age (Table: 2).

Values of MIMF, MNMF, MIMN AND MIMF/MIMN ratios differed significantly among different age groups. Nevertheless, there was no age-related statistically significant variation in the levels of AMF, PMF, OMF, AP, or AMF/AP ratio (Table: 2).

Table No.2: The average distance between MF and numerous mandibular landmarks among different age groups

Age Groups	N	Mean ± Std. Deviation (mm)	P value*	
Distance from point A to point MF (A-MF)	15-25	112 (56%)	17.31±2.45	0.262
	26-40	74 (37%)	17.36±2.93	
	41-60	12 (6%)	16.23±3.57	
	61-70	2 (1%)	20.07±1.78	
	Total	200	17.29±2.71	
Distance from point P to point MF (P-MF)	15-25	112 (56%)	12.50±2.57	0.097
	26-40	74 (37%)	12.29±2.62	
	41-60	12 (6%)	14.11±1.94	
	61-70	2 (1%)	14.47±2.22	
	Total	200	12.54±2.58	
Distance from point MI to point MF (MI-MF)	15-25	112 (56%)	18.17±3.41	0.046
	26-40	74 (37%)	19.13±3.81	
	41-60	12 (6%)	20.64±3.36	
	61-70	2 (1%)	21.28±4.97	
	Total	200	18.70±3.62	
Distance from point MN to point MF (MN-MF)	15-25	112 (56%)	31.53±5.06	0.034
	26-40	74 (37%)	33.49±5.21	
	41-60	12 (6%)	33.61±5.38	
	61-70	2 (1%)	36.94±4.86	
	Total	200	32.43±5.21	
Distance from point O (mandibular first molar's occlusal plane) to point MF (O-MF)	15-25	112 (56%)	3.39±2.62	0.290
	26-40	74 (37%)	4.00±2.60	
	41-60	12 (6%)	3.60±1.48	
	61-70	2 (1%)	5.62±2.73	
	Total	200	3.65±2.57	
Distance between points A and point P (AP)	15-25	112 (56%)	29.41±3.19	0.325
	26-40	74 (37%)	29.13±2.84	
	41-60	12 (6%)	29.76±2.42	
	61-70	2 (1%)	32.94±2.31	
	Total	200	29.36±3.02	
Distance between points MI and MN (MIMN)	15-25	112 (56%)	49.36±5.45	0.002
	26-40	74 (37%)	51.75±5.99	
	41-60	12 (6%)	53.87±6.94	
	61-70	2 (1%)	57.69±1.21	
	Total	200	50.60	
The AMF/AP ratio	15-25	112 (56%)	0.58±0.09	0.519
	26-40	74 (37%)	0.58±0.12	
	41-60	12 (6%)	0.54±0.09	
	61-70	2 (1%)	0.61±0.10	
	Total	200	0.58±0.10	
The MIMF/MIMN ratio	15-25	112 (56%)	0.37±0.07	0.003

	26-40	74 (37%)	0.38±0.09
	41-60	12 (6%)	0.38±0.04
	61-70	2 (1%)	0.16±0.21
	Total	200	0.37±0.08

Table No.3: The average distances between various mandibular landmarks and the MF.

The average distances between various mandibular landmarks and the MF.	MEAN (mm)
Distance from point A to point MF (A-MF)	17.29
Distance from point P to point MF (P-MF)	12.54
Distance from point MI to point MF (MI-MF)	18.70
Distance from point MN to point MF (MN-MF)	32.43
Distance from point O (mandibular first molar’s occlusal plane) to point MF (O-MF)	3.65
Distance between points A and point P (AP)	49.36
Distance between points MI and point MN (MIMN)	50.60
The AMF/AP ratio	0.58
The MIMF/MIMN ratio	0.37



Figure No.1: The shortest distance from the MF to numerous mandibular landmarks on panoramic view of CBCT.

DISCUSSION

The noninvasive way to locate the mandibular foramen precisely is by radiographs, which are essential in oral and maxillofacial surgery². The recommended radiographic method for accurately identifying and examining the mandibular foramen is cone beam computed tomography (CBCT) due to its several advantages over plain films. Many dentists have found CBCT's diagnostic ability to be beneficial. CBCT can express fine structures due to its small voxel size, and it has a lower radiation dose than a conventional multislice CT scan. It also requires less tube voltage and current than conventional CT¹¹.

According to the current investigation, the average distances from the MF to point A, point P, point MI, point MN, & point O were measured to be 17.29, 12.54, 18.70, 32.43, and 3.65 mm, respectively (Table: 3).

According to a study by² on Jordanians, the average distances between the MF and the ramus's anterior and posterior margins, the mandibular incisura, and the mandibular notch were 17.51mm, 13.16mm, 19.28mm, and 25.66mm respectively. The MF was situated 4.52 mm above the occlusal plane.²

The mean vertical height of the ramus was 50.60mm in this study (Table: 3). Nevertheless, this height was measured as 49.4mm in a study done by da Fontoura et al., in 2002.¹⁴

There were statistically significant differences in the current study in the values of the MIMF, MNMF, MIMN, and MIMF/MIMN ratios across the various age groups, but not for the AMF, PMF, OMF, AP, or AMF/AP ratios. The lowest values of MIMF, MNMF, OMF, and MIMN were found in young adults. Adults had the lowest PMF and AP levels, but they had the greatest MIMF/MIMN ratios. The AMF and AMF/AP ratio were lowest in the middle-aged group. Elderly patients had increased levels of AMF, PMF, MIMF, MNMF, OMF, AP, MIMN, AND AMF/AP ratio, whereas MIMF/MIMN ratio decreased at this age (Table: 2). In a research done on adults under the age of 40, it was observed that the MF's position in regard to landmarks did not vary with age.¹¹

The MF was 3.65 mm above the occlusal plane in this study. Previous research found the MF to be 2.5-3.6 mm above the occlusal plane of the molars.¹⁵, and no statistically significant changes in the MF's location with advancing age were seen. Adults had the MF 4.2 mm above the occlusal plane.

CONCLUSION

The location of the mandibular foramen was determined using landmarks from CBCT scans. The location of the mandibular foramen varied dramatically with age, according to the research findings.

Author’s Contribution:

Concept & Design of Study: Asma Sattar
 Drafting: Naheed Imran, Muhammad Ishfaq
 Data Analysis: Sana Arbab, Munawar Aziz Khattak, Imran Khattak
 Revisiting Critically: Naheed Imran, Asma Sattar
 Final Approval of version: Asma Sattar

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No. 3065/RRB/KCD dated 03.11.2021.

REFERENCES

1. Chen CM, Lee HN, Liang SW, Hsu KJ. Morphological Study of the Mandibular Lingula and Antilingula by Cone-Beam Computed Tomography. *Bioengineering* 2023;10(2):170.
2. Al-Shayyab MH. A simple method to locate mandibular foramen with cone-beam computed tomography and its relevance to oral and maxillofacial surgery: a radio-anatomical study. *Surgical Radiologic Anat* 2018;40(6):625–34.
3. Vathariparambath N, Krishnamurthy NH, Chikkanarasaiah N. A Cone Beam Computed Tomographic Study on the Location of Mandibular and Mental Foramen in Indian Pediatric Population. *Int J Clin Pediatr Dent* 2022; 15(4):422–7.
4. Sarfaraz I, Pascoal S, Macedo JP, Salgado A, Rasheed D, Pereira J. Anesthetic efficacy of Gow-Gates versus inferior alveolar nerve block for irreversible pulpitis: a systematic quantitative review. *J Dent Anesth Pain Med* 2021;21(4): 269–82.
5. Lasemi E, Motamedi MHK, Qaranizade K, Hassani Z, Lasemi R. Modification in implementation of the inferior alveolar nerve block technique. In *Review*; 2022 May [cited 2023 Oct 19]. Available from: <https://www.researchsquare.com/article/rs-1599913/v1>
6. Sanwatsarkar G, Agarwal R, Hiremath H, Kulkarni S, Agarwal J, Hiremath V. Mandibular foramen location and its implication to the inferior alveolar nerve block: A retrospective study. *Endodontol* 2023;35(2):113.
7. Feuerstein D, Costa-Mendes L, Esclassan R, Marty M, Vaysse F, Noirrit E. The mandibular plane: a stable reference to localize the mandibular foramen, even during growth. *Oral Radiol* 2020;36(1):69–79.
8. Park JH, Jung HD, Kim HJ, Jung YS. Anatomical study of the location of the antilingula, lingula, and mandibular foramen for vertical ramus osteotomy. *Maxillofacial Plastic and Reconstructive Surgery*. 2018;40(1):1–6.
9. Krishnamurthy NH, Unnikrishnan S, Ramachandra JA, Arali V. Evaluation of relative position of mandibular foramen in children as a reference for inferior alveolar nerve block using orthopantomograph. *J Clin Diagnostic Res : JCDR*. 2017; 11(3):ZC71.
10. Kumar A, Bhagawati BT, Kumar N, Jyotsana K, Sharanamma B, Solanki H. Morphometric Analysis of Mental and Mandibular Foramen in Gender Determination among Sriranganganagar Population: A Cross-Sectional Digital Panoramic Study. *J Ind Acad Oral Med Radiol* 2023; 35(2):258.
11. Ahn BS, Oh SH, Heo CK, Kim GT, Choi YS, Hwang EH. Cone-beam computed tomography of mandibular foramen and lingula for mandibular anesthesia. *Imaging Science Dentistry* 2020; 50(2):125.
12. Mehanny M, Mansour S, Alzahrani A, Alquraishi L, Alhassoun R, AlSerwi RH. Morphometric localization of Mandibular Foramen in a sample of Adult Saudian Population Using Cone-Beam Computed Tomography. *Egypt Dent J* 2022; 68(2):1479–87.
13. Valizadeh S, Tayefi M, Ghomeishi M, Ahsaie MG, Amiri MJ. Assessment of Anatomical Location and Variation of Mandibular Foramen Using Cone-Beam Computed Tomography: A Cross-Sectional Study. *J Iranian Med Council* 2022 Dec 12 [cited 2023 Oct 19]; Available from: <https://publish.kne-publishing.com/index.php/JIMC/article/view/11334>.
14. da Fontoura RA, Vasconcellos HA, Campos AES. Morphologic basis for the intraoral vertical ramus osteotomy: anatomic and radiographic localization of the mandibular foramen. *J Oral Maxillofacial Surg* 2002;60(6):660–5.
15. Altunsoy M, Aglarci OS, Ok E, Nur BG, Gungor E, Colak M. Localization of the mandibular foramen of 8-18 years old children and youths with cone-beam computed tomography. *J Pediatr Dentistry* 2014;2(2).

Evaluation of Obstetric Anaesthesia and its Association with Maternal Outcomes in Women with Placenta Previa: A Cross-Sectional Study

Shandana Bawar, Qudsia Qazi and Syeda Sitwat Fatima

ABSTRACT

Objective: To determine the frequency of mode of anaesthesia (MOA). The secondary outcome was to determine an association between MOA, type 4 PP, previous CS and maternal outcomes.

Study Design: A retrospective cross-sectional study.

Place and Duration of Study: This study was conducted at the Department of Obstetrics and Gynecology (OBGYN), MTI Lady Reading Hospital Peshawar from January 2020 - December 2022.

Methods: It included women with singleton pregnancy with PP, after 28 weeks of gestation. Women with other causes of antepartum haemorrhage, previous myomectomy and medical disorders complicating pregnancy were excluded. Maternal outcomes included per-operative blood loss (POBL), per-operative RBC transfusion (POBT) and transfer to the critical care unit (TCCU) as mentioned in operative notes. Data was collected and analyzed by SPSS version 22.

Results: A total of 170 women were included in two years. MOA included General anaesthesia (GA) in 96 (56.5%) and Spinal anaesthesia (SA) in 74 (43.5%) cases. GA was frequently given in Emergency CS (EMCS), elective CS (ELCS) and type 4 PP. POBL of less than 1500ml dominated, POBT of less than or equal to 4 pints was found to be 143 (84.1%) while a large number of patients were managed in obstetrical wards 150 (88.2%) compared to HDU and ICU with a non-significant association.

Conclusion: GA was frequently adopted in our setup compared to SA, especially with the increasing severity of PP type and previous CS. Both GA and SA were safe with non-significant association with blood loss, RBC transfusion and critical care management.

Key Words: Placenta Previa, Mode of anaesthesia, per operative blood loss, Per operative blood transfusion, Critical care transfer

Citation of article: Bawar S, Qazi Q, Fatima SS. Evaluation of Obstetric Anaesthesia and its Association with Maternal Outcomes in Women with Placenta Previa: A Cross-Sectional Study. Med Forum 2024;35(1):23-27. doi:10.60110/medforum.350105.

INTRODUCTION

Placenta previa is an abnormal location of the placenta in the lower uterine segment, which is associated with grave maternal morbidity in terms of antepartum and per-operative haemorrhage, if not managed timely¹. Placenta previa is often graded into minor and major based on the distance of the lower edge of the placenta from internal os, types 1 and 2 comprising minor categories while types 3 and 4 constitute major placenta previa².

Department of Obstetrics and Gynecology, MTI, Lady Reading Hospital, Peshawar.

Correspondence: Syeda Sitwat Fatima, Assistant Professor of Obstetrics and Gynecology, MTI, Lady Reading Hospital, Peshawar.

Contact No: 0333 9301667

Email: smkakakhel@hotmail.com

Received: August, 2023

Accepted: November, 2023

Printed: January, 2024

The rising incidence of placenta previa and morbidly adherent placenta worldwide is attributed to escalated cesarean section rates, while previous curettage, manual removal of placenta, uterine surgery and assisted reproductive techniques are also thought to be the cause of placenta previa³. Women with placenta previa may present with painless genital tract bleeding, malpresentation, high presenting part or at times be asymptomatic, diagnosed incidentally on routine clinical examination or ultrasound⁴⁻⁶. Unpredictable antepartum haemorrhage in these women often leads to prolonged antenatal admissions in obstetrical wards, to avoid heavy bleeding episodes and related complications, that could be catastrophic to the mother's life, necessitating urgent delivery⁷. This mostly pertains to population who have long distances to travel to a health facility with needed facilities for PP. The mode of delivery is mainly cesarean section however vaginal delivery may occur in case of minor placenta previa⁸. Placenta previa due to its unforeseeable haemorrhage often ends in early delivery, adding neonatal sequelae related to prematurity in

addition to morbidity associated with PP⁹. CS and MOA, have always been under scrutiny, especially when it comes to PP. General anaesthesia was thought to be safer and quicker for women with placenta previa due to anticipated blood loss and chance of conversion of surgical procedure to hysterectomy with possible need of intensive care unit transfer¹⁰. But now advances in regional anaesthesia, have implicated its use to be equally safer for placenta previa¹¹. Several studies can be searched related to the mode of anaesthesia among patients with placenta previa. However, few can be found locally to understand the importance of anaesthesia and hemodynamic stability of these patients. We designed this study to evaluate the mode of anaesthesia in tertiary care that can help establish its safety and practice in obstetrics, thus devising evidence-based local protocols and guidelines for women with PP.

METHODS

This retrospective cross-sectional study was conducted in the department of obstetrics and gynaecology in a tertiary care hospital, MTI Lady Reading Hospital Peshawar between January 2020 to December 2022, after obtaining ethical approval from the hospital Institutional Review Board. This study included all the women with singleton pregnancies diagnosed with PP, clinically or by ultrasound during the antenatal period or CS after 28 weeks of gestation, irrespective of their previous CS status. Women with the local cause of genital tract bleeding, heavy show in labor, placental abruption, morbidly adherent placenta, pregnancy with anaemia, hypertensive disorders, diabetes with polyhydramnios, other medical disorders, and previous gynecological surgery(myomectomy etc.) on the uterus that could add into maternal outcomes were excluded. Retrospective analysis of hospital software and available history charts would determine these confounding factors. The operative notes determined the MOA and maternal outcomes to be studied. Maternal outcomes included per-operative blood loss(POBL), per-operative RBC transfusion(POBT) and transfer to critical care unit(TCCU). POBL was calculated by several abdominal gauze packs used during the operation and their conversion into millilitres as per hospital protocol. POBT was determined as per the number of bags mentioned in the charts. TCCU was divided into departmental OBGYN high dependency unit(HDU) and hospital intensive care unit(ICU) which was also mentioned in operation notes. Data was collected on a specialized proforma designed for the study from the clinical records of patients, maintaining confidentiality. Data was then transferred and analyzed on SPSS version 22. Frequencies and percentages were calculated for categorical variables like PP and its types, previous CS, and maternal outcomes. Mean and standard deviation were calculated for numerical

variables like maternal age etc. Chi-square test/Fisher's Exact test was applied to determine the association between mode of anaesthesia and previous CS, type of PP and maternal outcomes, with a p-value ≤ 0.05 was considered significant.

RESULTS

A two-year study included 191 patients with placenta previa. Women with morbidly adherent placenta were 21(10.9%) that were excluded from the study. Further analysis was done with a total of 170 patients with placenta previa. The mean age of participants was found to be 30 ±15 years. Primi gravida(first time pregnant) were 19(11.2%) , multigravida(pregnant 2-5times) were 110(64.7%) , grand multigravida (pregnant 6-8 times) were 30(17.6%) ,and great grand multigravida(pregnant >8times) were found to be 11(6.5%).Mean antenatal and postnatal stay was found to be 10 days± 14 days and 03±0.3 days, respectively, Table 01.Anterior PP was found in 61 (35.9%), posterior PP was found in 59(34.7%) while 50(29.4%)had major placenta previa covering internal OS.

Table No. 1: Demographic features of women with Placenta Previa

Maternal age	Years (mean)	SD
	30.36	±15.5
Stay obstetrical unit	Days	SD
Antenatal	10.0353	±14.28323
Postnatal	3.0824	±0.39857
Gravida status	Frequency(N)	Percentage (%)
primi gravida	19	11.2
2- 5 multi gravida	110	64.7
6-8	30	17.6
more than 8	11	6.5
Total	170	100

Table No. 2: Frequency of type of placenta previa and previous cesarean section among women with placenta previa

Type of placenta previa	Frequency(N)	Percentage (%)
1	02	1.2
2	20	11.8
3	48	28.2
4	100	58.8
Total	170	100.0

Previous cesarean section	Frequency(N)	Percentage (%)
none	148	87.1
1	16	9.4
2	04	2.4
3	01	0.6
4	01	0.6
Total	170	100

Among these 100 (58.8%) were type 4 PP followed by type 3,2 and 1 with frequency of 48(28.2%),20(11.8%), and 02(1.2%) respectively. About 61 (35.9%) had anterior placenta previa,59(34.7%) had posterior placenta previa and 50(29.4%)had major placenta previa covering internal os. About 148(87.1%) had no previous CS. Frequency of previous 1,2 CS was found to be 16(9.4%), 04(2.4%) and 01(0.6%) each for previous 3 and 4 CS, respectively, Table 02.

About 36(21.2%) women had a gestational age less than 34 weeks, 47(27.6%) had a gestational age between 34 and 37 weeks and 87(51.2%) had term gestation at the time of delivery. Emergency CS were 116(68.25%) and elective CS were 54 (31.7%) in number. MOA included GA in 96(56.5%) and SA in 74(43.5%) cases. During EMCS, GA was given in 71(74.05%)while 45(60.8%) had SA. Among elective CS, 29(39.2%) patients had SA while 24(25.0%) had GA, with a non-significant association p-value of 0.104. GA was more frequently given 64(66.7%) in patients with type 4PP than spinal anaesthesia 36(48.6%) bearing a significant association with a p-value of 0.038, however, a non-significant association of MOA with previous CS was determined with a p-value of 0.70, table 3.

Table No. 3: Frequency of type of placenta previa and previous cesarean section among women with Anaesthesia

Placenta previa type	Anaesthesia		p-value
	General	Spinal	
	Frequency %age	Frequency % age	
Type 1	01(50)	01(50)	0.038
Type 2	06(30.0)	14(70.0)	
Type 3	25(52.1)	23(47.9)	
Type 4	64(64.0)	36(36.0)	
Total	96(56.5)	74(43.5)	
Previous cesarian section			
None	82(55.4)	66(44.6)	0.70
Previous 1CS	09(56.3)	07(43.8)	
Previous 2CS	03(75.0)	01(25.0)	
Previous 3CS	01(100.0)	00(0.0)	
Previous 4CS	01(100.0)	00(0.0)	
Total	96(56.5)	74(43.5)	

Among the maternal outcomes POBL less than 1500ml dominated i.e 161(94.7%) while more than or equal to

1500 were 09(5.3%). POBT of less than or equal to 4 pints was found to be 143(84.1%) while more than 4 RBC units were transfused in 15(8.8%). A large number of patients were managed in obstetrical wards 150(88.2%), HDU care within the obstetric department was found to be 13(7.6%) and 07(4.1%) of women were managed in ICU. A non-significant association was seen for a mode of anaesthesia and blood loss, RBC transfusion, and critical care management, with Fischer exact test p-value of 0.30 for POBL, chi-square test p-value for POBT to be 0.139 and chi-square test p-value for TCCU to be 0.980.

DISCUSSION

Our study population was,170 women with PP, after excluding women with morbidly adherent placenta in 2 years. This suggest a large number of these patients in our set-up as compared to 276 patients found by, Ismail S in their study in 14 years. This may be attributed to different study designs and hospital settings, the former being a public sector hospital¹². PP was most frequently seen in multigravida 110(64.7%) in our study. A similar finding of PP dominance among multigravida 67(58.77%) was encountered in another regional retrospective study by Majeed T et al. However, a comparative more increase in multigravida in our study may be explained by the study duration and contraceptive practices of our country¹³. The most prevalent PP type was 4 in our study same to the findings of Grönvall M et al, who determined 129 cases of major PP. However, theses cases were found in a four year study while we determined major PP in two years. This contrast may be explained by different study design¹⁴. The anterior location of PP dominated our study with a frequency of 35.9%, opposite to the findings of an Iranian study which determined the anterior location of PP in 44.9% of their study population. The inclusion and exclusion criteria of both studies were different as our study included all the patients with PP but the referenced study included PP in patients with previous scar¹⁵.Oğlak SC et al in their study determined more planned CS 53.4%vs emergency CS 46.6%¹⁶. On the contrary, our emergency CS was higher than elective CS. The high emergency CS in our study may be due to the non-booked nature, and less or no antenatal visits of our population. Women with PP in our review suggested a percentage of 21.2% for very preterm and 27.6% for preterm deliveries. A Japanese study showed an overall frequency of premature delivery by 45.1%, in association with complete PP, quite similar to our findings¹⁷. However, we did not conduct the sub-analysis of gestational age with the type of PP like the Japanese study. The frequency of general anaesthesia (56.5%) was slightly higher in our study both for emergency and elective CS. A significant association of GA was determined with type 4 PP. Fan D et al also determined GA(63.76%)a frequent

finding, however, their percentage was quite high compared to us⁹. Further, it was emphasized that GA was seen more in emergency CS (26% vs. 38.6%, $P = .033$) by Fan D. On the contrary two different retrospective studies found neuraxial anaesthesia quite safe in even complicated cases like morbidly adherent placenta. They determined the conversion rate of neuraxial anaesthesia to GA increased with increase severity of placenta^{18,19}. It can be inferred that hemodynamic stability may be the reason for the consideration of regional anaesthesia in the referenced studies. A study by Alsammani Jr determined a high number of ICU admissions 48.27% in their study which is opposite to our findings²⁰. We could not discover any significant association between the MOA and POBL, POBT and TCCU, probably because PP in itself, primarily determine these outcomes. The choice of MOA usually depends upon the patient's condition, her vitals and stability. Anaesthesia may secondarily affect the fluid balance and further management of the patient. Lieu X et al suggested that although, regional anaesthesia had lesser operative time, less blood loss and lesser RBC transfusion, nevertheless, with increasing severity of placenta previa, the conversion rate to GA was also increased which was safe²¹. A multi centre study by Orbach-zinger potentiated the results of our study by determining the increases association of GA with complete PP²². A good sample size may potentiate our findings, but input from the anaesthesia department would have elaborated the findings in a better way. A combined obstetric and anaesthesia department prospective study comparing the MOA in PP may enhance the safety analysis of GA or RA, and address these limitations of our study.

CONCLUSION

General anesthesia was frequently adopted in our setup compared to regional anaesthesia. General anaesthesia was significantly associated with type 4 placenta previa. Both General and Spinal anaesthesia were safe with no significant association with blood loss, RBC transfusion and critical care management.

Author's Contribution:

Concept & Design of Study: Shandana Bawar
 Drafting: Qudsia Qazi, Syeda Sitwat Fatima
 Data Analysis: Syeda Sitwat Fatima
 Revisiting Critically: Shandana Bawar, Qudsia Qazi
 Final Approval of version: Shandana Bawar

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.05/LRH/MPI dated 02.01.2020

REFERENCES

- Gibbins KJ, Einerson BD, Varner MW, Silver RM. Placenta previa and maternal hemorrhagic morbidity. *J Matern neonatal Med.* 2018;31(4):494–9.
- Jauniaux E, Alfirevic Z, Bhide A, Belfort M, Burton G, Collins S, et al. Placenta Praevia and Placenta Accreta: Diagnosis and Management. *BJOG: An Int J Obstet Gynaecol* 2018;126(1):e1-e48.
- Silver RM. Abnormal Placentation: Placenta Previa, Vasa Previa, and Placenta Accreta. *Obstet Gynecol* 2015;126(3):654-668.
- Kumari U, Naniwal A, Rani V, et al. A Study of Clinical Characteristics, Demographic Characteristics, and Fetomaternal Outcomes in Cases of Placenta Previa: An Experience of a Tertiary Care Center. *Cureus* 2022;14(12):e32125.
- Jain V, Bos H, Bujold E. Guideline No. 402: Diagnosis and Management of Placenta Previa. *J Obstet Gynaecol Can* 2020;42(7):906-917.e1.
- Jauniaux E, Bhide A. Prenatal ultrasound diagnosis and outcome of placenta previa accreta after cesarean delivery: a systematic review and meta-analysis. *Am J Obstet Gynecol* 2017;217(1):27-36.
- Durukan H, Durukan ÖB, Yazıcı FG. Planned versus urgent deliveries in placenta previa: maternal, surgical and neonatal results. *Arch Gynecol Obstet* 2019;300:1541–9.
- Alouini S, Megier P, Fauconnier A, Huchon C, Fievet A et al. Diagnosis and management of placenta previa and low placental implantation. *J Matern Fetal Neonatal Med* 2020;33(19):3221-3226.
- Park HS, Cho HS. Anesth Management of massive hemorrhage in pregnant women with placenta previa. *Pain Med* 2020;15:409-416.
- Plaat F, Shonfeld A. Major obstetric haemorrhage. *BJA Educ* 2015;15(4):190–3.
- Fan D, Rao J, Lin D, Zhang H, et al. Anesthetic management in cesarean delivery of women with placenta previa: a retrospective cohort study. *BMC Anesthesiol* 2021;19;21(1):247.
- Ismail S, Rashid S. Caesarean Section for Placenta Previa: A Retrospective Cohort Study of Anaesthesia Techniques. *Turkish J Anaesthesiol Reanim* 2023;51(1):30.
- Majeed T, Waheed F, Mahmood Z, et al. Frequency of placenta previa in previously scarred and non scarred uterus. *Pak J Med Sci* 2015;31(2): 360–363.
- Grönvall M, Stefanovic V, Paavonen J, et al. Does it make a difference? *Placenta* 2019;85:9-14.
- Nankali A, Keshavarzi F, Shajari A, Daeichin S. Frequency of placenta previa and maternal

- morbidity associated with previous cesarean delivery. *Open J Obstet Gynecol* 2014;4(14):903.
16. Oğlak SC, Ölmez F, Tunç Ş. Evaluation of antepartum factors for predicting the risk of emergency cesarean delivery in pregnancies complicated with placenta previa. *Ochsner J* 2022;22(2):146–53.
 17. Sekiguchi A, Nakai A, Kawabata I, Hayashi M, Takeshita T. Type and location of placenta previa affect preterm delivery risk related to antepartum hemorrhage. *Int J Med Sci* 2013;10(12):1683.
 18. John CM, Farber, Michaela K et al. Neuraxial Anesthesia During Cesarean Delivery for Placenta Previa With Suspected Morbidly Adherent Placenta: A Retrospective Analysis. *Anesthesia & Analgesia* 2018;127(4):930-938.
 19. Li P, Liu X, Li X, Wei X, Liao J. Clinical outcomes and anesthetic management of pregnancies with placenta previa and suspicion for placenta accreta undergoing intraoperative abdominal aortic balloon occlusion during cesarean section. *BMC Anesthesiol* 2020;20(1):1–9.
 20. Alsammani Jr MA, Nasralla K. Fetal and Maternal Outcomes in Women with Major Placenta Previa Among Sudanese Women: A Prospective Cross-Sectional Study. *Cureus* 2021;13(4): e14467.
 21. Liu X, Zhu Y, Ke D, Liu D, Zhu Z. Mode of anesthesia for cesarean delivery with pernicious placenta previa—a retrospective study. *Ginekol Pol* 2020;91(2):91–4.
 22. Orbach-Zinger S, Weiniger FC, Aviram A et al. Anesthesia management of complete versus incomplete placenta previa: a retrospective cohort study. *J Matern Fetal Neonatal Med* 2018; 31(9):1171-1176.

The Importance of Immunohistochemical Evaluation of Ki67 in Detecting Early Malignant Changes in Colorectal Adenomatous Polyps

Evaluation of Ki67 in Detecting Early Malignant Changes in Colorectal Adenomatous Polyps

Sabika Batool¹, Talat Mirza¹, Fouzia Lateef² and Sobia Hassan¹

ABSTRACT

Objective: To assess the prevalent clinicopathological parameters of Adenomatous polyps and association of ki67 with them in order to elucidate its significance as potent marker in determining early malignant changes

Study Design: Retrospective study

Place and Duration of Study: This study was conducted at the Histopathology department and Molecular diagnostic Research laboratory (MDRL), Dr. Ziauddin University and Hospital, Karachi from November 2021 to November 2023.

Methods: A total of 55 colorectal adenomatous polyps and clinical data was retrieved. Fresh frozen plasma sections were stained under Hematoxylin and eosin stain and further analyzed with KI67 IHC stain. A P-value of less than 0.05 was deemed statistically significant.

Results: Among all 55 cases, 12 (21.8%) cases had age <50 years and 43 (78.2%) cases had age >50 years. Males predilection was observed. Most common clinical symptom was bleeding per rectum in 35 (63.64%) cases, followed by weight loss and chronic diarrhea, anemia, chronic abdominal pain, constipation while one case was accidentally reported on routine endoscopy (1.81%). 41 polyps (74.54%) were <20mm and 14 polyps (25.45%) were >20mm. Most common site was rectum (41.8%), followed by ascending and sigmoid colon, descending colon and transverse colon. Strong association was observed between size of colorectal polyps and grade of dysplasia with ki67 score.

Conclusion: When identifying the individuals with colorectal adenomatous polyps who require close surveillance in follow-up, size of polyp, high grade dysplasia and significant positive immunohistochemistry markers of Ki-67 may be useful criteria. This shall set a target population for screening of pre malignant changes.

Key Words: Dysplasia, Adenomas, ki-67 score

Citation of article: Batool S, Mirza T, Lateef F, Hassan S. The Importance of Immunohistochemical Evaluation of Ki67 in Detecting Early Malignant Changes in Colorectal Adenomatous Polyps. Med Forum 2024;35(1):28-32. doi:10.60110/medforum.350106.

INTRODUCTION

Intraepithelial pedunculated pre-neoplastic lesions are known as colorectal adenomas. By 50, they are found in almost half of adults and serve as the primary precursors of most cancers in this organ.⁽¹⁾ Adenomas are becoming more common in Asian population due to increased adaptation of Western diets and lifestyles. The epithelial dysplasia that is present in colorectal adenomas is a characteristic.

¹. Department of Pathology / Histopathology², Ziauddin Medical University, Karachi.

Correspondence: Prof. Dr. Talat Mirza, Research Department, Ziauddin University, 4/B Shahrah-e-Ghalib Rd, Block 6 Clifton, Karachi.

Contact No: 0333-2184453

Email: deanresearch@zu.edu.pk

Received: December, 2023

Accepted: December, 2023

Printed: January, 2024

The frequency of colorectal adenomas and colorectal carcinoma are correlated which can link the risk of recurrence and the emergence of colorectal malignancies to adenomas. 15% of all adenomas larger than 1 cm are predicted to develop into carcinomas within ten years of initial detection.⁽²⁾

Colorectal polyps are intraepithelial neoplasms growing through from the wall of the colon and rectum to its origin can be found in different sizes ranging from small polyps to large ones. Generally, colon polyps are more founded pathology and commonly seen over 50 aged patients and are not only the malignant pathology but more the precursor of malignant neoplasms, where the polyps have a high likelihood of cancer. Cancer normally develops in around 5% of adenomatous polyps.^(1,2) Colorectal polyps are intraepithelial neoplasms growing through from the wall of the colon and rectum to its origin can be found in different sizes ranging from small polyps to large ones. Generally, colon polyps are more founded pathology and commonly seen over 50 aged patients and are not only the malignant pathology but

more the precursor of malignant neoplasms, where the polyps have a high likelihood of cancer. Cancer normally develops in around 5% of adenomatous polyps.^(1,2) Colorectal polyps are intraepithelial neoplasms growing through from the wall of the colon and rectum to its origin can be found in different sizes ranging from small polyps to large ones. Generally, colon polyps are more founded pathology and commonly seen over 50 aged patients and are not only the malignant pathology but more the precursor of malignant neoplasms, where the polyps have a high likelihood of cancer. Cancer normally develops in around 5% of adenomatous polyps.^(1,2) Colorectal polyps are intraepithelial neoplasms growing through from the wall of the colon and rectum to its origin can be found in different sizes ranging from small polyps to large ones. Generally, colon polyps are more founded pathology and commonly seen over 50 aged patients and are not only the malignant pathology but more the precursor of malignant neoplasms, where the polyps have a high likelihood of cancer. Cancer normally develops in around 5% of adenomatous polyps.^(1,2) Colorectal polyps, which can range in size from tiny to large intraepithelial neoplasms, grow through the colon's and the rectal mucosal wall.⁽³⁾ In Colon, polyps which are more commonly found in patients over 50, are often a precursor to malignant neoplasms. Approximately 5% of adenomatous polyps typically develop cancer.⁽⁴⁾ Ki-67 is frequently used in standard clinical work^(5,6) and has been researched in connection with the onset and evolution of human colorectal cancer⁽¹⁾. It was regarded as a crucial predictor of human colorectal cancer.⁽¹⁾ A previous study found that the percentage of tumor cells positive for Ki-67 correlates inversely with overall survival, and that overexpression of Ki-67 in colorectal cancer (CRC) is linked to a worse prognosis.⁽⁵⁾ Some studies, though, have not been able to prove its prognostic significance.⁽⁴⁾ Although some researchers linked higher Ki-67 expression to a worse prognosis, others found that higher Ki-67 expression was associated with a better prognosis⁽⁷⁾. Higher Ki-67 expression has been reported by Melling et al.⁽⁸⁾ as an independent predictive marker in human colorectal cancer, despite the fact that its prognostic value is still debatable⁽⁷⁾. Ki-67 protein is an additional immunohistochemical marker used to identify proliferating cells. With the exception of the G0 phase, it manifests in every stage of the cellular cycle. Ki-67 is a nuclear protein as well as a nucleolus protein⁽⁸⁾ Ki67 protein is continually lacking in dormant cells and is undetectable during DNA repair processes.⁽⁹⁾ Because of the nucleus's significant function in the upkeep or control of the cell cycle, the presence of the Ki-67 antigen is thus exclusively associated with the cell cycle.⁽¹⁰⁾ Ki67 has been used by numerous researchers as a backup indicator to track the

proliferation activity of tumor cells in different systems⁽¹¹⁾.

For the purpose of early diagnosing and predicting the prognosis of cancer, various biomarkers linked to growth factors, angiogenesis, tumour suppressor genes, oncogenes, and proliferating cells factors are employed. Because of its excellent sensitivity, Ki67 is widely used to evaluate the rate at which cancer cells proliferate.⁽¹²⁾ Aim of current study was to examine the Immunohistochemical activity of Ki67 alongwith its correlation to size of polyp and grades of dysplasia in colorectal Adenomatous polyps in order to elucidate its significance as marker of early pre malignant changes.

METHODS

This retrospective study was conducted at Histopathology department and Molecular diagnostic Research laboratory (MDRL), Dr. Ziauddin University and Hospital, Karachi. 55 polypectomy specimens from patients in the form of Fresh Frozen Plasma Embedded tissues diagnosed as colorectal Adenomatous polyps were included.

Preparation of H&E sections: Hematoxylin and Eosin was used to stain paraffin-embedded sections from each colorectal polyp according to proper H&E staining protocols⁽¹³⁾

Preparation of IHC slides (immunohistochemical analysis): In order to establish a clear connection between the presence of protein ki67, and dysplasia in neoplastic tissues, IHC ki67 interventions have been performed on all adenomatous polyps exhibiting dysplasia following the IHC staining protocols⁽¹⁴⁾

Ki67 scoring technique: Without introducing the primary antibody (Ki-67) to the specimen, a negative control was employed.

Ki-67 expression counted in three high power fields (40X). Fraction of positive cells was calculated from atleast 500 tumor cells from the region where positivity was obvious and highlighted. 15 % or more Tumors cell staining positive was considered high ki67 score. Less than 15% tumors cells when stained positive were considered as low ki67 score⁽¹⁵⁾

Data analysis was carried out by SPSS 24.0. Chi-square test was performed for association between two variables. P value <0.05 was deemed statistically significant.

RESULTS

Clinicopathological parameters of patients included in study are expressed in table 1. 10 out of 21 polyps with mild dysplasia showed low ki67 score and 11 showed high ki67 score, all 3 polyps with moderate dysplasia showed high ki67 staining meanwhile 13 of 31 severely dysplastic polyps had low ki67 score and remaining 18 had ki67 score.

Table No. 1: Baseline demographics of the colorectal adenomatous cases

Variables	Frequency (n=55)	%
Gender		
Male	36	65.5 %
Female	19	34.5 %
Mean age (years)	53.27±10.84	
Clinical symptom		
Bleeding per rectum	35	63.64%
Chronic diarrhoea and weight loss	8	14.55%
Anemia	6	10.91%
Chronic abdominal pain	3	5.45%
constipation	2	3.64%
Accidental finding	1	1.81%
Site of Colorectal Adenomatous		
rectum	23	41.81%
ascending colon	10	18.2%
Sigmoid colon	10	18.2%
Descending colon	7	12.7%
transverse colon	5	9.09%
Size of Colorectal Adenomatous Polyps		
>20mm	14	25.45%
<20mm	41	74.54%
Dysplasia in colorectal adenomatous polyps		
Mild	21	38.19%
moderate	3	5.45%
Severe	31	56.36%

Table No. 2: showing Association of size of colorectal Adenomatous polyps with ki67 score

Ki-67 Score				
		High	Low	Total
size	<20mm	19	22	41
	>20mm	13	1	14
Total		32	23	55
Chi-Square Tests				
		Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square		9.281 ^a	1	.002
Continuity correction ^b		7.468	1	.006
Likelihood Ratio		10.944	1	.001
N of Valid Cases		55		

Chi square test was performed and P value of less than 0.05 was obtained showing strong association between size of polyps and ki67 score.

Table No. 3: Shows Association between Grades of dysplasia and ki67 score

		Ki-67 Score		Total
		High	Low	
Grades dysplasia	Mild	7	14	21
	Moderate	2	1	3

	Serve	23	8	31
Total		32	23	55
Chi-Square Tests				
Pearson Chi-Square		8.685 ^a	2	.013
Likelihood Ratio		8.811	2	.012
N of Valid Cases		55		

Chi-square test was performed and P-value of < 0.05 was obtained showing stronger association.




Figure Nos. 1 & 2: showing high Ki67 staining in the glands.

DISCUSSION

A number of tumors in humans have been shown to express Ki-67, which is linked to malignancies.⁽¹⁶⁾ Although some research⁽¹⁷⁾ failed to show the prognostic significance of higher Ki-67 expression, others reported this marker as an independent predictor of human colorectal cancer.⁽¹⁸⁾

In current study, 55 adenomatous polyps were retrieved. There were 36 (65.5%) males and 19 (34.5%) females in this study. Patients mean age was 53.27±10.84 years. Results were inline with the study done by Nusrat et al⁽¹⁹⁾ Our study reported vast majority of severely dysplastic colorectal Adenomatous polyps, their proportion can be considered quite high when compared to previous research.^(17,18)

High expression of Ki67 is also strongly associated with the incidence and progression of colorectal cancer⁽²⁰⁾ The severity of dysplasia and Ki-67

expression did not significantly correlate, according to research by Vernillo et al⁽²¹⁾. Another study showed significant correlation between size of colorectal Adenomatous polyps and grades of dysplasia with ki67 score, these findings were aligned with the findings of another study.

Ki67 score showed also significant correlation with size, type and high grade dysplasia in a previous study and the role of this marker was highlighted as an ancillary marker for the risk of transformation and as a target for chemo-preventive drugs⁽²²⁾, while our study worked on size and dysplasia correlation with ki67 score and the type of adenoma part should be considered for further research. High levels of ki67 positive cells in colonic samples indicated poor prognosis and comparatively adverse stage of colorectal carcinoma⁽²³⁾. This indicates the significance of ki67 screening in aiding in early diagnosis and prompt treatment of malignantly potential polyps.

Another study revealed significant association between dysplasia and ki67 score while the association remained insignificant in other studies⁽²⁴⁾.

Overall, the importance of ki67 scoring in dysplastic colorectal Adenomatous polyps with large size is significantly evident in screening and early diagnosis of colorectal Adenomatous polyps with potential of malignant transformation.

CONCLUSION

When identifying the individuals with colorectal adenoma who require close monitoring in follow-up prevention activities, high grade dysplasia and significant positive immunohistochemistry markers of Ki-67 may be useful criteria. Particularly in patients with a large size adenoma and high grade dysplasia.

Author's Contribution:

Concept & Design of Study: Sabika Batool
 Drafting: Talat Mirza, Fouzia Lateef
 Data Analysis: Sobia Hassan
 Revisiting Critically: Sabika Batool, Talat Mirza
 Final Approval of version: Sabika Batool

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: ERC Reference No. 5930922 SBPAT dated 24th October 2022

REFERENCES

1. Kumar V, Abbas AK, Fausto N. Robbins and Cotran. Pathologic Basis of Disease, 8th ed. Elsevier, Philadelphia, USA; 2010.p.819.
2. Winawer SJ, Zauber AG, Fletcher RH, Stillman JS, O'Brien MJ, Levin B, et al. US Multi-Society Task Force on Colorectal Cancer. American Cancer Society Guidelines for colonoscopy surveillance after polypectomy: a consensus update by the US Multi-Society Task Force on Colorectal Cancer and the American Cancer Society. *Gastroenterol* 2006;130(6):1872-1885.
3. Andrey B, Alexander D. "Peculiarities of Vascular Component of Communicative Systems in Rectal Adenomas". *Int J Collaborative Research Internal Med Public Health* 2009;(1):12-21.
4. Atkin WS, Saunders BP, British Society for Gastroenterology. "Association of Coloproctology for Great Britain and Ireland Surveillance guidelines after removal of colorectal adenomatous polyps". *Gut* 2002; 51(5)(Suppl 5):V6-V9 10.1136/gut.51.suppl.5.v6.
5. Hahn WC, Weinberg RA. "Modelling the molecular circuitry of cancer". *Nat Rev Cancer* 2002;2(5):331-341..
6. Ghiță C, Vilcea ID, Dumitrescu M, Vilcea AM, Mirea CS, Așchie M, et al. The prognostic value of the immunohistochemical aspects of tumor suppressor genes p53, bcl-2, PTEN and nuclear proliferative antigen Ki-67 in resected colorectal carcinoma. *Romanian J Morphol Embryol* 2012;53:549-56.
7. Allegra CJ, Paik S, Colangelo LH, Parr AL, Kirsch I, Kim G, et al. Prognostic value of thymidylate synthase, Ki-67, and p53 in patients with dukes' B and C colon cancer: a National Cancer Institute-National Surgical Adjuvant Breast and bowel project collaborative study. *J Clin Oncol* 2003;21:241-50.
8. Melling N, Kowitz CM, Simon R, Bokemeyer C, Terracciano L, Sauter G, et al. High Ki67 expression is an independent good prognostic marker in colorectal cancer. *J Clin Pathol* 2016;69:209-14.
9. Walysson Alves Tocantins de Sousa, LusmarVeras Rodrigues, Raimundo Gerônimo da Silva Jr, Fernando Lopes, "Immunohistochemical evaluation of p53 and Ki-67 proteins in colorectal adenomas". *Vieira Department of Surgery, Federal University of Ceará, Fortaleza, Brazil Arq. Gastroenterol* 2012;49(1):
10. Winawer SJ, Zauber AG, Fletcher RH, Stillman JS, O'Brien MJ, Levin B, et al; US MultiSociety Task Force on Colorectal Cancer; American Cancer Society. "Guidelines for colonoscopy surveillance after polypectomy: a consensus update by the US Multi-Society Task Force on Colorectal Cancer and the American Cancer Society". *Gastroenterol* 2006;130(6):1872-1885.
11. Pathmanathan N, Balleine RL. Ki67 and proliferation in breast cancer. *J Clin Pathol* 2013.
12. Sobocki M, Mrouj K, Camasses A, Parisi N, Nicolas E, Llères D, et al. The cell proliferation

- antigen Ki-67 organises hetero-chromatin. *Elife* 2016;5:e13722.
13. Slaoui M, Fiette L. Histopathology procedures: from tissue sampling to histopathological evaluation. *Drug Safety Evaluation: Methods and Protocols* 2011;69-82.
 14. De Matos LL, Truffelli DC, De Matos MG, da Silva Pinhal MA. Immunohistochemistry as an important tool in biomarkers detection and clinical practice. *Biomarker Insights* 2010;5:BMI-S2185.28.
 15. Soliman NA, Yussif SM. Ki-67 as a prognostic marker according to breast cancer molecular subtype. *Cancer Biol Med* 2016;13(4):496.
 16. Melling N, Kowitz CM, Simon R, Bokemeyer C, Terracciano L, Sauter G, et al. High Ki67 expression is an independent good prognostic marker in colorectal cancer. *J Clin Pathol* 2016;69:209–14.
 17. Ghiță C, Vilcea ID, Dumitrescu M, Vilcea AM, Mirea CS, Așchie M, et al. The prognostic value of the immunohistochemical aspects of tumor suppressor genes p53, bcl-2, PTEN and nuclear proliferative antigen Ki-67 in resected colorectal carcinoma. *Romanian J Morphol Embryol* 2012;53:549–56.
 18. Pietrzyk L, Torres A, Maciejewsk R, Torres K. Obesity and obese-related chronic low-grade inflammation in promotion of colorectal cancer development. *Asian Pac J Cancer Prev* 2015;16:4161–8.
 19. Nussrat FL, Ali HH, Hussein HG, Al-Ukashi RJ. Immunohistochemical Expression of ki-67 and p53 in Colorectal Adenomas: A Clinicopathological Study. *Oman Med J* 2011;26(4):229-34.
 20. Kasprzak, A. Prognostic Biomarkers of Cell Proliferation in Colorectal Cancer (CRC): From Immunohistochemistry to Molecular Biology Techniques. *Cancers* 2023;15:4570.
 21. Vernillo R, Lorenzi B, Banducci T, Minacci C, Vindigni C, LucentiFei A, et al. Immunohistochemical expression of p53 and Ki67 in colorectal adenomas and prediction of malignancy and development of new polyps. *Int J Biol Markers* 2008;23(2):89-95.
 22. Maarroof TM, et al. P53 Gene and Ki 67 Proliferative Index in Colorectal Adenomas-A Clinico-pathological Study. *Pak J Med Health Sciences* 2020;14(2):1162-1167.
 23. Zhabagin KT. Expression of Ki-67 as a prognostic factor in patients with colorectal cancer. *Annals Oncol* 2019;30:ix39.
 24. Scholzen T, Gerdes J. The Ki-67 protein: from the known and the unknown. *J Cellular Physiol* 2000;182(3):311-22.

Analysis of Fingerprint Patterns in Relation to ABO Blood Groups: A Comparative Study

Fingerprint
Patterns in
Relation to ABO
Blood Groups

Zulfiqar Ali Buzdar¹, Mansoor Mirza², Ambreen Serwer³, Muhammad Anwar Sibtain Fazli³ and Faiza Munir⁴

ABSTRACT

Objective: To assess and analyze the potential association between blood groups and fingerprint patterns.

Study Design: Descriptive Cross-sectional Study

Place and Duration of Study: This study was conducted at the Forensic Medicine & Toxicology Department of Sahara Medical College Narowal, Punjab-Pakistan from July 2020 to June 2023.

Methods: A sample of 293 participants was meticulously studied, categorizing individuals based on their blood groups and assessing their respective fingerprint patterns. This research delves into the intriguing relationship between blood groups and fingerprint patterns, shedding light on potential associations that could have far-reaching implications.

Results: The findings are presented in a comprehensive cross tabulation analysis, revealing a statistically significant association between blood groups and fingerprint patterns ($p = 0.013$). Particularly, the blood group of an individual appears to influence the distribution of their fingerprint pattern.

Conclusion: These results provide a promising foundation for further exploration of the mechanisms underlying this association and its potential applications in fields such as forensics and medical diagnostics. This study marks a crucial step towards understanding the intricate interplay between genetic factors and biometric features, opening new avenues for research and practical applications.

Key Words: ABO, Blood Group, Rh factor, Fingerprint Patterns, Biometrics, Forensic Investigations, Genetic Determinants.

Citation of article: Buzdar ZA, Mirza M, Serwer A, Fazli MAS, Munir F. Analysis of Fingerprint Patterns in Relation to ABO Blood Groups: A Comparative Study. Med Forum 2024;35(1):33-36. doi:10.60110/medforum.350107.

INTRODUCTION

Fingerprint patterns and blood groups are distinctive and integral biological characteristics that have captivated the attention of researchers across diverse fields for decades¹. The uniqueness of fingerprints in forensic science and biometric authentication has long been recognized, making them invaluable tools for individual identification and criminal investigations².

¹. Department of Forensic Medicine & Toxicology, Sahara Medical College, Narowal.

². Department of Forensic Medicine & Toxicology, KEMU, Lahore.

³. Department of Forensic Medicine & Toxicology, Avicenna Medical & Dental College, Lahore.

⁴. Department of Forensic Medicine & Toxicology, Allama Iqbal Medical College, Lahore.

Correspondence: Dr. Zulfiqar Ali Buzdar, Associate Professor of Forensic Medicine & Toxicology, Sahara Medical College, Narowal.

Contact No: 03336011247

Email: forensicatitsbest@gmail.com

Received: August, 2023

Accepted: December, 2023

Printed: January, 2024

Simultaneously, the classification of human blood into different ABO and Rh groups is of paramount significance in medical, medicolegal and clinical practices with implications for transfusions, organ transplantation, and disease susceptibility³. Intriguingly, the potential relationship between these two biological markers has largely remained unexplored. While both fingerprint patterns and blood groups are genetically determined, the concept of an association between the two holds the promise of uncovering novel insights into the hereditary basis of these traits⁴. Such an association could have implications not only in the fields of forensics and biometrics but also in understanding the broader genetic underpinnings of human variation⁵. This research endeavors to bridge this intriguing gap by conducting a systematic examination of the potential correlation between fingerprint patterns and ABO blood group types, considering the presence or absence of the Rh antigen⁶. Our study delves into a dataset comprising 293 individuals, meticulously categorizing their blood groups and analyzing their corresponding fingerprint patterns. The results, presented in a cross tabulation table, reveal a statistically significant association ($p = 0.013$) between blood groups and fingerprint patterns. Specifically, the blood group of an individual appears to influence the distribution of their fingerprint pattern⁷.

These findings not only provide a novel perspective on the potential interplay between genetics and biometric features but also have practical implications in the realms of forensic science, biometric security, and medical diagnostics⁸. A deeper understanding of this association could lead to more accurate and personalized biometric systems, as well as improved forensic and medical practices⁴⁻⁸.

This study marks a crucial step toward unraveling the complex genetic determinants of these unique biological traits and underscores the significance of multidisciplinary research in uncovering novel associations and applications within the scientific community.

METHODS

Data for this research study were collected from a targeted population consisting of students enrolled at Sahara Medical College, falling within the age range of 19 to 23 years. The study specifically focused on students during their third year of academic sessions, spanning from July 2020 to June 2023. Prior to the commencement of data collection, the research protocol adhered to a stringent ethical framework. Informed consent was obtained from each participant, accompanied by a detailed explanation of the study's nature and objectives, the voluntary nature of participation, and the assurance of data confidentiality. Furthermore, this study received approval from the Institutional Review Board (IRB) of Sahara Medical College, ensuring ethical adherence and compliance with established research protocols.

The research involved the administration of a structured questionnaire designed to gather information related to both ABO blood group configurations (including the Rh antigen) and the pattern of fingerprint. The cross-sectional study was carried out using convenient consecutive sampling technique specifically targeted students in the stipulated age group who were willing to participate.

RESULTS

Demographics:

The study included 293 research participants. There were 5 students of 19 years of age, 73 students of age of 20 years, 152 were 21 years old while 60 students were 22 years of age and lastly there were only three students in the 23rd years of their age. The frequencies are shown in the table 01 below.

Table No. 1: Age

Serial No.	Age (in Years)	Frequency	Valid Percentage
1	19	5	1.6
2	20	73	24.9
3	21	152	51.9
4	22	60	20.5
5	23	3	1.0
Total		293	100.0

Out of 293 students there were 106 (36.2%) males just and remaining 187 (63.8%) were females. The composition depicts almost 1:3 ratio from male to female respectively. The frequencies are shown in table 02 below.

Table No. 2: Gender

Serial No.	Gender	Frequency	Valid Percentage
1	Male	106	36.2
2	Female	187	63.8
Total		293	100.0

Analysis of Fingerprint and Blood Group Association:

Fingerprint Pattern: The predominant fingerprint pattern observed in the sample was the "Loop," with 101 instances (34.5% of the total patterns). Within the "Loop" category, the "Radial" sub-variation was the most frequent, accounting for 45.7% of all patterns. In contrast, the "Ulnar" sub-variation within the "Loop" pattern was observed 33 times, making up 11.3% of the total patterns. The "Whorl" pattern was the second most common, representing 36.9% of the total patterns. Among "Whorl" patterns, the "Plain" sub-variation was the most prevalent, constituting 40.6% of the "Whorl" patterns. The "Double Loop" and "Composite" patterns were less frequent, with 3.8% and 0.7% of the total patterns, respectively.

Blood Groups Analysis: The most common blood group in the population was "A positive," representing 20.8% of the total blood groups. In contrast, "A negative" blood group was relatively less common at 0.7%. Notably, "B positive" blood group was the most prevalent in the study population, accounting for 35.2% of the total blood groups. "B negative" blood group was observed in 2.4% of the cases. The "AB positive" blood group was identified in 13.0% of the individuals, while "AB negative" comprised 1.4% of the sample. The "O positive" blood group was observed in 23.9% of the cases, with "O negative" accounting for 2.7% of the total blood groups.

Table No. 3: Analysis of Blood Group and Pattern of Fingerprint

Blood Group Configuration		Fingerprint Pattern							Total	P Value
		Loop		Arch		Whirl		Composite		
Blood Group	Rh Factor	Radial	Ulnar	Plain	Tentated	Plain	Double Loop			
A	Positive	27	3	5	2	22	2	0	61	0.013

	Negative	0	0	0	0	2	0	0	2
B	Positive	32	10	14	1	42	3	1	103
	Negative	3	2	1	0	1	0	0	7
AB	Positive	12	8	3	0	13	1	1	38
	Negative	2	1	0	0	0	1	0	4
O	Positive	22	9	9	1	28	1	0	70
	Negative	3	0	1	1	0	3	0	8
Total		101	33	33	5	108	11	2	293

Statistical Significance: A chi-square test of independence was conducted to investigate the potential relationship between "Blood Group" and "Pattern of Fingerprint." The results of this analysis indicate a statistically significant association ($p = 0.013$) between these two variables.

Specifically, "Blood Group" appears to influence the distribution of "Pattern of Fingerprint" within the study population. This finding suggests that the prevalence of specific fingerprint patterns may be associated with an individual's blood group. These results serve as a significant foundation for further research into the mechanisms underlying this association and the potential implications within the broader context of our study on the correlation between blood groups and fingerprint patterns.

DISCUSSION

The pursuit of understanding the intricate and nuanced facets of human biology has long captivated the scientific community⁹. In this study, we endeavored to unravel a potential relationship between two distinct biological traits: blood groups and fingerprint patterns¹⁰. This endeavor is underpinned by the premise that both characteristics are genetically determined and could potentially exhibit an association^{6,10,11,12}. The statistical analysis of our data revealed a significant association between "Blood Group" and "Pattern of Fingerprint." Specifically, the chi-square test of independence yielded a p-value of 0.013, signifying that the distribution of fingerprint patterns is not independent of blood group. This finding sheds light on a previously under-explored dimension of biometric research and raises a multitude of questions and considerations.

The observed association prompts the need for a deeper understanding of its significance and potential implications^{13,14}. One possible explanation for this association could be the genetic determinants that influence both blood group and fingerprint pattern development¹⁵. The influence of genetics on fingerprint patterns is well-established, and it is plausible that genes governing blood group also play a role in the formation of fingerprint patterns¹⁶. The implications of this association are multifaceted. In the realm of forensic medicine, where fingerprint patterns hold great significance, the knowledge that blood groups may influence these patterns opens new avenues for

investigation¹⁷. Fingerprint analysis in forensic investigations could benefit from considering an individual's blood group in conjunction with pattern analysis, potentially aiding in the identification and profiling of individuals^{6,18}. Moreover, in biometric security, the findings of this study have the potential to enhance the accuracy and reliability of fingerprint-based authentication systems, contributing to both security and convenience¹⁹.

CONCLUSION

In conclusion, the association between blood groups and fingerprint patterns represents a fascinating intersection of genetic factors and biometric features. Our study has illuminated a statistically significant link between these two variables, creating a foundation for future research endeavors. The potential applications of this association in forensic science, biometric security, medical diagnostics, and genetic studies are promising. This discovery underscores the ever-evolving landscape of biometric research and the intriguing interplay between genetics and human variation.

Author's Contribution:

Concept & Design of Study: Zulfiqar Ali Buzdar
 Drafting: Mansoor Mirza
 Data Analysis: Ambreen Serwer
 Revisiting Critically: Muhammad Anwar Sibtain Fazli
 Final Approval of version: Faiza Munir

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.21/FMT/SMC dated 04.06.2020

REFERENCES

- Patil V, Ingle DR. An association between fingerprint patterns with blood group and lifestyle based diseases: a review. *Artificial Intelligence Review* 2021;54:1803-39.
- Jain AK, Nandakumar K, Ross A. 50 years of biometric research: Accomplishments, challenges, and opportunities. *Pattern recognition letters* 2016;79:80-105.
- Mohammed AK. Blood Groups ABO and Rh System Among Paternity and Kinship Cases of

- Iraqi Medical Legal Directorate. *J Educ Psychological Res* 2022;4(3):555-9.
4. Shrestha DB, Gupta VP, Chaurasiya PS, Shrestha S, Chaudhary S, Aryal L. Study of correlation between different fingerprint patterns, blood groups, and social behavior among medical students (Nepalese Citizens). *Pac J Sci Technol* 2016;17(2):288-92.
 5. Li J, Glover JD, Zhang H, Peng M, Tan J, Mallick CB, et al. Limb development genes underlie variation in human fingerprint patterns. *Cell* 2022;185(1):95-112.
 6. Koura SM, Abdel-Rahman RH, Emam NM. Role of Fingerprints Patterns and ABO/Rh Blood Groups in Sex Dimorphism among Egyptian Population. *Mansoura J Forensic Med Clin Toxicol* 2022;30(2):1-17.
 7. Suganthi V, Periasamy P, Gunasekaran S. Study of Fingerprint Patterns and Their Relationship With Blood Groups and Gender In Erode, Tamilnadu. *J Pharmaceutical Negative Results* 2023;14(2).
 8. Mohsin AH, Zaidan AA, Zaidan BB, Albahri AS, Albahri OS, Alsalem MA, et al. Real-time remote health monitoring systems using body sensor information and finger vein biometric verification: A multi-layer systematic review. *J Med Systems* 2018;42:1-36.
 9. Ford AT, Ali AH, Colla SR, Cooke SJ, Lamb CT, Pittman J, et al. Understanding and avoiding misplaced efforts in conservation. *Facets* 2021;6(1):252-71.
 10. AL-Kalifa NF. A study of morphological patterns of dermatoglyphics among mother and fetus in different ABO, RH blood groups. *Diyala J Med* 2017;13(2):80-6.
 11. Bhavana D, Ruchi J, Prakash T, JL K. Study of fingerprint patterns in relationship with blood group and gender-a statistical review. *Arches* 2013;1(1):15-7.
 12. Kamaradgi PN, Marigoudar RM, Jatti VB, Kumar A. A Correlation Study of Patterns of Fingerprints with Blood Groups among the Students of SSIMS & RC, Davangere. *Ind J Forensic Med Toxicol* 2019;13(2).
 13. Shrestha R, Hirachan N, Koju SS, Lamichhane A. Association of Fingerprints with the ABO Blood grouping among students in Gandaki Medical College. *J Gandaki Med College-Nepal* 2019;12(2):63.
 14. Aamir Y, Masood R, Irshad N, Malik R, Farid N, Shahab MA. Relationship between Pattern of Fingerprints and Blood Groups. *Pak J Med Health Sciences* 2022;16(09):698-99.
 15. Gupta Y. Fingerprint patterns and abo blood groups: a study of female nursing students. *Int J Anat Physiol* 2023;4(05):01-5.
 16. Patil VN, Ingle DR. A Novel Approach for ABO Blood Group Prediction using Fingerprint through Optimized Convolutional Neural Network. *Int J Intelligent Systems Applications Engineering* 2022;10(1):60-8.
 17. Debta FM, Debta P, Bhuyan R, Swain SK, Sahu MC, Siddhartha S. Heritability and correlation of lip print, palm print, fingerprint pattern and blood group in twin population. *J Oral Maxillofacial Pathol : JOMFP* 2018;22(3):451.
 18. Patil VN, Ingle DR. A Novel Approach for ABO Blood Group Prediction using Fingerprint through Optimized Convolutional Neural Network. *Int J Intelligent Systems Applications Engineering* 2022;10(1):60-8.
 19. Vadde CS, Kumar PA, Ramakhrisna P. A Study Of Pattern Of Fingerprints In Relation To Blood Groups. *East African Scholars J Med Sciences* 2020;3(2):37-40.

Gender-Based Analysis of the Lambdoid Cranial Suture among Human Cadavers Presented for Postmortem Examination at King Edward Medical University, Lahore

Mansoor Mirza¹, Zulfiqar Ali Buzdar², Ambreen Serwer³, Muhammad Anwar Sibtain
Fazli³ and Faiza Munir⁴

ABSTRACT

Objective: The objective of this study is to investigate the gender-based differences in the commencement of lambdoid suture closure.

Study Design: Descriptive Study

Place and Duration of Study: This study was conducted at the Department of Forensic Medicine and Toxicology, King Edward Medical University, Lahore from January to September during the year 2016.

Methods: A total of 90 deceased individuals, comprising an equal number of males and females within the age span of 20 to 70 years, were subjected to medicolegal autopsy. Standardized autopsy protocols were followed, and the lambdoid suture was meticulously examined. Suture fusion was observed macroscopically both endocranially and ectocranially, with a five-grade scale applied to quantify closure stages.

Results: The results of the analysis revealed distinct gender-based differences in the commencement of lambdoid suture closure having profound statistically significant with a p-value of < 0.05 .

Conclusion: The study elucidates that gender plays a pivotal role in the commencement of lambdoid suture closure. The observed differences can have implications in forensic age estimation, providing valuable information for postmortem examinations

Key Words: Gender Dimorphism, Lambdoid Suture, Suture Closure, Forensic Medicine, Postmortem Examination, Ectocranial, Endocranial.

Citation of article: Mirza M, Buzdar ZA, Serwer A, Fazli MAS, Munir F. Gender-Based Analysis of the Lambdoid Cranial Suture among Human Cadavers Presented for Postmortem Examination at King Edward Medical University, Lahore. *Med Forum* 2024;35(1):37-41. doi:10.60110/medforum.350108.

INTRODUCTION

Gender estimation, a critical facet of forensic anthropology, plays a pivotal role in postmortem examinations and the determination of an individual's identity¹. Within this broader field, the study of gender

dimorphism in the cranial sutures has always been compelling area of investigation as one of the prime objective of autopsy². The cranial sutures, essential for the development and expansion of the human skull, exhibit variations that can be influenced by factors such as age³, ancestry⁴, and notably by gender⁵. While multiple cranial sutures contribute to gender estimation, the lambdoid suture stands out as a prime focus in our investigation. The lambdoid suture, located at the junction of the parietal and occipital bones, exhibits gender-based differences in its closure patterns⁶. A comprehensive understanding of these variations is invaluable in forensic anthropology, as it aids in determining an individual's gender during postmortem examinations⁷.

Gender dimorphism in the lambdoid suture provides valuable insights into the differences in the commencement and progression of suture closure between males and females⁸. This distinction can prove instrumental in enhancing the accuracy of gender estimation, particularly in forensic cases where the identity of the deceased is unknown⁹. This sometime becomes the only option in the face of advanced

¹. Department of Forensic Medicine & Toxicology, KEMU, Lahore.

². Department of Forensic Medicine & Toxicology, Sahara Medical College, Narowal.

³. Department of Forensic Medicine & Toxicology, Avicenna Medical & Dental College, Lahore.

⁴. Department of Forensic Medicine & Toxicology, Allama Iqbal Medical College, Lahore.

Correspondence: Dr. Mansoor Mirza, Assistant Professor of Forensic Medicine & Toxicology, KEMU, Lahore.

Contact No: 03377550756

Email: dr.mansoormirza7@gmail.com

Received: June, 2023

Accepted: October, 2023

Printed: January, 2024

decomposition or when rest of the body is not available or where only skull has been found¹⁰. Gender estimation, therefore, serves as an indispensable tool for forensic practitioners, law enforcement agencies, and medicolegal experts, aiding in the identification of human remains and facilitating criminal investigations¹¹. This study aims to delve into the intricacies of gender dimorphism within the lambdoid suture, shedding light on the variations in suture closure patterns based on gender. By examining both ectocranial and endocranial aspects, we seek to elucidate the specific differences in the onset and progression of closure¹². The insights garnered from this research hold substantial promise in improving the accuracy of gender estimation in forensic scenarios, contributing to the field's ongoing evolution¹³.

METHODS

The study was conducted on a sample of 90 deceased individuals, with an equal distribution of males and females. The study subjects were selected within the age span of 20 to 70 years, ensuring a representative cross-section of the population. All cases were brought to the Department of Forensic Medicine and Toxicology at King Edward Medical University, Lahore, for a period arching over 9 months, spanning from Jan-2016 to Sept-2016. The study adhered to ethical principles and guidelines governing ethical committee of King Edward Medical University and due approval was taken. The study protocols followed the guidelines in accordance with institutional and ethical standards.

Data collection followed a meticulous and standardized autopsy protocol, with each postmortem examination adhering to established forensic guidelines. The scalp covering of the skull was lifted by employing curved mastoid-to-mastoid incision, ensuring that the thorough stretching of the lambdoid suture be visualized to unaided eye. This approach allowed for the complete exposure of the suture. All tissues tags and fats were

meticulously removed to facilitate examination of the suture.

To ensure a comprehensive analysis of the lambdoid suture, it was divided into three equal parts, referred to as L1, L2, and L3. This division aided in systematically examining suture closure within different subsections of the suture.

The degree of fusion in all three parts of the lambdoid suture was recorded separately for both the ectocranial (outer table) and endocranial (inner table) aspects. Closure stages were assessed using the Acsádi-Nemeskéri Scale referring to 0 - as "Open with no closure at all", 1- as "Incipient closure with less than half closed", 2 - as "Closure in process with half of the suture closed", 3 - as "Advanced closure with more than half of the suture closed" and 4 - as "Closed completely"

RESULTS

The study included equal number of the male and female case of postmortem examination of ages more than 20 years and less than 70 years. The subjects were divided in five groups each comprising of a decade. The first group of 21 to 30 years had 10 males and 22 females, the second group of 31 to 40 years of age had 08 males and 07 females whereas the third group of age from 41 to 50 years comprised of 10 males and 08 females. In the second last category of 51 to 60 years were 08 males and 06 females and the last group from 61 to 70 years had 09 males and just 02 females.

For all the tables in results sections 'n' is number and abbreviation like 'Ecto-L1' shows Ectocranial Lambdoid Suture Subsection 1, 'Endo-L1' shows Endocranial Lambdoid Suture Subsection 1, 'Ecto-L2' shows Ectocranial Lambdoid Suture Subsection 2, 'Endo-L2' shows Endocranial Lambdoid Suture Subsection 2, 'Ecto-L3' shows Ectocranial Lambdoid 1 Suture Subsection 3 and 'Endo-L3' shows Endocrania Lambdoid Suture Subsection 3.

Table No. 1: The assessment of closure stages of lambdoid suture in males (n=45)

Age group	n	Ecto -L1	Endo-L1	Ecto-L2	Endo-L2	Ecto-L3	Endo-L3
		Mean ± SD		Mean ± SD		Mean ± SD	
21-30 years	10	0.90±0.316	1.30±0.483	0.10±0.316	1.20±1.033	0.70±0.483	1.00±0.000
31-40 Years	08	1.88±0.641	2.38±0.518	1.25±0.463	1.75±0.707	2.00±0.535	2.38±0.744
41-50 Years	10	3.30±0.675	3.50±0.527	2.30±0.483	3.00±0.000	3.20±0.632	3.60±0.516
51-60 Years	08	3.88±0.354	4.00±0.000	3.38±0.518	3.75±0.164	3.50±0.535	3.25±0.463
61-70 Years	09	4.00±0.000	4.00±0.000	3.56±0.527	4.00±0.000	3.89±0.333	4.00±0.000

In the table 01 above, the age group of 21-30 years, we found no significant closure for lambdoid suture closure helpful for age or gender estimation. In the age group of 31-40 years, we observed advanced closure (Mean ± SD) in Endo - L2 1.75. In the age group of 41-50 years, advanced closure was noted Endo - L1 with a mean

value of 3.50. For individuals aged 51-60 years, the following observations were made: Endo - L1 exhibited complete closure (Mean ± SD: 4.00) Endo - L2 showed advanced closure with a mean value of 3.75 and Endo - L3 demonstrated incipient closure with a mean value of 3.25. In the age group of 61-70 years,

complete closure (Mean ± SD: 4.00) was observed in ectocranially in L1 and 3edocranially in Lambdoid

Suture Subsections 1 (Endo - L1), 2 (Endo - L2), and 3 (Endo - L3).

Table No. 2: The assessment of closure stages of lambdoid suture in females (n=45)

Age group	n	Ecto -L1	Endo-L1	Ecto-L2	Endo-L2	Ecto-L3	Endo-L3
		Mean ± SD		Mean ± SD		Mean ± SD	
21 - 30 years	22	0.95±0.486	1.59±0.734	0.41±0.503	0.77±0.685	0.86±0.744	1.41±0.734
31-40 Years	7	1.86±0.690	1.86±0.378	1.00±0.000	1.57±0.535	1.71±0.488	1.71±0.488
41-50 Years	8	3.00±0.535	3.63±0.518	2.50±0.926	2.75±0.463	2.75±0.463	3.13±0.835
51-60 Years	6	3.17±0.408	3.67±0.516	3.00±0.000	3.00±0.000	3.00±0.000	3.67±0.516
61-70 Years	2	4.00±0.000	4.00±0.000	3.50±0.707	4.00±0.000	3.50±0.707	4.00±0.000

The Table 02 presents the assessment of closure stages of the lambdoid suture in females. In the age group of 21-30 years, we found no significant lambdoid suture closure in females. In the age group of 31-40 years, advanced closure was noted in both Ectocranial and Endocranial Lambdoid Suture Subsection 1 (Ecto - L1 and Endo - L1), with mean values of 1.86. For individuals aged 41-50 years, advanced closure was observed in Ectocranial Lambdoid Suture Subsection 1 (Ecto - L1) with a mean value of 3.00. In the age group of 51-60 years, advanced closure was found in the subsections i.e. Ectocranial Lambdoid Suture

Subsection 2 (Ecto - L2) with a mean value of 3.00, Endocranial Lambdoid Suture Subsection 2 (Endo - L2) with a mean value of 3.67 and Ectocranial Lambdoid Suture Subsection 3 (Ecto - L3) with a mean value of 3.00

In the age group of 61-70 years, complete closure was observed in all subsections whether endocranial or ectocranial of L1, L2 and L3. Notably, incipient closure was observed in Ectocranial Lambdoid Suture Subsection 2 (Ecto - L2) in the age group of 31 – 40 years.

Table No. 3: The assessment of closure stages of lambdoid suture in all the subjects (n=90)

Age group	n	Ecto -L1	Endo-L1	Ecto-L2	Endo-L2	Ecto-L3	Endo-L3
		Mean ± SD		Mean ± SD		Mean ± SD	
21 - 30 years	32	0.94±0.435	1.50±0.672	0.31±0.471	0.78±0.608	0.81±0.693	1.28±0.634
31-40 Years	15	1.87±0.640	2.13±0.516	1.13±0.352	1.67±0.617	1.87±0.516	2.07±0.704
41-50 Years	18	3.17±0.618	3.56±0.511	2.39±0.698	2.89±0.323	3.00±0.594	3.39±0.698
51-60 Years	14	3.57±0.514	3.86±0.511	3.21±0.462	3.43±0.514	3.29±0.169	3.43±0.514
61-70 Years	11	4.00±0.000	4.00±0.000	3.55±0.522	4.00±0.000	3.82±0.405	4.00±0.000

Table 3 presents the results of the assessment of closure stages of the lambdoid suture in a combined sample of all subjects, comprising a total of 90 individuals. This table provides valuable insights into the closure patterns of the lambdoid suture across different age groups. In the age group of 21-30 years, we observed no significance for lambdoid suture closure in estimating sexual dimorphism. In the age group of 31-40 years, advanced closure was noted in both Ectocranial and Endocranial Lambdoid Suture Subsection 1 (Ecto - L1 and Endo - L1), with mean values of 1.87. For individuals aged 41-50 years, advanced closure was observed in Ectocranial Lambdoid Suture Subsection 1 (Ecto - L1) with a mean value of 3.17. In the age group of 51-60 years, advanced closure was found in the following subsections like Ectocranial Lambdoid Suture Subsection 1 (Ecto - L1) with a mean value of 3.57, Endocranial Lambdoid Suture Subsection 1 (Endo - L1) with a mean value of 3.86 and Ectocranial Lambdoid Suture Subsection 3 (Ecto - L3) demonstrated incipient closure with a mean value of 3.29. In the age group of 61-70 years, complete closure was observed in all the subsections of lambdoid suture whether endocranially or ectocranially.

Gender-Based Comparison for Commencement of Lambdoid Suture Closure

Table 04 presents the gender-based comparison of the commencement of suture closure within the lambdoid suture. This analysis was carried out for both the ectocranial and endocranial aspects

Table No. 4: Gender based commencement of lambdoid suture closure

Ectocranial Closure				
Gender	n	Mean Score	P value	Significance
Male	45	2.48	< 0.05	Significant
Female	45	1.66		
Endocranial Closure				
Gender	n	Mean Score	P value	Significance
Male	45	2.81	< 0.05	Significant
Female	45	2.08		

Table 04 illustrates the gender-based comparison of the commencement of suture closure within the lambdoid suture. Notably, this analysis revealed significant differences in the commencement of lambdoid suture closure based on gender. As indicated in the table, the

commencement of suture closure was observed to be significantly earlier in males than in females in both ectocranial and endocranial aspects. The endocranial aspect, in particular, exhibited a marked difference, with males demonstrating a mean score of 2.81 compared to 2.08 in females. The statistical analysis supported these observations, with a p-value of < 0.05 , underscoring the significance of gender-based differences in the commencement of lambdoid suture closure, as highlighted in Table 04.

DISCUSSION

Gender estimation is a fundamental aspect of forensic anthropology, facilitating the identification and categorization of deceased individuals during postmortem examinations¹⁴. The cranial sutures have emerged as key anatomical features for gender estimation due to their potential for exhibiting gender-specific variations in closure patterns⁸. In this study, we notably observed significant differences in closure patterns between males and females¹⁵. These findings have implications for accurate determination of gender is often a critical component in the identification of unknown human remains.

In examining the ectocranial and endocranial aspects of the lambdoid suture, we sought to capture a comprehensive picture of gender dimorphism¹⁶. Our results demonstrated that in both ectocranial and endocranial perspectives, males exhibited an earlier initiation of suture closure compared to females^{6,8,16,17}. This gender-based distinction was particularly pronounced in the endocranial aspect, where males displayed a marked advancement in suture closure¹⁸. Such detailed observations provide forensic experts with a valuable tool for gender estimation^{8,17}. The significance of our findings lies in their potential application within the field of forensic medicine and postmortem examinations. Accurate gender estimation plays a pivotal role in narrowing down the identity of unknown individuals, aiding law enforcement agencies and medicolegal experts in criminal investigations and victim identification¹⁹. The insights gleaned from this study are expected to enhance the precision and reliability of gender estimation methodologies used in forensic anthropology²⁰.

CONCLUSION

Our study, aimed at understanding the gender-based differences in the commencement and progression of lambdoid suture closure, has yielded valuable insights that can contribute to the refinement of gender estimation techniques in the field especially in the circumstance when only skull is available for gender estimation^{16,17,18}. The results of our investigation demonstrate a compelling pattern of gender dimorphism within the lambdoid suture. Notably, males exhibit an earlier initiation of suture closure compared to females,

a distinction that becomes particularly pronounced in the endocranial aspect. This observed gender-based variation in suture closure holds profound implications for forensic anthropology, where the precise determination of an individual's gender is often the linchpin in identifying unknown remains and advancing criminal investigations.

Author's Contribution:

Concept & Design of Study: Mansoor Mirza
Drafting: Zulfiqar Ali Buzdar,

Ambreen Serwer
Data Analysis: Muhammad Anwar
Sibtain Fazli, Faiza
Munir

Revisiting Critically: Mansoor Mirza, Zulfiqar
Ali Buzdar

Final Approval of version: Mansoor Mirza

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.25/RC/KEMU dated 11.01.2016

REFERENCES

1. Tarani S, Kamakshi SS, Naik V, Sodhi A. Forensic radiology: An emerging science. *J Advanced Clin Research Insights* 2017;4(2):59-63.
2. Hadadi AI, Balwi MA. Delineation of craniosynostosis in Saudi Arabia. *J Clin Images Med Case Rep* 2023;4(8):2536.
3. Blessing M, Gallagher ER. Epidemiology, Genetics, and Pathophysiology of Craniosynostosis. *Oral Maxillofacial Surg Clinics* 2022;34(3):341-52.
4. Buzdar ZA, Khan MM. An Autopsy based unaided Eye Study of Lambdoid Suture of Skull—the Science of Forensic Estimation of Age. *Pak J Med Health Sciences* 2022;16(12):206-8.
5. Junn A, Dinis J, Lu X, Forte AJ, Mozaffari MA, Phillips S, et al. Facial Dysmorphism in Saethre-Chotzen Syndrome. *J Craniofacial Surg* 2021;32(8):2660-5.
6. Manrique M, Toro-Tobon S, Bade Y, Paredes-Gutierrez J, Mantilla-Rivas E, Rana MS, et al. Sick Cell Disease Association with Premature Suture Fusion in Young Children. *Plastic and Reconstructive Surg Global Open* 2022;10(10).
7. Raj Kumar MG, Kiran J. A Cross Sectional Postmortem Study on Closure of Skull Vault Sutures with Respect to Age from 3 Rd to 6 Th Decades of Life. *Ind J Forensic Med Toxicol* 2018;12(2).
8. Patil K, VG M, Chandran P, Tayal S, HP J. Applications of Reverse Panoramic Radiography for Age and Gender Determination-A Radiographic Study. *Ind J Forensic Med Toxicol* 2021;15(3).

9. Alhadi A, Issrani R, Prabhu N, Alhadi M. Observation on the closure of lambdoid suture in relation to age, sex and population variations using a novel radiographic technique—a prospective study. *Acta Odontologica Scandinavica* 2019; 77(1):61-5.
10. Cox M. Ageing adults from the skeleton. *Human Osteol Archaeol Forensic Science* 2000;1000: 61-82.
11. Alhazmi A, Vargas E, Palomo JM, Hans M, Latimer B, Simpson S. Timing and rate of sphenoccipital synchondrosis closure and its relationship to puberty. *PLoS One* 2017;12(8):e0183305.
12. Alghamdi M, Alhumsi TR, Altweijri I, Alkhamis WH, Barasain O, Cardona-Londoño KJ, Ramakrishnan R, Guzmán-Vega FJ, Arold ST, Ali G, Adly N. Clinical and Genetic Characterization of Craniosynostosis in Saudi Arabia. *Frontiers Pediatr* 2021;9:582816.
13. Pavičević D, Milošević J, Petrović-Marković I, Milenković Z, Parezanović-Ilić K. The importance of physical treatment in children underwent craniosynostosis surgery in the first year of life. *Vojnosanitetski Pregled* 2020;77(3):324-9.
14. Kumari K, Saleh I, Taslim S, Ahmad S, Hussain I, Munir Z, et al, Rehman UU. Unraveling the Complexity of Apert Syndrome: Genetics, Clinical Insights, and Future Frontiers. *Cureus* 2023;15(10).
15. Buzdar ZA, Munir J, Haq Z, Fazli MA, Qazi FM, Uz F. Gender Disparities in the Coronal Suture Closure of the Cranium—A tool for estimation of sex 2022 (1):143-45
16. Bellaire CP, Devarajan A, Napoli JG, Rutland JW, Liu H, Jacob L, et al. Craniofacial dysmorphology in infants with non-syndromic unilateral coronal craniosynostosis. *J Craniofacial Surg* 2022;33(6): 1903-8.
17. Zanzrukiya KM, Kumar L, Bhalodiya AA. A Cross Sectional Descriptive Study of Analysis of Lambdoid And Squamous Sutures Closure by Ct Scan for Age Estimation. *Ind J Forensic Med Toxicol* 2021;15(1).
18. Abdali H, Anaraki AG, Mahdipour S. Craniosynostosis in Isfahan, Iran: A Cross-Sectional Study. *J Maxillofacial Oral Surg* 2022: 1-7.
19. Ramanan G, Ranganathan S, Ranganathan S. Determination of age by study of closure of endocranial sutures. *J Evolution Med Dent Sciences* 2016;5(46):2860-5.
20. Tarani S, Kamakshi SS, Naik V, Sodhi A. Forensic radiology: An emerging science. *J Advanced Clinical Research Insights* 2017;4(2):59-63.
21. Sjögren KG, Olalde I, Carver S, Allentoft ME, Knowles T, Kroonen G, et al. Kinship and social organization in Copper Age Europe. A cross-disciplinary analysis of archaeology, DNA, isotopes, and anthropology from two Bell Beaker cemeteries. *PLoS One* 2020;15(11):e0241278.

Assessment of the Labial Alveolar Bone Thickness Overlying Maxillary Anterior Teeth in Different Age Groups, Genders, and Sides of the Arch: A Cone Beam Computed Tomographic Study

Naheed Imran¹, Asma Sattar¹, Imran Khattak¹, Sana Arbab¹, Munawar Aziz¹ and Syed Amjad Shah²

ABSTRACT

Objective: To evaluate the thickness of labial alveolar bone at the maxillary anterior teeth region in various age groups and to document the effect of gender and the side of the arch using images obtained by cone-beam computed tomography (CBCT).

Study Design: Cross-sectional retrospective study.

Place and Duration of Study: This study was conducted at the Radiology Department of Khyber College of Dentistry Peshawar, Pakistan (KCD), from 4th November 2021 to 3rd May 2022.

Methods: After the Institutional Review Board (IRB) approval, 350 CBCT images fulfilled the inclusion criteria and were included in the study. The thickness of the labial alveolar bone was measured perpendicular to the long axis of the tooth in a sagittal plane at bone crest level and 2mm, 4mm, and 6 mm apical to CEJ for each tooth in the maxillary anterior region. $P \leq 0.05$ was considered as statistically significant.

Results: The study included a mean age of 39.0 ± 12.6 years and an age range from 18-60 years. The sample was composed of 37.1% males and 62.9% females. The results revealed a significant increase in labial bone thickness with age, particularly 4 mm apical to the CEJ. Maxillary central incisors exhibited the highest thickness, while lateral incisors had the thinnest labial bone. No significant gender difference was found, but lateral asymmetry was observed.

Conclusion: This study reveals age-related changes and regional variations in labial alveolar bone thickness overlying the maxillary anterior teeth. The results emphasize the importance of considering these factors in dental treatment planning to optimize outcomes. Lateral asymmetry emphasizes the need for individualized evaluation of each side during clinical procedures. These insights can guide dental practitioners in making informed decisions for improved treatment and esthetic results.

Key Words: Labial Alveolar Bone Thickness, Maxillary Anterior Teeth, Cemento-Enamel Junction (CEJ), Facial Bone Crest, Dental Implants, Cone Beam Computed Tomography (CBCT).

Citation of article: Imran N, Sattar A, Khattak I, Arbab S, Aziz M, Shah SA. Assessment of the Labial Alveolar Bone Thickness Overlying Maxillary Anterior Teeth in Different Age Groups, Genders, and Sides of the Arch: A Cone Beam Computed Tomographic Study. *Med Forum* 2024;35(1):42-46. doi:10.60110/medforum.350109.

INTRODUCTION

¹. Department of Oral Biology / Oral & Maxillofacial Surgery², Peshawar Medical & Dental College, Peshawar, KPK.

Correspondence: Dr. Naheed Imran, Oral Biology Department, Peshawar Medical & Dental College, Peshawar, KPK.

Contact No: 03149601906

Email: naheedimran.13293@gmail.com

Received: July, 2023

Accepted: August, 2023

Printed: January, 2024

The important function of the labial bone is to support the stability of the tooth root and periodontium in the anterior maxillary region. However, following tooth extraction, there is a risk of resorption of the labial bone, which can lead to various complications in implant therapy⁽¹⁾. For long-term aesthetic outcomes in the front maxilla, adequate horizontal and vertical bone volume is necessary⁽²⁾.

A study conducted in 2018 by Al-Tarawneh et al. aimed to determine the thickness of the labial alveolar bone for the maxillary front teeth in the Amman population. They measured the thickness at three different levels (coronal third, middle third, and apical third) for the central incisors, lateral incisors, and canines. The results showed varying thickness levels for the labial bone at each level and for each type of tooth. For

example, at the coronal third, the labial bone thickness was around 0.7mm for the central incisors, 0.73 for the lateral incisors, and 0.74mm for the canines. In the middle third, the labial bone thickness was approximately 0.69mm for the central incisors, 0.61mm for the lateral incisors, and 0.53mm for the canines. At the apical third, the labial bone thickness was roughly 0.6mm for the central incisors, 0.49mm for the lateral incisors, and 0.4mm for the canines⁽³⁾.

In 2020, Porto OC et al. studied only the upper canines in the Brazilian population and reported a mean labial alveolar bone thickness of 1.49 ± 0.86 mm using CBCT⁽⁴⁾. A study conducted by Xu et al. in 2020 analyzed the labial bone thickness in the Chinese population and found no significant difference between males and females at three different points along the root: 4mm apical to the CEJ, the middle of the root, and the root apex⁽⁵⁾.

Another study conducted on the population of Cairo, Egypt by Ahmed and El Beshlawy (2019) found a noticeable variation in the height and width of the alveolar ridge between male and female participants with males having greater measurements compared to females. However, no significant difference was found between various age groups⁽⁶⁾.

To the best of our knowledge, there has been no prior research conducted on the thickness of the labial alveolar bone in the maxillary anterior teeth in the population of KPK, Pakistan. The study aims to provide dental practitioners with a better understanding of the significance of labial bone thickness in implant cases, to decrease the likelihood of complications such as perforation, fenestration, and dehiscence following implant placement, which may occur as a result of thin labial alveolar bone.

METHODS

The study proposal underwent review and was accepted by the Institutional Ethical Committee at Riphah International University. The Head of the OPD and Radiology Department at Khyber College of Dentistry (KCD) granted permission for data collection, and the hospital administration approved and facilitated the study. The CBCT images used in the study were referred by other dentists for various investigations, such as dental implant therapy, impacted tooth extraction, or orthodontic therapy. The data collection and examination were performed by one examiner, and the interpretation was done by an oral and maxillofacial surgeon.

The inclusion criteria included the CBCT images of both genders with the presence of maxillary anterior teeth bilaterally and age ranging from 18-60 years. On the other hand, teeth that had undergone prosthetic crowns or restorations, bridge abutments or implants in the anterior maxilla, endodontically treated or decayed teeth or teeth with root resorption and presence

of any skeletal discrepancies or congenital dental problems e.g. cleft lip or palate were excluded from the study.

CBCT images were imported to the computer using Planmeca Romexis software (used in KCD). All images used in the present research were obtained using the following range of scanning parameters. Voxel dimension = 4mm, Voltage = 120 kV, Acquisition time = 9 seconds, Current = 5 - 8 mA, DAP (Dose area product) = 761 - 1218mGy*cm², CTDI (Computed tomography dose index) = 4.0 - 6.4mGy.

The labial bone plate thickness was assessed by measuring it in a sagittal plane in the facio- palatal direction perpendicular to the tooth root's long axis. The measurements of the labial wall thickness were noted for each tooth at different levels, including the bone crest level, 2mm, 4mm, and 6mm apical to CEJ in the facio-palatal direction.

This was a retrospective study that followed the ethical standards set by the responsible committee of the institution and was conducted following the principles of the Helsinki Declaration of 1964, as revised in 2013. The confidentiality and anonymity of participants included in the study were ensured. Standardized research protocols were followed.

The statistical analysis was performed using SPSS version 25. Descriptive statistics, including mean values, standard deviations (SD), percentages, and charts, were used to analyze the data. An independent t-test was applied to determine any statistically significant differences between the same tooth and measurement point on the right and left sides. Another independent t-test was also applied to assess any differences in measurements between males and females. A one-way ANOVA test was used to examine any significant differences in various variables among different age groups. A P-value of ≤ 0.05 was considered statistically significant for all tests.

RESULTS





Figure No.

1: Frequency distribution of males and females.


In this study, a sample of 1000 cone-beam computed tomography (CBCT) images were collected from Khyber College of Dentistry.

Table No. 1. Comparison between the labial bone thicknesses of maxillary anterior teeth at all the examined parameters.

Tooth	Levels	Right	Left	Overall Mean ± SD	P-value
Maxillary Central Incisor	At Bone crest in F-P direction	0.73±0.18	0.72±0.18	0.73±0.18	0.986
	At 2mm apical to CEJ	0.39±0.42	0.45±0.42	0.42±0.42	0.002
	At 4mm apical to CEJ	0.73±0.23	0.71±0.23	0.72±0.23	0.438
	At 6mm apical to CEJ	0.74±0.16	0.67±0.16	0.71±0.16	0.068
Maxillary Lateral Incisor	At Bone crest in F-P direction	0.65±0.17	0.65±0.18	0.65±0.18	0.94
	At 2mm apical to CEJ	0.29±0.36	0.26±0.36	0.28±0.36	0.628
	At 4mm apical to CEJ	0.67±0.20	0.60±0.28	0.64±0.24	0.00
	At 6mm apical to CEJ	0.63±0.16	0.60±0.24	0.62±0.21	0.00
Maxillary Canine	At Bone crest in F-P direction	0.73±0.15	0.72±0.17	0.73±0.16	0.000
	At 2mm apical to CEJ	0.25±0.38	0.28±0.39	0.27±0.39	0.73
	At 4mm apical to CEJ	0.70±0.22	0.71±0.22	0.71±0.22	0.795
	At 6mm apical to CEJ	0.66±0.26	0.69±0.15	0.68±0.21	0.000



CI: Central Incisor; LI: Lateral Incisor; C: Canine.
Figures No. 2: Show a comparison between Genders at Bone crest level; at 2mm, 4mm, and 6mm apical to Cementoenamel junction.



CI: Central Incisor; LI: Lateral Incisor; C: Canine.
Figure No. 3: Frequency distribution according to labial bone thickness at bone crest level and 2mm, 4mm, and 6mm apical to CEJ in Facio-Palatal Direction.

Following the application of the inclusion criteria, a total of 350 cone beam computed tomography (CBCT) images were selected for analysis. The study population consisted of 130 (37.1%) male and 220 (62.9%) female participants between the ages of 18 to 60 years, with a mean age of 39.0 ± 12.6 years. The study evaluated 2100 anterior teeth in the maxillary region, including 700 central incisors, 700 lateral incisors, and 700

canines. It is worth noting that the labial bone thickness was greater at 4mm apical to CEJ than at 6mm for all examined locations. Moreover, the maxillary central incisors showed the highest values among the examined regions, while the lateral incisor regions showed the thinnest labial bone as illustrated in Table 1.

In terms of the comparison between gender and labial bone thickness, there was no statistically significant difference found between males and females (P value > 0.05), as illustrated in Figure 2.

Significant differences were observed between the right and left sides for all examined locations (P value <0.05) (Figure 3).

In terms of the correlation between age groups and the examined parameters, a highly statistically significant difference found (P value = 0.000), as shown in Figure 4. This indicates that there are significant variations in the labial bone thickness in maxillary anterior teeth among different age groups.




Figure No. 4: Frequency distribution between age groups at various parameters.

DISCUSSION

This study aimed to measure the thickness of the labial alveolar bone in the six maxillary anterior teeth of the

population in Peshawar, Pakistan. Cone beam computed tomography (CBCT) is commonly used to assess the bone volume and morphology before tooth extraction, to ensure adequate knowledge for future implant placement⁽⁷⁾. CBCT has a good reputation for image clarity and linear measurement accuracy at a lower radiation dosage compared to standard CT⁽⁸⁾. According to the current study analyses the mean labial alveolar bone thickness measured at 2mm, 4mm, and 6mm showed greater values at 4mm followed by 6mm apical to CEJ. The maxillary central incisors showed the uppermost values among the regions examined, followed by canine, but at all the examined parameters the thinnest labial bone was found at the lateral incisor region. These results are consistent with those seen in the majority of investigations where in maxillary anterior region the thickness of labial alveolar bone values below 1 mm were noted. According to the study by AlAli et al., 2022, more than 80% of the sites had an LBT of less than 1 mm. Similar results were found in earlier studies by dos Santos et al., 2019; Gakonyo et al., 2018, with 76% to 89% of sites having LBT of not more than 1 mm at the central incisor in maxillary region^(2,9,10). These results are reliable with the results of the current study, which found that all of the evaluated central incisors had an LBT of less than 1 mm. According to the research by H. Sheerah et al. 2019, one-third of entirely canines and nearly half of entirely incisors have bone wall thin, of less than 1mm⁽¹¹⁾. These findings support our understanding of front maxillary sites with labial alveolar bone thicknesses of less than 1mm.

Additionally in-depth investigation of our findings demonstrated a tendency towards the existence of an increased thickness of labial alveolar bone at 4mm apical to CEJ when compared to 2mm and 6mm labial bone thickness. The data published by H. Sheerah et al. 2019 reported that the apical 3rd of the labial alveolar bone give the idea to have the most favorable thickness, which is opposite to the current study and the study done by AlAli et al., 2022; El Nahass & N. Naiem, 2015; Ghassemian et al., 2012.

In our analysis, gender did not appear to have an impact on labial alveolar bone thickness. This appears to be consistent with the outcomes of other published studies^(9,12). There have also been conflicting reports about the effect of gender on labial alveolar bone thickness, with some research reporting an increased thickness in men^(11,13). The variance of the sample and the population of interest differ, which might lead to contradictory results⁽¹¹⁾.

Additionally, unlike previous research by AlTarawneh et al., 2018; Sheerah et al., 2019, discovered significant changes in the labial alveolar bone thickness between the right and the left sides in the current study. Our research suggests that aging affects labial bone thickness, which is consistent with prior studies that

found that aging was related to lower labial alveolar bone thickness values^(2,10,13). However, other research found no association between the age and the labial bone thickness^(9,11).

CONCLUSION

In light of the results and limitations of this study, the following conclusions can be drawn: The thickness of the labial alveolar bone in the maxillary anterior teeth demonstrated a significant increase with age (P-value = 0.000). The greatest thickness was observed 4 mm apical to the CEJ, with the maxillary central incisors showing the highest values among the examined regions. However, the lateral incisor regions had the thinnest labial bone at the bone crest level and at 2mm, 4mm, and 6mm apical to the CEJ. The results of this study suggested that there were no statistically significant differences in labial alveolar bone thickness between males and females (P-value <0.05). Nevertheless, a highly statistically significant difference was found between the right and left sides of maxillary anterior teeth (P-value <0.05).

Author's Contribution:

Concept & Design of Study:	Naheed Imran
Drafting:	Asma Sattar, Imran Khattak
Data Analysis:	Sana Arbab, Munawar Aziz, Syed Amjad Shah
Revisiting Critically:	Naheed Imran, Asma Sattar
Final Approval of version:	Naheed Imran

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: PRIME/IRB/2021-357 dated 10.09.2021

REFERENCES

1. Rojo-Sanchis J, Soto-Peñaloza D, Peñarrocha-Oltra D, Peñarrocha-Diago M, Viña-Almunia J. Facial alveolar bone thickness and modifying factors of anterior maxillary teeth: a systematic review and meta-analysis of cone-beam computed tomography studies. *BMC Oral Health* 2021;21(1):143.
2. Gakonyo J, Mohamedali A, Kabubei E. Cone Beam Computed Tomography Assessment of the Buccal Bone Thickness in Anterior Maxillary Teeth: Relevance to Immediate Implant Placement. *Int J Oral Maxillofacial Implants* 2018;33:880–7.
3. AlTarawneh S, AlHadidi A, Hamdan AA, Shaqman M, Habib E. Assessment of Bone Dimensions in the Anterior Maxilla: A Cone Beam Computed Tomography Study: Assessment of

- Maxillary Bone Dimensions. *J Prosthodontics* 2018;27(4):321–8.
4. Porto OCL, Silva BS de F, Silva JA, Estrela CR de A, Alencar AHG de, Bueno M dos R, et al. CBCT assessment of bone thickness in maxillary and mandibular teeth: an anatomic study. *J Appl Oral Sci* 2020 Feb 7 [cited 2023 Jan 18];28. Available from: <http://www.scielo.br/j/jaos/a/T763GbCy67X9Tfw5X8bVbJs/?format=html&lang=en>
 5. Xu D, Xie C, Yu H, Zhang Z, Zheng J, Xu S. Evaluation of factors affecting alveolar ridge height and facial bone thickness in Chinese maxillary central incisors by cone beam CT. *J Dent Sciences* 2021;16(1):229–35.
 6. Ahmed DF, El Beshlawy DM. Assessment of anterior maxillary alveolar bone dimensions and morphology for immediate implant planning: A retrospective study using Planmeca Promax CBCT unit. *Egypt Dent J* 2019;65(Issue 2-April (Oral Medicine, X-Ray, Oral Biology&Oral Pathology):1267–78.
 7. Di Stefano DA, Arosio P, Capparè P, Barbon S, Gherlone EF. Stability of Dental Implants and Thickness of Cortical Bone: Clinical Research and Future Perspectives. *A Systematic Review. Materials* 2021;14(23):7183.
 8. Chandak DM. Comparative Evaluation Of Accuracy Of Dental Operating Microscope With Conventional Vernier Caliper In Conservation Of Tooth Structure For Root Canal Treatment. *Clin Med* 2020;7(07):7.
 9. AlAli F, Atieh MA, Hannawi H, Jamal M, Harbi NA, Alsabeeha NHM, et al. Anterior Maxillary Labial Bone Thickness on Cone Beam Computed Tomography. *Int Dent J* 2022 May 6 [cited 2023 Jan 3]; Available from: <https://www.sciencedirect.com/science/article/pii/S0020653922000703>
 10. dos Santos JG, Oliveira Reis Durão AP, de Campos Felino AC, de Faria de Almeida RMCL. Analysis of the Buccal Bone Plate, Root Inclination and Alveolar Bone Dimensions in the Jawbone. A Descriptive Study Using Cone-Beam Computed Tomography. *JOMR* 2019 Jun 30 [cited 2023 Jan 3];10(2). Available from: <http://www.ejomr.org/JOMR/archives/2019/2/e4/v10n2e4ht.htm>
 11. Sheerah H, Othman B, Jaafar A, Alsharif A. Alveolar bone plate measurements of maxillary anterior teeth: A retrospective Cone Beam Computed Tomography study, AlMadianh, Saudi Arabia. *The Saudi Dental J* 2019;31(4):437–44.
 12. Lim HC, Kang DU, Baek H, Hong JY, Shin SY, Chung JH, et al. Cone-beam computed tomographic analysis of the alveolar ridge profile and virtual implant placement for the anterior maxilla. *J Periodontal Implant Sci* 2019;49(5):299–309.
 13. A T, J T, B de BB, A G, Ab B, J C, et al. Buccal bone thickness of maxillary anterior teeth: A systematic review and meta-analysis. *J Clin Periodontol* 2020 Nov [cited 2023 Jan 3];47(11). Available from: <https://pubmed.ncbi.nlm.nih.gov/32691437>.

Delayed Presentation of Large Goitre, A Cross Sectional Study at a Tertiary Care Hospital in Karachi

Delayed
Presentation of
Large Goitre

Mariam Imran, Saad Abdul Razzaq, Zahid Mehmood, Ghansham Rawtani and
Hazrat Bilal Burki

ABSTRACT

Objective: To identify the contributing factors which lead to delayed presentation of goitres.

Study Design: Descriptive Cross Sectional Study

Place and Duration of Study: This study was conducted at the Department of General Surgery ward 25, from January 2022 to 31st August 2022.

Methods: All patients presenting to thyroid OPD with a history of thyroid goitre for at least 3 years were included. All data was recorded in a Performa.

Results: Out of 120 participants, 82.5% were male and the rest were females. Fear of surgery (31%) was the most common reason for delayed presentation, followed by advice from hakeem (20%) and advice from friends (12.8%). There was significant correlation between the educational and socio-economic status and time of presentation for intervention giving the p-value of <0.001.

Conclusion: Lack of education regarding the surgical management of thyroid diseases was found to be the most mentioned cause of delayed presentation followed by advice from hakeem/faith healers, peers and physicians.

Key Words: Giant goitre, delayed presentation, education level.

Citation of article: Imran M, Abdul Razzaq S, Mehmood Z, Rawtani G, Burki HB. Delayed Presentation of Large Goitre, A Cross Sectional Study at a Tertiary Care Hospital in Karachi. Med Forum 2024;35(1):47-49. doi:10.60110/medforum.350110.

INTRODUCTION

Goitre or a benign enlargement of a thyroid gland that can be asymptomatic in some patients but can cause compressive symptoms like dysphagia and dyspnoea in others¹. The incidence of goitre depends upon the iodine intake of the patient, familial background, iodine deficiency, goiterogens and radiation exposure². In long standing cases, nodularity develops frequently. It has been noted in the past that multinodular goitre remains asymptomatic for a longer period of time unless there is underlying malignancy³. Therefore, some people from developing countries usually ignore the enlarging thyroid gland and prefer no treatment until it becomes enlarged and cause symptoms⁴. In contrary to this, in the western world the incidence of thyroid carcinoma has rapidly increased in the recent time due to early presentation and better diagnostic modalities and early intervention, in case of benign smaller toxic thyroid

swellings are usually dealt with radio iodine or thyroxin or a combination of both but multinodular large goitre usually does not respond to medical management and requires surgical intervention^{5,6}. Further medical management include mainly iodine replacement, thyroid hormone replacement, thyroid hormone suppressive therapy, and radioactive iodine, whereas refractory to medical therapy in case of large goitre, surgical options are available. Even in experienced hands difficulties are usually encountered during thyroidectomy for the huge multinodular goitre and chances of injury to the vital structures can be expected including tracheal, oesophageal and recurrent laryngeal nerve and hematoma formation⁷. Furthermore, in the post-operative period, these patients can experience tracheomalacia and hypocalcaemia due to inability to preserve parathyroid glands⁸. To prevent these complications, early surgical intervention is advised. It has been observed in many high volume thyroid centres that people are presenting with huge thyroid glands with long standing histories despite being symptomatic⁹. Therefore, this study is conducted to find out the reason behind delayed presentation so that the awareness can be provided to the general population to consult with physicians for the evaluation of enlarging thyroid gland as soon as they observe it.

Department of General Surgery, Jinnah Postgraduate Medical Centre, Karachi.

Correspondence: Dr. Mariam Imran, General Surgery, Jinnah Postgraduate Medical Centre, Karachi.

Contact No: 03343624746

Email: mariam.imran93@gmail.com

Received: July, 2023

Accepted: September, 2023

Printed: January, 2024

METHODS

This descriptive cross sectional study was conducted in surgical ward 25, Jinnah Postgraduate Medical Centre,

Karachi, Pakistan, from January 2022 to 31st August 2022. Patients with large goitre at presentation having a history of at least 3years are included in the study. Patients with early presentation(less than 3 years), known thyroid malignancy, solitary thyroid nodule, are excluded from the study. Data was collected in a pre-designed proforma after taking written and informed consent. Data was analysed via SPSS 23 and P value of less than 0.01 was considered significant.

RESULTS

Total of 120 patients were included in the study as per the inclusion criteria, out of which 99(82.5%) were female and 21(17%) were male.

Table No.1: Percentile of patients

	Total no. of Patients	Percentile
Male	21	17%
Female	99	82.5%

Patient's socio-economic status was determined on the basis of monthly income and expenditure. Most of the patients belong to the middle class (63.3%) followed by lower class (34.2%).

Table No.2: Socioeconomic Status

Socioeconomic Status	Percentile
Middle Class	63.3%
Lower Class	34.2%)

Furthermore, majority of the patients were uneducated (53.3%), with only 31.7% of people who studied in school.

Table No.3: Percentile of patients with education and un-education

Total no. of Patients	Percentile
Uneducated	53.3%
Educated	31.7%

The average time of presentation was found to be 7-8years. Patients are divided in to four age groups i.e. <20years, 20-40years, 41-60years and >60 years. Around 48.3% patient belong to 20-40years group followed by 44.2% in age group of 41-60years. The reasons for delayed presentation included fear from surgery (31.7%), advise from hakeem/faith healer (20%), advise from physician(10.8%) and advise from relative/friends(12.5%). In addition to these, around 20% of patient did not give any reason for their late presentation. Significant correlation between the socio-economic and educational level and delayed presentation has been found with the p-value of less than 0.001. Simply stated, patients belonging to the lower socioeconomic status and who were uneducated were more likely to have delayed presentation. This could be due to a lack of awareness amongst the population regarding the surgical treatment of an enlarged thyroid gland.

DISCUSSION

Massively enlarged thyroid glands are becoming infrequent in the western world, however, it is still prevalent in some geographical locations. It can cause compressive symptoms including dysphagia, dyspnoea and hoarseness of voice¹⁰. Also, the proposed treatment option is surgical and it requires high level of surgical expertise in order to prevent disastrous complications including permanent hypocalcaemia and tracheostomy in case of bilateral RLN injury.

A study conducted in Baghdad showed the incidence of differentiated thyroid carcinoma to be 21.7% in patients with long standing MNGs¹¹. This study finding was also close to the findings presented in the study conducted in Saudi Arabia by Al-Sala

mah et al. showing 21.3% incidence of differentiated thyroid cancer in MNG detected by FNA and confirmed by subsequent histopathology¹². One study suggested that , patient with papillary cancer have better survival upto 75% and the incidence of complications were found to be on lower site , this study took a brief review of 200 cases over the period of 30 years and show despite delayed presentation adverse symptoms appear late for which patient seek medical attention.¹³ another cross sectional study conducted in India to compare individuals that are iodine deficient or ones having normal iodine intake and it was found that a significant proportion of population with insufficient iodine intake had thyroid related disorders and total prevalence of goitre was 12.2%.¹⁴

Another study conducted in US based on surveillance among 318,318 participant undergoing various surgeries, the incidence of refusal was found to be 3.5% and mainly among blacks, advance age, unmarried and uninsured individuals. However, racial and ethnic injustice in healthcare domain is one the major talked issue in United states up till date but in our setup still opting for live saving procedures prior to marriages is major dilemma¹⁵ Furthermore, the transformation of papillary thyroid carcinoma in to anaplastic thyroid carcinoma is well documented in the literature. Although, it only comprises 3.8% of the thyroid carcinomas, it is the most aggressive form and have a very high mortality rate. Keeping in mind the high incidence of malignancy in MNG and anaplastic transformation, early evaluation of thyroid nodules is indicated¹⁶ still the incidence of conversion of papillary to anaplastic carcinoma with life threatening complications are minimal as it requires series of further genetic mutations¹⁷, that seems one of the another possibility of delayed presentation, as individuals are unaware of complications that results in early search for medical guidance. Unfortunately, in some eastern part of the world, people are found to be hesitated to seek medical attention for a longer time for thyroid diseases.

This study conducted in one of the surgical unit of Jinnah post graduate medical centre has identified few reasons for the late presentation of large multinodular goitres. Amongst these, fear from the surgery is the most common reason by the patients followed by advise given by faith healers, friends/family and even by some physicians. On top of it, an important co-relation was

found between the late presentation and socio-economic status and educational level. The majority of the studied population belonged to the low socioeconomic status and had a very low level of education, therefore, most people were not even aware about the consequences associated with surgery of large thyroid gland and high incidence of carcinoma in MNG.

CONCLUSION

Lack of education regarding the surgical management of thyroid diseases was found to be the most mentioned cause of delayed presentation followed by advice from hakeem/faith healers, peers and physicians. This is mandating the organization of awareness programs amongst the general population and different health care professionals to highlight the importance of early evaluation of thyroid diseases which in turn will potentially lead to a decrease the morbidity associated with the total thyroidectomy for huge multinodular goitre, especially those with the retrosternal extension.

Acknowledgments: Authors contributions: Dr. Saad Abdul Razzaq contributed to the conception and design of the study acquisition of data, Dr Mariam Imran help in drafting the manuscript and interpretation of data..., Dr. Zahid Mehmood, , Dr. Ghansham, Dr Hazrat Bilal Burki contributed to the acquisition, analysis and interpretation of data along. All authors approved the final version of the manuscript.

Author's Contribution:

Concept & Design of Study:	Mariam Imran
Drafting:	Saad Abdul Razzaq, Zahid Mehmood
Data Analysis:	Ghansham Rawtani, Hazrat Bilal Burki
Revisiting Critically:	Mariam Imran, Saad Abdul Razzaq
Final Approval of version:	Mariam Imran

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No. F.2-81/2021-GENL/70132/
JPMC Dated 30.12.2021

REFERENCES

- Lockhart ME, editor. Endocrine Imaging, An Issue of Radiologic Clinics of North America, E-Book. Elsevier Health Sciences. Bron LP, O' Brien CJ. 2020 Oct 23.
- Barczyński M, Stopa-Barczyńska M. Hemithyroidectomy for benign euthyroid asymmetric nodular goitre. *Best Pract Res Clin Endocrinol Metab* 2019;33(4):101288.
- Censi S, Salmaso L, Ceccato F, Manso J, Fedeli U, Saia M, et al. Hyperthyroidism incidence in a large population-based study in northeastern Italy. *Endocrine Connections* 2023;12(12).
- Illouz F, Chanson P, Sonnet E, Brue T, Ferriere A, Raffin Sanson ML, et al. Somatostatin receptor ligands induce TSH deficiency in thyrotropin-secreting pituitary adenoma. *Eur J Endocrinol* 2021;184(1):1-8.
- Maniakas A, Davies L, Zafereo ME. Thyroid disease around the world. *Otolaryngologic Clinics North Am* 2018;51(3):631-42.
- Orloff LA, Parangi S. History of Thyroid Surgery in the Last Century. *Thyroid* 2023;33(9):1029-38.
- Chen AY, Bernet VJ, Carty SE, Davies TF, Ganly I, Inabnet III WB, Shaha AR. American Thyroid Association statement on optimal surgical management of goiter. *Thyroid* 2014;24(2):181-9.
- Canu GL, Medas F, Cappellacci F, Soddu C, Romano G, Erdas E, et al. Intact parathyroid hormone value on the first postoperative day following total thyroidectomy as a predictor of permanent hypoparathyroidism: a retrospective analysis on 426 consecutive patients. *Endokrynologia Polska* 2022 Feb 8.
- Ghafouri A, Hawsawi H, Alanzi A, Alzaidi S. The Influence of Preoperative Thyroid Gland Volume On Intraoperative Time and Postoperative Complications for Total Thyroidectomy Patients. *J Pathol Res Reviews Reports SRC/JPR-147*. DOI: doi.org/10.47363/JPR/2022 (4). 2022;138:2-4.
- Imran M, Mehmood Z, Baloch MN, Altaf S. Endoscopic thyroid lobectomy vs Conventional open thyroid lobectomy. *Pak J Med Sciences* 2020;36(4):831.
- Ghadhban BR. Incidence of differentiated thyroid carcinoma in multinodular goiter patients. *Int J Surg Open* 2018;15:18-24.
- Al-Salamah SM, Kamran Khalid FC, Bismar HA, CAES C. nodular goiter. *Saudi Med J* 2002;23(8):947-52.
- Nicolson NG, Carling T. Reply to "Differences in the Impact of Age on Mortality in Well-Differentiated Thyroid Cancer". *Ann Surg Oncol* 2019;26(2):697.
- Menon U, Sundaram KR, Unnikrishnan AG, Jayakumar RV, Nair V, Kumar H. High prevalence of undetected thyroid disorders in an iodine sufficient adult south Indian population. *J Ind Med Assoc* 2009;107(2):72-7.
- Hu X, Ye H, Yan W, Sun Y. Factors Associated With Patient's Refusal of Recommended Cancer Surgery: Based on Surveillance, Epidemiology, and End Results. *Front Public Health* 2022; 9:785602.
- Amacher AM, Goyal B, Lewis Jr JS, El-Mofty SK, Chernock RD. Prevalence of a hobnail pattern in papillary, poorly differentiated, and anaplastic thyroid carcinoma: a possible manifestation of high-grade transformation. *Am J Surgical Pathol* 2015;39(2):260-5.
- Capdevila J, Mayor R, Mancuso FM, Iglesias C, Caratù G, Matos I, et al. Early evolutionary divergence between papillary and anaplastic thyroid cancers. *Annals Oncol* 2018;29(6): 1454-60.

Percutaneous Nephrolithotomy: A Single Center Experience of 162 Cases of Standard Percutaneous Nephrolithotomy

Zohair, Akhtar Nawaz, Waqas, Siddique Akbar and Sardar Alam

ABSTRACT

Objective: This study aims to share our experience of Percutaneous nephrolithotomy regarding its safety and efficacy for renal stones in a tertiary care hospital.

Study Design: A Retrospective Study

Place and Duration of Study: This study was conducted at the department of Urology Unit IKD Peshawar from July 2021 to July 2022.

Methods: This retrospective study included all those patients who underwent Percutaneous nephrolithotomy in the Urology Unit. The data was collected retrospectively from the hospital record system and analysis was done with IBM SPSS version 20.

Results: A total of 162 patients including 67.3% male and 32.7% female were part of this study. Fifty-five percent of the patients had a previous history of renal stones. Pre-operative Ultrasound findings were multiple stones in 54.3%, single stones in 40.1%, staghorn stones in 3.7%, a duplex system in 1.2 %, and horseshoe kidney in 0.6% (n=1). Sixty percent had stone sizes ranging from 15 mm to 30mm, 22.8% had stone sizes less than 15mm and 16.7% had stone sizes greater than 30mm. In the majority, the location of stones was renal pelvis (46.3%), lower pole 18.5%, Pelvis plus lower pole 16% and staghorn 7.4%. Pre-operative mean Hemoglobin was 12.9g/dl and post-operative 11.8g/dl with a mean drop of 1.1g/dl. Complete stone clearance was achieved in 86.4%, partial stone clearance in 11.7% and the procedure was abandoned in 1.9% (n=3). The majority of the procedure was uneventful (87.7%) while 12.3% had complications in the form of bleeding requiring blood transfusion 5.5%(n=9), sepsis 3.08%, pleural injury 1.2%, peritoneal injury and stone fragment migration to ureter with subsequent ureteral obstruction 0.61% (n=1) each. One patient died of sepsis with multi-organ failure. The mean hospital stay was 3.06 days with a minimum of 2 days and a maximum of 11 days.

Conclusion: Percutaneous nephrolithotomy is an effective way of treating a wide range of stone sizes, in different renal locations, in anomalous kidneys with a high clearance rate and acceptable complication rate.

Key Words: Renal stone, Percutaneous nephrolithotomy

Citation of article: Zohair, Nawaz A, Waqas, Akbar S, Alam S. Percutaneous Nephrolithotomy: A Single Center Experience of 162 Cases of Standard Percutaneous Nephrolithotomy. Med Forum 2024;35(1):50-53. doi:10.60110/medforum.350111.

INTRODUCTION

The landscape of urological interventions has witnessed a tremendous evolution, with percutaneous nephrolithotomy (PCNL) rising as a pivotal player in the complete control of renal stones¹. As renal stone incidence continues to pose a considerable burden on global healthcare, exploring the nuances of procedural safety and efficacy becomes imperative².

This retrospective observe serves as a meticulous exploration of the reports and effects associated with

PCNL in the context of a tertiary care clinic³. Renal stones, a recurrent and regularly debilitating condition, necessitate nuanced tactics for their powerful decision. Among the array of available interventions, PCNL has garnered interest for its versatility in addressing a diverse spectrum of stone sizes, places, and patient profiles⁴. The complicated stability between reaching most appropriate stone clearance and mitigating capacity complications underscores the want for a thorough investigation into the procedural dynamics⁵. The number one objective of this take a look at is to provide a comprehensive assessment of the protection and efficacy of PCNL, drawing insights from a retrospective analysis of instances spanning from July 2021 to July 2022 in the Urology Unit of our tertiary care clinic⁶. By delving into the demographic traits, stone profiles, and postoperative effects, this observe endeavors to make contributions treasured insights to the existing body of information surrounding PCNL⁷. Through meticulous statistics series and subsequent evaluation making use of IBM SPSS model

Department of Urology and Renal Transplant, IKD Peshawar.

Correspondence: Akhtar Nawaz, Assistant Professor Urology and Renal Transplant, IKD, Peshawar.

Contact No: 0333 9382176

Email: nawaz157@gmail.com

Received: August, 2023

Accepted: November, 2023

Printed: January, 2024

20, this take a look at aspires to provide a nuanced know-how of PCNL's position in coping with renal stones. The inclusion of numerous cases, encompassing various stone sizes, places, and affected person histories, pursuits to present a holistic attitude at the technique's applicability and effectiveness⁸.As the introduction units the level for the approaching exploration, the look at anticipates losing mild on PCNL as a cornerstone inside the armamentarium of urological interventions. By navigating via the intricacies of patient demographics, stone characteristics, and procedural results, this investigation strives to make a contribution substantively to the existing literature, offering precious insights that can inform scientific choice-making and decorate the general first-rate of take care of people grappling with renal stones⁹.

METHODS

This retrospective observe encompasses sufferers undergoing Percutaneous Nephrolithotomy (PCNL) at the Urology Unit in our tertiary care sanatorium from July 2021 to July 2022. The comprehensive records, which includes demographic info, stone traits, and postoperative effects, become meticulously retrieved from clinic facts. Analysis turned into conducted using IBM SPSS model 20, ensuring statistical robustness. The chosen timeframe provides a image of current PCNL practices, facilitating an in-intensity exam of its protection and efficacy in managing renal stones. This methodological method ambitions to provide a treasured contribution to the existing understanding base surrounding PCNL effects in a actual-global clinical setting.

Data collection: Patient information, comprising demographics, stone features, and postoperative results, become systematically amassed thru a retrospective overview of hospital data. The look at period, spanning from July 2021 to July 2022, ensured a representative photo of Percutaneous Nephrolithotomy (PCNL) practices. This meticulous information collection bureaucracy the foundation for a strong analysis of PCNL outcomes in our tertiary care

Statically analysis:The accumulated statistics underwent rigorous statistical analysis the usage of IBM SPSS model 20. Descriptive records had been employed to explain patient demographics, stone traits, and postoperative results. This analytical method affords a quantitative framework for comprehensively comparing the safety and efficacy of Percutaneous Nephrolithotomy (PCNL) in our tertiary care health facility throughout the specified timeframe.

RESULTS

In this examine, a cohort of 162 sufferers underwent Percutaneous Nephrolithotomy (PCNL), with

67.3% men and 32.7% women. Stone characteristics found out diverse sizes (15 mm to 30 mm in 60% of cases), varied locations (predominantly renal pelvis), and 55% with a records of renal stones. PCNL proven a high efficacy with complete stone clearance in 86.4%, partial clearance in 11.7%, and abandonment in 1.9%. Complications, encountered in 12.3% of cases, protected bleeding (5.5%), sepsis (3.08%), and pleural harm (1.2%). Notably, one patient succumbed to sepsis with multi-organ failure. The imply health facility stay changed into 3.06 days, putting forward PCNL as an powerful and appropriate intervention for renal stones.

Table No. 1: Patient Demographics

Parameter	Total Patients (n=162)
Gender	
- Male	67.3%
- Female	32.7%
Age (Mean ± SD)	

Table No. 2: Stone Characteristics

Parameter	Distribution (%)
Stone History	55%
Stone Size	
- 15 mm to 30 mm	60%
- <15 mm	22.8%
- >30 mm	16.7%
Stone Location	
- Renal Pelvis	46.3%
- Lower Pole	18.5%
- Pelvis + Lower Pole	16%
- Staghorn Stones	7.4%
Other Anomalies	

Table No. 3: Pre-operative Hemoglobin Levels

Parameter	Mean ± SD
Pre-operative Hemoglobin	12.9g/dl

Table No. 4: Stone Clearance Rates

Parameter	Clearance Rate (%)
Complete Stone Clearance	86.4%
Partial Stone Clearance	11.7%
Abandoned Procedures	1.9%

Table No. 5: Complications and Outcomes

Complications	Incidence (%)
Bleeding	5.5%
Sepsis	3.08%
Pleural Injury	1.2%
Peritoneal Injury	0.61% (n=1)
Stone Migration to Ureter	0.61% (n=1)
Mortality	0.61% (n=1)
Mean Hospital Stay	3.06 days

DISCUSSION

The results of this observe shed light at the protection and efficacy of Percutaneous Nephrolithotomy (PCNL) within the context of renal stone control¹⁰. The dialogue will delve into key elements, integrating insights from current literature for context and evaluation. The male predominance determined in our cohort is consistent with established literature highlighting a higher occurrence of renal stones in males (Litwin et al., 2007)¹⁰. This gender disparity underscores the significance of thinking about demographic factors in the evaluation and management of renal stones, aligning with broader epidemiological traits. The diverse distribution of stone sizes and places mirrors the multifaceted nature of renal stone displays. Previous research, such as those by using Assimos et al. (2016)¹¹, emphasize the importance of tailoring interventions based on man or woman stone profiles. The occurrence of fifty five% with a history of renal stones underscores the recurrent nature of the condition, warranting interest to preventive techniques. The excessive complete stone clearance price of 86.Four% aligns with the efficacy confirmed in research by using Preminger et al. (2007)². However, the eleven.7% partial clearance fee warrants attention, emphasizing the want for lengthy-term observe-up to cope with capacity residual fragments and save you recurrence¹². These findings underscore the need for endured refinement of PCNL strategies to optimize stone clearance. The located headaches, specially bleeding (5.5%), sepsis (3.08%), and mortality (0.Sixty one%), resonate with current literature emphasizing the capability risks related to PCNL (Lopes et al., 2017)¹². Vigilant intraoperative and postoperative management is vital to mitigate these dangers. This study contributes to the continued speak concerning the stability between procedural effectiveness and safety.¹³⁻¹⁵ The imply clinic live of 3.06 days aligns with studies advocating for shorter hospitalization periods without compromising affected person results (Chen et al., 2016)¹⁸. This underscores the feasibility of PCNL as a minimally invasive method with a extraordinarily speedy postoperative recuperation, contributing to fee-effectiveness and progressed affected person experience¹⁶. In end, the records supplied on this look at contribute to the developing body of evidence assisting the function of PCNL within the management of renal stones¹⁸. The nuanced interaction between patient demographics, stone traits, and procedural effects necessitates a customised technique, aligning with modern-day urological guidelines (European Association of Urology, 2021)¹⁷. However, the look at's retrospective nature and potential selection bias warrant validation through larger potential studies for a greater comprehensive expertise of PCNL outcomes¹⁹.

CONCLUSION

This study illuminates percutaneous nephrolithotomy's effectiveness in dealing with renal stones, showcasing excessive clearance rates and appropriate headaches. The nuanced exploration of patient demographics, stone profiles, and procedural outcomes adds treasured insights to the present literature. Acknowledging PCNL's function as a cornerstone in urological interventions, this research contributes substantively to scientific choice-making. The commitment to methodological rigor and numerous case inclusion enhances the have a look at's relevance, urging a persisted consciousness on customized approaches. Overall, the findings verify PCNL as a vital and powerful intervention in the comprehensive management of renal stones.

Acknowledgement: We would like to thank the hospitals administration and everyone who helped us complete this study.

Author's Contribution:

Concept & Design of Study:	Akhtar Nawaz
Drafting:	Zohair
Data Analysis:	Waqas
Revisiting Critically:	Siddique Akbar
Final Approval of version:	Sardar Alam

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.212 dated 20.02.2021

REFERENCES

1. Türk C, Knoll T, Petrik A, Sarica K, Skolarikos A, Straub M, et al. Guidelines on Urolithiasis. Eur Assoc Urol 2015. Retrieved from <https://uroweb.org/guideline/urolithiasis/>
2. Preminger GM, Tiselius HG, Assimos DG, Alken P, Buck C, Gallucci M, et al. 2007 guideline for the management of ureteral calculi. Eur Urol 2007;52(6):1610-1631.
3. Litwin MS, Saigal CS. Urologic Diseases in America Project. Urologic diseases in America project: Analytical methods and principal findings. J Urol 2007;177(3):1151-1156.
4. Tzelvels L, Geraghty RM, Hughes T, Juliebø-Jones P, Somani BK. Innovations in Kidney Stone Removal. Research Reports Urol 2023;Dec 31:131-9.
5. Amparore D, Campi R, Checcucci E, Sessa F, Pecoraro A, Minervini A, et al. Forecasting the future of urology practice: a comprehensive review of the recommendations by international and European associations on priority procedures during the COVID-19 pandemic. Eur Urol Focus 2020;6(5):1032-48.

6. Qin P, Zhang D, Huang T, Fang L, Cheng Y. Comparison of mini percutaneous nephrolithotomy and standard percutaneous nephrolithotomy for renal stones > 2cm: a systematic review and meta-analysis. *Int Braz J Urol* 2022;48:637-48.
7. Kallidonis P, Tsaturyan A, Lattarulo M, Liatsikos E. Minimally invasive percutaneous nephrolithotomy (PCNL): Techniques and outcomes. *Turkish J Urol* 2020;46(Suppl 1):S58.
8. Jiao B, Luo Z, Huang T, Zhang G, Yu J. A systematic review and meta analysis of minimally invasive vs. standard percutaneous nephrolithotomy in the surgical management of renal stones. *Exp Therapeutic Med* 2021;21(3):1.
9. Deng J, Li J, Wang L, Hong Y, Zheng L, Hu J, Kuang R. Standard versus mini-percutaneous nephrolithotomy for renal stones: a meta-analysis. *Scandinavian J Surg* 2021;110(3):301-11.
10. Litwin MS, Saigal CS. Urologic Diseases in America Project. Urologic diseases in America project: Analytical methods and principal findings. *J Urol* 2007;177(3):1151-1156.
11. Assimos D, Krambeck A, Miller NL, Monga M, Murad MH, Nelson CP, et al. Surgical management of stones: American urological association/endourological society guideline, PART I. *J Urol* 2016;196(4):1153-1160.
12. Lopes T, Sangam K, Alken P, Barroilhet G, Averch TD, Bariol SV, et al. Percutaneous nephrolithotomy worldwide: results from the CROES percutaneous nephrolithotomy global study. *J Urol* 2017;188(2):306-311.
13. Sadiq AS, Atallah W, Khusid J, Gupta M. The surgical technique of mini percutaneous nephrolithotomy. *J Endourol* 2021;35(S2):S-68.
14. Kandemir E, Savun M, Sezer A, Erbin A, Akbulut MF, Sarılar Ö. Comparison of miniaturized percutaneous nephrolithotomy and standard percutaneous nephrolithotomy in secondary patients: a randomized prospective study. *J Endourol* 2020;34(1):26-32.
15. Sahalevych A, Sergiychuk R, Ozhohin V, Vozianov O, Khrapchuk A, Dubovyi Y, et al. Mini-percutaneous nephrolithotomy in surgery of nephrolithiasis. *Ukrainian J Nephrol Dialysis* 2021;(3 (71)):44-52.
16. Feng D, Hu X, Tang Y, Han P, Wei X. The efficacy and safety of miniaturized percutaneous nephrolithotomy versus standard percutaneous nephrolithotomy: A systematic review and meta-analysis of randomized controlled trials. *Investigative and Clinical Urol* 2020;61(2):115-26.
17. Zeng G, Cai C, Duan X, Xu X, Mao H, Li X, et al. Mini percutaneous nephrolithotomy is a noninferior modality to standard percutaneous nephrolithotomy for the management of 20–40 mm renal calculi: a multicenter randomized controlled trial. *Eur Urol* 2021;79(1):114-21.

Complications of Ultrasound Guided Percutaneous Nephrostomy in Adults: A retrospective Study

Ultrasound
Guided
Percutaneous
Nephrostomy

Akhtar Nawaz, Waqas, Zohair, Siddique Akbar and Sardar Alam

ABSTRACT

Objective: To identify the indications for percutaneous nephrostomy (PCN) placement and assess the success rate as well as the incidence of various complications associated with ultrasound-guided percutaneous nephrostomy.

Study Design: A retrospective Study

Place and Duration of Study: This study was conducted at the Institute of Kidney Diseases, Peshawar from April 2021 to April 2022.

Methods: This retrospective look at carried out at the Institute of Kidney Diseases, Peshawar, focused on patients who underwent ultrasound-guided percutaneous nephrostomy (PCN) for obstructive uropathy between April 2021 to April 2022. Data from HIS records and Urology Department registers have been analyzed. The inclusion standards encompassed patients aged >sixteen years, with exclusion standards involving incomplete documentation, pregnancy, and PCN for motives apart from obstructive uropathy. Statistical evaluation turned into achieved using IBM SPSS, calculating suggest, preferred deviation, frequency, and percentage. The take a look at layout aimed to offer complete insights into PCN-associated headaches, contributing to the development of urological care.

Results: The observe included 1702 sufferers with an average age of 38.50 ± 14 . Seventy eight years. Successful PCN placement changed into executed in ninety six.1%, with 67 patients requiring more than one try. Complications have been mentioned in 18.9%, comprising 17.6% minor and 1.3% major headaches. Macroscopic hematuria changed into the maximum not unusual (6.9%), followed by way of PCN dislodgment (1.9%) and tube occlusion (1.8%). Statistical analysis found out associations among complications and variables including gender, number of attempts, and age. The consequences contribute valuable insights into the efficacy and safety of ultrasound-guided PCN, assisting in refining procedural protocols for top-rated affected person care.

Conclusion: This study underscores the effectiveness and protection of ultrasound-guided percutaneous nephrostomy in dealing with obstructive uropathy. With urinary stones diagnosed as a important motive, the findings contribute to enhancing affected person care and procedural protocols at the Institute of Kidney Diseases, Peshawar, Pakistan.

Key Words: Obstructive Uropathy, Percutaneous Nephrostomy, Complications

Citation of article: Nawaz A, Waqas, Zohair, Akbar S, Alam S. Complications of Ultrasound Guided Percutaneous Nephrostomy in Adults: A retrospective Study. Med Forum 2024;35(1):54-57. doi:10.60110/medforum.350112.

INTRODUCTION

Obstructive uropathy, a structural hindrance to the flow of urine ⁽¹⁾ accounts for 10% of the causes of acute renal failure and 4% of chronic end stage renal failure ⁽²⁾. The common causes of obstructive uropathy in adults are urinary stones, malignancy and iatrogenic benign stricture ⁽³⁾.

Department of Urology and Renal Transplant IKD Peshawar.

Correspondence: Zohair, Consultant Urology and Renal Transplant, IKD, Peshawar.

Contact No: 03332362625

Email: doctorzohair1@gmail.com

Received: July, 2023

Accepted: September, 2023

Printed: January, 2024

Percutaneous nephrostomy (PCN), a minimally invasive procedure, first documented by a urologist Dr. Willard Goodwin in 1955 is used to decompress the obstructed renal collecting system in order to preserve the renal function. This involves the insertion of a tube through the skin into the renal collecting system to drain the urine from the affected kidney ⁽⁴⁾. Percutaneous Nephrostomy (PCN) can be performed under fluoroscopic and ultrasound guidance. However, ultrasound guided PCN has lower complications and is as effective as fluoroscopic guided. Hence it is the most common method used to manage obstructive uropathy ⁽⁵⁾. After the placement of PCN, renal functions return to normal within 15 days in two-third of the patients with azotemia secondary to obstruction ⁽⁶⁾. Apart from relieving the urinary obstruction, which accounts for 85% to 95% of the cases, it is also utilized for other purposes. These include providing access to endourologic procedures,

diagnostic testing and urinary diversion⁽³⁾. The technical success of PCN varies depending on the clinical situation⁽⁷⁾. For obstructed dilated systems, the reported success rate is 96-100%. For nondilated collecting systems it is 82-96% and for complex stone disease it is 82-85%. Although the procedure is generally safe, some minor or major complications may occur. Quality Improvement Guidelines for Percutaneous Nephrostomy reported major complications in 0.1-10% of those undergoing the procedure. Similarly, high success rates of PCN have also been reported in Pakistan; however, the rates of complications vary from 4.66% to 17.3% which include both minor and major complications⁽⁸⁻¹¹⁾. The Institute of Kidney Diseases (IKD) in Peshawar is a leading provider of care for urological patients in the region. A study done from 2011-2012 at IKD⁽¹²⁾ reported that Urinary Tract Infection (UTI) was the most frequent complication (35%) after Percutaneous Nephrostomy (PCN), followed by macroscopic haematuria (21.4%). Catheter dislodgment occurred in 17% of the patients, while sepsis affected 13% of patients. Despite this valuable data, there remains a gap in understanding the current status and trends of PCN-related complications.¹³ Our study aims to address this gap by evaluating complications related to ultrasound-guided percutaneous nephrostomy in patients with obstructive uropathy, ultimately contributing to improved patient care and outcomes.

METHODS

A retrospective study was conducted at Institute of Kidney Diseases, Peshawar, Pakistan. After approval from Institutional Ethical & Review Board, the HIS records and registers of Department of Urology were accessed for patients who underwent Ultrasound guided Percutaneous Nephrostomy (PCN) between April 2021 to April 2022. The study focused on patient demographics, indications for PCN, and procedure-related complications. The inclusion criteria included all patients aged >16 years who underwent percutaneous nephrostomy (PCN) for obstructive uropathy. The exclusion criteria were patients with incomplete documentation, pregnant women, and who had PCN for reasons other than obstructive uropathy. Data was collected using self-made proforma and analysed using IBM SPSS for Windows version 26. Mean and standard deviation were calculated for age. Frequency and percentage was calculated for gender, cause of obstruction and complications. Chi-square test was done to determine the association between complications and variables of gender, age and number of attempts. The p-value of 0.05 or less was considered statistically significant.

Data collection: Data collection involved having access to Health Information System (HIS) records and Urology Department registers at the Institute of Kidney

Diseases, Peshawar. The study duration ranged from April 2021 to April 2022. Comprehensive information on patient demographics, symptoms for PCN, and headaches become systematically retrieved for evaluation.

Statistical Analysis: Statistical evaluation turned into carried out the usage of IBM SPSS for Windows model 26. Mean, popular deviation, frequency, and percent had been calculated. The Chi-square check determined associations among complications and gender, range of tries, and age, with a importance stage set at zero.05 or less.

RESULTS

There were 1702 patients in our study with a mean age of 38.50 ± 14.78 and ranged from 16 to 83 years. 995 (58.5%) were male and 707 (41.5%) were female. 1152 (67.7%) patients were aged 16 to 45, while 550 (32.3%) were aged 46 and above. 1628 unilateral and 74 bilateral percutaneous nephrostomies were performed. The indications for PCN included Urinary Stones in 1150 (67.6%), malignancy in 407 (23.9%), Pyonephrosis in 60 (3.5%), Stricture in 54 (3.2%), Iatrogenic Ureteric Injury in 16 (0.9%), Emphysematous Pyelonephritis in 10 (0.6%) and Ureteric Ligation in 05 (0.3%).




Figure No. 1: Bar chart showing causes of obstructive uropathy

Successful PCN was placed in the first attempt in 1635 (96.1%) patients and only 67 (3.6%) patients required more than single attempt. Complications were reported in 320 (18.9%) of patients. Among these 300 (17.6%) were minor complications and 22 (1.3%) were Major complications. 95.3% (286/300) of the minor complications occurred in patients who had their PCN placed in the first attempt, while 54.5% (12/22) of the major complications occurred in patients who required more than one attempt for PCN placement. Macroscopic haematuria was the most common complication which was experienced by 6.9% of the patients, while 1.9% and 1.8% of the patients experienced PCN dislodgement and occlusion, respectively. PCN tube occlusion with associated pain was reported in 1.2% of patients. Sepsis was observed

in 13 (0.8%) patients, and major haemorrhage requiring transfusion was noted in 2 (0.1%) patients. Out of the 300 minor complications, 176 (58.7%) were observed

in patients aged 16 to 45. Similarly, out of the 22 major complications, 14 (63.6%) were observed in patients aged 46 and above.

Table No. 1: Incidence of complications of Percutaneous Nephrostomy

		Urinary Stones	Malignancies	Pyonephrosis	Ureteric Stricture	Iatrogenic Ureteric Injury	Emphysematous Pyelonephritis	Ureteric Ligation	Incidence N (%)	
	No Complications	964	306	43	44	11	8	4	1380 (81.1)	
Minor Complications	Macroscopic Hematuria	75	29	5	6	2	0	0	117 (6.9)	
	Pain, Macroscopic Hematuria	29	14	1	0	0	1	0	46 (2.2)	
	PCN Tube Dislodgement	11	19	1	1	1	0	0	33 (1.9)	
	PCN Tube Occlusion	15	9	2	2	1	1	1	31 (1.8)	
	Pain	14	9	1	0	0	0	0	24 (1.4)	
	Urine Leak	17	3	1	0	0	0	0	21 (1.2)	
	Pain, PCN Tube Occlusion	11	7	2	0	1	0	0	21 (1.2)	
	Infection at PCN Insertion Site	3	1	2	0	0	0	0	6 (0.4)	
	Major Complications	Sepsis	5	7	0	1	0	0	0	13 (0.8)
		Puncture of adjacent organ	2	1	1	0	0	0	0	4 (0.2)
Pleural effusion		2	0	0	0	0	0	0	2 (0.1)	
Major Haemorrhage		1	1	0	0	0	0	0	2 (0.1)	
Urinoma		0	1	0	0	0	0	0	1 (0.1)	
Pneumothorax		1	0	0	0	0	0	0	1 (0.1)	

Table No. 2: Association of Gender, No. of attempts and Age with Complications

		Complications			p-value
		None N (%)	Minor N (%)	Major N (%)	
Gender	Male	807 (81.1)	173 (17.4)	15 (1.5)	0.627
	Female	573 (81.0)	127 (18.0)	7 (1.0)	
Attempt	1	1298 (83.3)	252 (16.2)	9 (0.6)	<0.001
	>1	82 (57.3)	48 (33.6)	13 (9.1)	
Age	16 to 45	968 (84.0)	176 (15.3)	8 (0.7)	<0.001
	46 and over	412 (74.9)	124 (23.5)	14 (2.5)	

DISCUSSION

In discussing the findings of the study on ultrasound-guided percutaneous nephrostomy (PCN) complications at the Institute of Kidney Diseases in Peshawar,

Pakistan, it is crucial to contextualize them in the existing literature. The examine aligns with previous studies, confirming the efficacy and protection of PCN, with a fulfillment price of 96.1%, steady with pronounced quotes ranging from eighty two% to a hundred%^(14,15,7,11). This reaffirms the reliability of PCN as a treasured intervention for obstructive uropathy. The occurrence of urinary stones because the leading cause of obstructive uropathy echoes findings from earlier research (thirteen, 12). Notably, the take a look at highlights a statistically vast association between age and complications, emphasizing the importance of age as a capability chance component. This is consistent with the observations that sixty three.6% of predominant complications befell in sufferers elderly forty six and above (Table 2).The suggested principal headaches (1.3%) fall within the mounted range of 0.1% to 10% outlined in fine development tips for PCN (eight). Noteworthy is the meticulous breakdown of headaches, with macroscopic hematuria being the most commonplace, going on in 6.9% of patients. This aligns with the findings of a preceding study carried out at IKD⁽¹²⁾. Comparisons with different regional research in Pakistan, which includes the one from 2011-2012, monitor a consistency in the prevalence of headaches, with urinary tract infections (UTI) being the maximum

common (35%) (thirteen). The current observe reinforces UTI as a significant complication, taking place in 21.6% of sufferers elderly 20-50 years. However, it's miles important to well known the limitations of this look at, inclusive of its retrospective nature and the absence of an assessment of long-time period complications. Furthermore, the impact of tube length and the degree of hydronephrosis on complications was no longer assessed, representing capability areas for future studies. Our study findings make contributions valuable insights into the headaches related to ultrasound-guided PCN in adults. The alignment with existing literature reinforces the reliability of PCN as a vital intervention for obstructive uropathy. The affiliation between age and complications emphasizes the need for individualized care in precise age groups, guiding future studies and refining procedural protocols.^{16,17}

CONCLUSION

This study establishes that urinary stones are the most prevalent cause of obstructive uropathy. Percutaneous Nephrostomy (PCN) is a comparatively simple, secure, and quick technique for temporary urinary diversion in cases of obstructive uropathy. This technique has a high success rate and results in fewer minor and major complications.

Acknowledgement: We would like to thank the hospitals administration and everyone who helped us complete this study.

Author's Contribution:

Concept & Design of Study: Akhtar Nawaz
 Drafting: Zohair
 Data Analysis: Waqas
 Revisiting Critically: Siddique Akbar
 Final Approval of version: Sardar Alam

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.122 dated 12.02.2021

REFERENCES

- Roth KS, Koo HP, Spottswood SE, Chan JC. Obstructive Uropathy: An Important Cause of Chronic Renal Failure in Children. *Clinical Pediatr* 2002;41(5):309-314.
- Siddiqui MM, McDougal WS. Urologic assessment of decreasing renal function. *Med Clin North Am* 2011;95(1):161-8.
- Dagli M, Ramchandani P. Percutaneous nephrostomy: technical aspects and indications. *Semin Intervent Radiol* 2011;28(4):424-37.
- Beiko D, Razvi H, Bhojani N, et al. Techniques – Ultrasound-guided percutaneous nephrolithotomy: How we do it. *Can Urol Assoc J* 2020;14(3):E104-E110.
- Yang YH, Wen YC, Chen KC, Chen C. Ultrasound-guided versus fluoroscopy-guided percutaneous nephrolithotomy: a systematic review and meta-analysis. *World J Urol* 2019;37(5):777-788.
- Pappas P, Stravodimos KG, Mitropoulos D, et al. Role of percutaneous urinary diversion in malignant and benign obstructive uropathy. *J Endourol* 2000;14(5):401-405.
- Pabon-Ramos WM, Dariushnia SR, Walker TG, Janne d'Othée B, Ganguli S, Midia M, et al. Quality Improvement Guidelines for percutaneous nephrostomy. *J Vascular Interventional Radiol* 2016;27(3):410-4.
- Haris MS, Hussain SA, Islam F. An evaluation of complications of ultrasound guided percutaneous nephrostomy in cases of obstructive uropathy. *Pak J Med Health Sciences* 2023;17(1):530-2.
- Ali SM, Mehmood K, Faiq SM, Ali B, Naqvi SA, Rizvi AU. Frequency of complications in image guided percutaneous nephrostomy. *J Pak Med Assoc* 2013;63(7):816-820.
- Ahmad I, Pansota MS. Comparison between double J (DJ) ureteral stenting and percutaneous nephrostomy (PCN) in obstructive uropathy. *Pak J Med Sciences* 2013;29(3). Doi:10.12669/pjms.293.3563
- Ahmad I, Pansota MS, Tariq M, et al. Complication of Percutaneous Nephrostomy (PCN) in Upper Obstructive Uropathy: Our Experience. *J Univ Med Dent Coll* 2014 Jun. 3 [cited 2023Dec.17];5(1):55-0. Available from: <https://jumdc.com/index.php/jumdc/article/view/275>
- Farooq K, Ahmad B, Shahab M. Frequency of common complications in patients after percutaneous nephrostomy for obstructive uropathy. *Khyber J Med Sciences* 2016;09(01): 72-6.
- Efesoy O, Saylam B, Bozlu M, Çayan S, Akbay E. The results of ultrasound-guided percutaneous nephrostomy tube placement for obstructive uropathy: A single-centre 10-year experience. *Turk J Urol* 2018;44(4):329-334.
- Montvilas P, Solvig J, Johansen TEB. Single-centre review of radiologically guided percutaneous nephrostomy using “mixed” technique: Success and complication rates. *Eur J Radiol* 2011;80(2):553-8.
- Naeem M, Jan MA, Ullah A, Ali L, Khan S, Haq A, et al. Percutaneous Nephrostomy for the Relief of Upper Urinary Tract Obstruction: An Experience with 200 Cases. *J Postgrad Med Inst* 2011 Oct 13 [cited 2024 Jan. 4];24(2). <https://jpmi.org.pk/index.php/jpmi/article/view/1056>
- Saeed K, Qureshi F, Hussain I, Tariq M. Frequency of complications of percutaneous nephrostomy in upper obstructive uropathy. *Sepsis* 2016;1:0-51.
- New FJ, Deverill SJ, Somani BK. Outcomes Related to Percutaneous Nephrostomies (PCN) in Malignancy-Associated Ureteric Obstruction: A Systematic Review of the Literature. *J Clin Med* 2021;10(11):2354.

Comparison of Treatment Response of Different Drugs in Common Migraine

Salman Khan, Imran Ahmad, Muhammad Jalil, Ibrar Saleem, Fawad Ahmad and
Khuram Haq Nawaz

ABSTRACT

Objective: The study was carried out to assess and contrast the efficacy of different pharmacological interventions in individuals who have received the diagnosis of common migraine.

Study Design: A longitudinal observational cohort research study

Place and Duration of Study: This study was conducted at the Department of Neurology, Pak Emirates Military Hospital in Rawalpindi from Jan 2022 to Jan 2023.

Methods: The cohort comprised 234 individuals diagnosed with common migraine according to the criteria of International Classification of Headache Disorders (ICHD-3). The medication regimens of the patients, which included Tricyclic antidepressants, Topiramate, beta-blockers, calcium channel blockers and antiepileptics, were utilized to classify them. The evaluation of treatment response was conducted at one, three, and six months after treatment using Migraine Treatment Response Score (MTRS).

Results: The study revealed that females comprised the majority at 65.8%. The average age of the participants was 40 ± 12 years. Over the course of six months, Tricyclic antidepressants demonstrated the most substantial enhancement in MTRS, as evidenced by scores increasing from 4.5 ± 1.2 to 5.8 ± 1.0 ($p < 0.05$). Significant declines in frequency and intensity of attacks were noted in response to Tricyclic antidepressants: the former decreased by 4 ± 2 to 2 ± 1 , while latter decreased by 7 ± 2 to 4 ± 2 ($p < 0.05$). Although improvements in attack frequency and severity were observed across all drug classes, Tricyclic antidepressants exhibited the most significant efficacy. 50% of antiepileptic-treated patients reported being affected post-treatment, with Tricyclic antidepressants influencing 28%.

Conclusion: After six-month evaluation of medications examined, Tricyclic antidepressants demonstrated the highest level of efficacy in management of common migraines. However, symptom relief was observed in all drug categories to varying degrees, emphasizing the importance of tailoring treatment plans to patient-specific characteristics.

Key Words: Antidepressants; Antiepileptics; Beta-Blockers; Calcium Channel Blockers.

Citation of article: Khan S, Ahmad I, Jalil M, Saleem I, Ahmad Fm, Nawaz KH. Comparison of Treatment Response of Different Drugs in Common Migraine. Med Forum 2024;35(1):58-62. doi:10.60110/medforum.350113.

INTRODUCTION

Migraine is a neurovascular disorder that causes significant global burden, characterized by recurrent episodes of severe cephalalgia that affect millions of people, majority of whom are women¹. These episodes are characterized by intense, pulsating pains that are typically localized to one side of cranium.

Department of Neurology, Pak Emirates Military Hospital, Rawalpindi.

Correspondence: Dr. Salman Khan, Trainee Neurology, Pak Emirates Military Hospital, Rawalpindi.

Contact No: 03339698830

Email: salman.imdc1@gmail.com

Received: July, 2023

Accepted: September, 2023

Printed: January, 2024

Nausea, vomiting and an acute sensitivity to light and sound are frequent accompanying symptoms². Numerous victims encounter auras, which manifest as visual impairments or blind areas, serving as indicators of impending assault. Migraines may persist for several hours to days, causing significant impairment to an individual's daily functioning. While the precise mechanism underlying migraines remains unknown, it is hypothesized that they are caused by secretion of inflammatory mediators in the vicinity of cranial nerves and blood vessels.

Migraines have consequences that extend beyond mere physical distress; they impose substantial socio-economic burden through the hindrance of work performance and personal relationships³. Migraine is often neglected and inadequately managed, resulting in significant number of individuals being compelled to endure the agony without any alleviation. Nevertheless, ongoing progress in migraine research is revealing more about the disorder's biological foundations and potential treatments, providing glimmer of hope for

individuals afflicted with this incapacitating condition and enhancing their quality of life⁴⁻⁵.

Significant advancements have been made in the treatment of migraines over the last few decades. Historically, treatment approaches for migraines have been divided into two categories: acute and preventive. Acute strategies seek to alleviate or terminate ongoing attacks, while preventive strategies strive to diminish the frequency and intensity of migraines⁶. Acute treatments comprise the variety of pharmaceutical interventions, including Topiramate, ergots, antiemetics and basic analgesics. In addition to calcium channel blockers, antiepileptic drugs, beta-blockers and antidepressants, preventive treatments have expanded to include monoclonal antibodies that target calcitonin gene-related peptide (CGRP) pathway⁷. Nevertheless, patient response to these interventions is extraordinarily variable and decision-making continues to be empirical, frequently predicated on trial and error⁸.

Through the assessment of response to various treatments, potential biomarkers of response can be identified, thereby augmenting the accuracy of migraine management. This is consistent with the rapidly expanding domain of pharmacogenomics, which utilizes the genetic underpinnings of drug response to customize therapeutic interventions⁹⁻¹⁰.

The objective of this study was to investigate the varying responses of common migraine patients to different medication treatments and to identify potential contributors to this variability, such as genetic predispositions, comorbid conditions, and phenotypic manifestation of the migraine.

METHODS

The research described herein was undertaken at Pak Emirates Military Hospital in Rawalpindi for the duration of one year, specifically from Jan 2022 to Jan 2023. The study population comprised 234 individuals who were diagnosed with common migraine as defined by ICHD-3¹¹. Age between 18 and 65 years, migraine diagnosis spanning at least one year, and minimum of one migraine episode per month during the previous three months constituted the inclusion criteria. Exclusion criteria for this study included patients who presented with chronic migraine, secondary headache disorders or significant comorbidities including cardiovascular disease, renal impairment or hepatic dysfunction.

Intervention Procedure: The medication regimen administered to patients was utilized to classify them into the following categories: Tricyclic antidepressants, Topiramate, beta-blockers, calcium channel blockers and antiepileptic pharmaceuticals. The treatment regimen was in accordance with standard clinical practice guidelines¹².

Data Collection: Clinical and baseline demographic information, including age, gender, migraine frequency,

duration and intensity was gathered. The evaluation of treatment response was conducted using MTRS, which was recorded at three, one, and six months after treatment commenced. Through clinical evaluations and patient self-reports, adverse effects were documented.

Outcome Measures: At each time point, primary outcome indicator was change in MTRS. The frequency of migraine attacks, alterations in the severity of attacks and occurrence of treatment-related adverse events constituted secondary outcomes.

Statistical Analysis: The data were analyzed with version 25.0 of SPSS. Descriptive statistics were utilized to provide the summary of sample's demographic and clinical attributes. Using repeated measures ANOVA for continuous variables and Chi-square test for categorical variables, treatment responses were compared. A p-value below 0.05 was deemed to indicate statistical significance.

Ethical Approval: Following evaluation and approval by Institutional evaluation Board of Pak Emirates Military Hospital, Rawalpindi, study protocol was implemented. All procedures conducted in this study adhered to the ethical guidelines set forth by the institutional research committee, Helsinki Declaration of 1964 and comparable standards of ethics.

RESULTS

The analysis comprised 234 patients who were diagnosed with common migraine. The average age of these patients was 40±12 years. The participants were predominantly composed of females (65.8%), with only slight variations observed among the treatment groups. The antidepressant cohort comprised participants having age of 43±15, whereas CGRP inhibitor group comprised patients of age 38±11 years. The mean number of migraine attacks per month was four, with CGRP inhibitor and beta-blocker groups experiencing fewer attacks (3±1). The mean duration of migraine history was 9±7 for antiepileptic users, while antidepressant users reported the minimum duration of 5 years. The migraine intensity exhibited mean value of 7±2. It is worth mentioning that antiepileptic groups documented reduced intensity 8±2 and 8±1, respectively, in contrast to the Tricyclic antidepressants group (Table 1).

Tricyclic antidepressants demonstrated consistent superior efficacy in treatment of migraines throughout the six-month study period. Response scores improved significantly from 4.5±1.2 at one month to 5.8±1.0 at six months (p<0.05). The mean response scores of Topiramate, beta-blockers, calcium channel blockers and antidepressants all increased moderately with time; however, none of these medications achieved the same level of effectiveness as Tricyclic antidepressants. With antiepileptics, least improvement was observed. The results of statistical analysis revealed that differential treatment responses became more significant as the

time points progressed; the p-values decreased from 0.045 at one month to 0.001 at six months, indicating the distinct trends (Table 2). Following treatment with Tricyclic antidepressants, there was notable decrease in frequency and severity of migraine attacks. Specifically, frequency of attacks decreased from 4±2 to 2±1 (p<0.05), while severity decreased from 7±2 to 4±2 (p<0.05). Topiramate also reduced assault frequency and severity by significant margins, from 4.2 to 3.1 (p<0.05) and 7.1 to 5.2 (p<0.05), respectively. Although beta-blockers did not have significant impact on attack frequency, they did marginally reduce severity from 7.3 to 6.3% (p<0.05). The frequency and severity of calcium channel blockers were reduced from 4.2 to 3.1 (p<0.05) and 7.2 to 5.0 (p<0.05),

respectively, bearing positive albeit less pronounced impact. Antiepileptics demonstrated efficacy by reducing the frequency by 5±3 to 4±2 (p<0.05) and severity by 8±1 to 6±2 (p<0.05). Antidepressants demonstrated a modest benefit by reducing severity from 6±3 to 5±2 (p<0.05) and frequency from 4±2 to 3±1 (p<0.05). Statistical improvements in both frequency and severity were observed across all pharmacological categories following treatment, with Tricyclic antidepressants demonstrating the greatest efficacy (Table 3). 50% who were prescribed antiepileptics. Topiramate influenced 38% of the patients, while beta-blockers influenced 45%. 30% of the patients were affected by Tricyclic antidepressants and Calcium Channel Blockers (Figure 1).

Table No. 1: Baseline demographic and clinical characteristics of participants

Variable	Total (N=234)	Tricyclic antidepressants	Topiramate	Beta-Blockers	Calcium Channel Blockers	Antiepileptics
Age (Mean±SD) years	40±12	38±11	39±10	41±14	40±12	37±9
Female n(%)	154 (65.8)	22 (68.8)	25 (64.1)	28 (70)	26 (65)	15 (62.5)
Migraine frequency/month (Mean±SD)	4±2	3±1	4±2	3±1	4±2	5±3
Migraine duration (Mean±SD) years	7 ±5	6±4	7±6	6±3	7±5	9±7
Migraine intensity (Mean±SD)	7±2	6±2	7±1	7±3	7±2	8±1

Table No. 2: Migraine Treatment Response Score (MTRS) at each time point

Time Point	Tricyclic antidepressants	Topiramate	Beta-Blockers	Calcium Channel Blockers	Antiepileptics	p-value
1 Month	4.5±1.2	4.0±1.3	3.5±1.2	3.8±1.4	2.5±1.3	0.045*
3 Months	5.2±1.1	4.5±1.1	4.0±1.3	4.2±1.2	3.0±1.4	0.012*
6 Months	5.8±1.0	5.0±1.2	4.5±1.1	4.7±1.1	3.5±1.5	0.001*

*indicated the significant values

Table No. 3: Frequency and severity of migraine attacks post-treatment

Drug Category	Baseline Attack Frequency	Post-Treatment Attack Frequency	p-value	Baseline Attack Severity	Post-Treatment Attack Severity	p-value
Tricyclic antidepressants	4±2	2±1	0.001*	7±2	4±2	0.001*
Topiramate	4±2	3±1	0.012*	7±1	5±2	0.018*
Beta-Blockers	3±1	3±1	0.020*	7±3	6±3	0.053
Calcium Channel Blockers	4±2	3±1	0.045*	7±2	5±2	0.051
Antiepileptics	5±3	4±2	0.031*	8±1	6±2	0.045*

*indicated the significant values




Figure No. 1: Incidence of Treatment-Related Adverse Events.

DISCUSSION

The current investigation examined the relative efficacy of different medications in managing patients with common migraines for a duration of six months. A significant finding was the notable effectiveness of Tricyclic antidepressants, especially when evaluated for extended periods of time, which suggests they have the potential to be a formidable tool for managing migraines¹³.

The study's female participants comprised the majority (65.8%), which is consistent with previous research indicating that migraines are more prevalent among women than men¹⁴. The age discrepancies observed among various drug categories, specifically in the CGRP inhibitor group in comparison to the antidepressant cohort, may be attributed to pharmacokinetic and adverse effect profiles of the drugs, which may have an impact on prescribing practices¹⁵.

Tricyclic antidepressants exhibited the most substantial decrease in both the frequency and severity of migraine attacks; nevertheless and Topiramate also demonstrated noteworthy effectiveness. The reason for this is their firmly established functions in the treatment of acute migraines⁷. Further corroboration of prior research underscores the importance of extended treatment durations for thorough evaluation, as evidenced by the upward trend in response scores for the majority of pharmaceuticals over time¹⁶.

Consistent with previous research¹⁷, beta-blockers substantially diminished the severity of attacks while having only the marginal effect on their frequency. Although noteworthy, advantages associated with calcium channel blockers, antiepileptics, and antidepressants were comparatively subdued. The risk-benefit ratio of these treatments should be diligently assessed, particularly when prescribing for extended periods of time, in light of these findings¹⁷.

The observed reduction in migraine intensity among the antiepileptic groups, as opposed to the Tricyclic antidepressants and antidepressants, could potentially be attributed to the distinct mechanisms of action exhibited by these pharmaceuticals. In contrast, antiepileptics alter neurotransmitter release by

modulating voltage-gated sodium and calcium channels¹⁸.

Additionally, improvements in both frequency and severity were observed across all pharmacological categories following treatment, according to the study. This consistency indicates that although certain drug categories may provide more substantial benefits, all drug categories offer some degree of alleviation. In contrast, the fact that Tricyclic antidepressants exhibited the greatest efficacy highlights their increasing significance in the treatment of migraines. The relatively low proportions of Tricyclic antidepressants in the sample may be attributed to their recent introduction to the market or possible financial obstacles¹⁹. Conducting additional randomized controlled trials would be advantageous in solidifying these findings.

CONCLUSION

In contrast to other drug classes, Tricyclic antidepressants significantly diminished the frequency and severity of migraine attacks. Nevertheless, despite the potential of Tricyclic antidepressants as primary therapeutic intervention, each drug class exhibited varying levels of alleviation, underscoring the significance of individualized treatment strategies. In order to customize therapeutic interventions for migraines, patient-specific factors, potential medication interactions, and adverse effects must be considered owing to the wide array of migraine manifestations and triggers.

Author's Contribution:

Concept & Design of Study: Salman Khan
 Drafting: Imran Ahmad, Muhammad Jalil
 Data Analysis: Ibrar Saleem, Fawad Ahmad, Khuram Haq Nawaz
 Revisiting Critically: Salman Khan, Imran Ahmad
 Final Approval of version: Salman Khan

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.478/2021 dated 05.09.2021

REFERENCES

- Amiri P, Kazeminasab S, Nejadghaderi SA, Mohammadinasab R, Pourfathi H, Araj-Khodaei M, et al. Migraine: A Review on Its History, Global Epidemiology, Risk Factors, and Comorbidities. *Front Neurol* 2022;12:800605.
- Khan J, Al Asoom LI, Al Sunni A, Rafique N, Latif R, Al Saif S, et al. Genetics, pathophysiology,

- diagnosis, treatment, management, and prevention of migraine. *Biomed Pharmacotherapy* 2021; 139:111557.
3. Burch R, Rizzoli P, Loder E. The prevalence and impact of migraine and severe headache in the United States: Updated age, sex, and socioeconomic-specific estimates from government health surveys. *Headache* 2021;61(1):60-68.
 4. Andersson M, Persson M, Kjellgren A. Psychoactive substances as a last resort-a qualitative study of self-treatment of migraine and cluster headaches. *Harm Reduct J* 2017;14(1):60.
 5. Scaini G, Valvassori SS, Diaz AP, Lima CN, Benevenuto D, Fries GR, Quevedo J. Neurobiology of bipolar disorders: a review of genetic components, signaling pathways, biochemical changes, and neuroimaging findings. *Braz J Psychiatr* 2020;42(5):536-551.
 6. Antonaci F, Ghiotto N, Wu S, Pucci E, Costa A. Recent advances in migraine therapy. *Springerplus* 2016;5:637.
 7. Gawde P, Shah H, Patel H, Bharathi KS, Patel N, Sethi Y, et al. Revisiting Migraine: The Evolving Pathophysiology and the Expanding Management Armamentarium. *Cureus* 2023;15(2):e34553.
 8. Molina-Mula J, Gallo-Estrada J. Impact of Nurse-Patient Relationship on Quality of Care and Patient Autonomy in Decision-Making. *Int J Environ Res Public Health* 2020;17(3):835.
 9. D'Amico D, Tepper SJ. Prophylaxis of migraine: general principles and patient acceptance. *Neuropsychiatr Dis Treat* 2008;4(6):1155-67.
 10. Ong JJY, De Felice M. Migraine Treatment: Current Acute Medications and Their Potential Mechanisms of Action. *Neurotherapeutics* 2018;15(2):274-290.
 11. Kim BK, Cho SJ, Kim BS, Sohn JH, Kim SK, Cha MJ, et al. Comprehensive Application of the International Classification of Headache Disorders Third Edition, Beta Version. *J Korean Med Sci* 2016;31(1):106-13.
 12. Zobdeh F, ben Kraiem A, Attwood MM, Chubarev VN, Tarasov VV, Schiöth HB, et al. Pharmacological treatment of migraine: Drug classes, mechanisms of action, clinical trials and new treatments. *Br J Pharmacol* 2021;178(23): 4588-4607.
 13. Masoud AT, Hasan MT, Sayed A, Edward HN, Amer AM, Naga AE, et al. Efficacy of calcitonin gene-related peptide (CGRP) receptor blockers in reducing the number of monthly migraine headache days (MHDs): A network meta-analysis of randomized controlled trials. *J Neurol Sci* 2021;427:117505.
 14. Babateen O, Althobaiti FS, Alhazmi MA, Al-Ghamdi E, Alharbi F, Moffareh AK, et al. Association of Migraine Headache With Depression, Anxiety, and Stress in the Population of Makkah City, Saudi Arabia: A Cross-Sectional Study. *Cureus* 2023;15(5):e39788.
 15. Boyce RD, Handler SM, Karp JF, Hanlon JT. Age-related changes in antidepressant pharmacokinetics and potential drug-drug interactions: a comparison of evidence-based literature and package insert information. *Am J Geriatr Pharmacother* 2012; 10(2):139-50.
 16. Anghel LA, Farcas AM, Oprean RN. An overview of the common methods used to measure treatment adherence. *Med Pharm Rep* 2019;92(2):117-122.
 17. Rothwell PM, Howard SC, Dolan E, O'Brien E, Dobson JE, Dahlöf B, et al. ASCOT-BPLA and MRC Trial Investigators. Effects of beta blockers and calcium-channel blockers on within-individual variability in blood pressure and risk of stroke. *Lancet Neurol* 2010;9(5):469-80.
 18. Russo M, De Rosa MA, Calisi D, Consoli S, Evangelista G, Dono F, et al. Migraine Pharmacological Treatment and Cognitive Impairment: Risks and Benefits. *Int J Mol Sci* 2022;23(19):11418.
 19. Russell FA, King R, Smillie SJ, Kodji X, Brain SD. Calcitonin gene-related peptide: physiology and pathophysiology. *Physiol Rev* 2014;94(4): 1099-142.

Effect of Informed Consent on Patient's Anxiety Regarding Third Molar Surgery

Kiran Bashir¹, Sadia Paiker², Erum Riaz¹, Wajiha Walayat³, Syed Muhammad Zaki Mehdi⁴ and Fatima Khattak¹

ABSTRACT

Objective: To assess mean change in anxiety of a patient undergoing 3rd molar surgery before and after disclosure of information regarding procedure.

Study Design: A descriptive study

Place and Duration of Study: This study was conducted on 50 patients that were presented to the outpatient department of a tertiary care hospital in FUCD between December 2018 to November 2019.

Methods: Patients reporting to OMFS department of FUCD for surgical removal of impacted 3rd molars were selected after history, clinical examination & radiographic evaluation. Informed consent form was given to the patients starting by explaining the brief overview of procedure, benefits and racial complications. Patients were asked to sign the informed consent form. Anxiety of patient after taking informed consent is evaluated by STAI form. The level of anxiety by using STAI on first visit and later again using the STAI on the day of procedure was compared.

Results: A total of 50 patients where 13(26.0) males and 37 (74.0) female patients having mean age (years) 29.00 ± 6.96 were included in the study. Our study finding showed that mean change in anxiety of patient undergoing 3rd molar surgery and after disclosure of information regarding procedure was 35.06 ± 2.25 and 45.60 ± 2.50 respectively with mean change 10.53 ± 0.25 .

Conclusion: The study concluded that patient experienced anxiety when underwent molar surgery. So there is a need of patient counseling about the procedural risks and associated complications regarding 3rd molar surgery in a sequential manner.

Key Words: Surgical extraction, Anxiety, Impaction

Citation of article: Bashir K, Paiker S, Riaz E, Walayat W, Mehdi SMZ, Khattak F. Effect of Informed Consent on Patient's Anxiety Regarding Third Molar Surgery. Med Forum 2024;35(1):63-67. doi:10.60110/medforum.350114.

INTRODUCTION

The significance of a patient-doctor relationship may be highlighted by educating patients about the processes that must be done during any surgery. This enlightenment occurs prior to the actual process. To correct some inconsistencies and malpractices observed

on the part of medical professionals throughout the years, it became important to keep patients aware of the whole process prior to medical operations, as well as gain patients agreement, which all indicate patients being involved in their own medical care.¹

An informed consent is the process of engaging in a dialogue between the patient and healthcare practitioner about a proposed medical treatment, including the nature of the treatment, its potential benefits, harms, and risks, as well as alternative healthcare services to a patient after which they have granted permission, and it includes three critical components: voluntarism, information disclosure, and decision-making capacity.²

Patients must have a thorough understanding of the necessary steps involved in their treatment before the actual procedure. This is crucial both to prevent any discrepancies or malpractices by medical professionals and to involve patients in their own medical care. It provides them with a sense of control, increased confidence, and a feeling of cooperation in the surgical process. Ultimately, informed consent strengthens the bond between patient and doctor, promotes a more positive surgical experience, and improves treatment outcomes.³

¹. Department of Oral & Maxillofacial Surgery, Foundation university dental college Islamabad.

². Department of Oral & Maxillofacial Surgery, Islamabad Medical & Dental College SZABMU, Islamabad.

³. Department of Dental Surgery, Pakistan Institute of Medical Sciences, SZABMU, Islamabad.

⁴. Department of Oral & Maxillofacial Surgery, Rawal Institute of Health Sciences SZABMU, Islamabad.

Correspondence: Dr. Sadia Paiker, Senior Registrar Department of Oral & Maxillofacial Surgery, Islamabad Medical & Dental College SZABMU, Islamabad.

Contact No: 0308-5801680

Email: sadia_paiker@hotmail.com

Received: September, 2023

Accepted: November, 2023

Printed: January, 2024

Studies have revealed that patient anxiety levels can be significantly reduced when they are provided with thorough preoperative information. This is essential as anxiety is directly related to the perception and tolerance of pain. Elevated anxiety levels may also impair how well the practitioner performs delicate and complex treatment procedures. Dentists, in particular, prioritize patient comfort and stress reduction, which greatly contributes to the technical superiority of the treatment.⁴

For instance, anxiety is an undesirable psychological experience that provokes tension, sweating, and increased pulse rate. Before any surgical intervention, patients may experience anxiety which affects their overall wellbeing. Hence, it is critical to obtain informed consent to access the anxiety level of a patient before surgery. The surgical extraction of third molars is the most common minor oral surgical procedure performed worldwide; making it a useful model for studying informed consent.⁵

Patients undergoing the surgery experience anxiety and it is essential to provide them with relevant information such as the procedure, its benefits, risks, complications, and alternatives.

However, communicating intricate details about surgical procedures and their potential complications can prove problematic. It can often be challenging to relay complex medical information to patients, which can have negative consequences.⁶ One study found that patients became more anxious when presented with step-by-step information before the surgical removal of impacted third molars. This information typically details the potential unfavorable outcomes of the surgical procedure, such as temporary or permanent nerve damage, infection, dry socket, hemorrhage, trismus, mandible fractures, adjacent tooth damage, and pain. Disclosing this information to patients before surgery can be distressing, thereby increasing anxiety levels^{7,8}.

Therefore, the purpose of this study is to investigate the level of anxiety patients experience before and after being counseled about the procedural risks, associated complications, and alternative management strategies regarding third molar surgery in a sequential manner. This study will help improve the quality of healthcare services by providing patients with adequate information about the procedures and their potential consequences.

METHODS

Patients between the age range of 18-45 years, requiring at least one lower third molar that showed symptoms of mild to severe inflammation or decay and the absence of a prior history of third molar surgery will be considered for inclusion. Whereas patients having systemic diseases or compromised immune systems, pregnant females and those who refuse to participate in

the study were excluded. The research has received ethical approval from the hospital's Ethical Committee. At their first appointment, patients were asked to evaluate their anxiety by completing the Spielberger's State-Trait Anxiety Inventory (STAI) form. The STAI-S is a 20-item self-evaluation questionnaire that analyses transitory emotional states or situations as characterized by subjective emotions of tension and anxiety that can fluctuate in duration and intensity. It is scored using a 4-level frequency scale ranging from 1 to 4 (1 calm, 2 somewhat anxious, 3 fairly anxious, 4 extremely anxious). Informed consent form was given to the patients starting by explaining the brief overview of procedure, benefits and racial complications. Then patient was also informed about following options whether to postpone the extraction or to undergo extraction of 3rd molar. Once the patient was confirmed that they understood the procedure. They were asked to sign the informed consent form. All patients were asked to evaluate their anxiety prior to surgery.

The minimum required sample size ($n=50$) was calculated with by Open Epi collections of epidemiologic calculators, considering a 95% level of confidence, 5% alpha error, 90% study power, Population mean after informed consent=42.46, Population SD after informed consent=7.076.

Results were analyzed using SPSS version 20.0 for quantitative variable like age whereas qualitative variables like gender, education and socioeconomic status (SES) was measured as frequency and percentage. Paired sample t-test was applied to compare pre and post STAI score. Effect modifier like age, gender, SES, education level was controlled by stratification. Post stratification paired sample t-test was applied. P-value <0.05 was considered significant.

RESULTS

Out of 50 patients, 13 (26%) were males and 37 (74%) were females as shown in Fig No 01. Mean age (years) of the patients in this study was 29.00 ± 6.96 . Frequency and percentage of education level was assessed in the study in terms of education (primary, middle, matric, and graduation, post-graduation) and non-education. Majority of the patients have education background, following by 12 (24%) primary education, 10 (20%) middle education whereas there were 11 (22%) patients had no education background. Socio economic status was in the study in terms of frequency and percentage of low ($<20,000$ PKR), middle (20,000-50,000) and upper level ($>50,000$) of socio economic status. Majority of the patients 37 (74.0) belonged from low income status and 12 (24.0) patients belong to middle income status (20,000-50,000), as shown in Table 1.

The objective of the study was to assess mean change in anxiety of patient undergoing 3rd molar surgery and after disclosure of information regarding procedure. Our study finding showed that mean change in anxiety

of patient undergoing 3rd molar surgery and after disclosure of information regarding procedure was 35.06 ± 2.25 and 45.60 ± 2.50 respectively with mean change 10.53 ± 0.25 , as shown in Table. No.02.

Table No.1: Descriptive statistics of Demographic data.

Variable		Frequency	Percentage %	Mean age 29 ± 6.96
Gender	Male	13	26.0%	
	Female	37	74.0%	
	Total	50	100%	
Education level	Uneducated	11	22.0%	
	Primary	12	24.0%	
	Middle	10	20.0%	
	Matric	8	16.0%	
	Graduated	8	16.0%	
	Post graduate	1	2.0%	
	Total	50	100%	
Socio-Economic status	Lower class	37	74.0%	
	Middle class	12	24.0%	
	Upper class	1	2.0%	
	Total	50	100%	

Effect modifier like age was stratified and compared with mean change in anxiety of patient undergoing 3rd molar surgery and after disclosure of information regarding procedure. Among patient with age 31 – 45 years, mean change in anxiety of patient undergoing 3rd molar surgery and after disclosure of information regarding procedure was 35.09 ± 1.9 and 45.44 ± 2.13 respectively with mean change 10.35 ± 2.11 . Effect modifier like gender was stratified and compared with mean change in anxiety of patient undergoing 3rd molar surgery and after disclosure of information regarding 3rd molar surgery and after disclosure of information regarding procedure was 35.03 ± 2.59 and 45.78 ± 2.92 respectively with mean change 10.74 ± 2.14 .

Effect modifier like education level was stratified and compared with mean change in anxiety of patient undergoing 3rd molar surgery and after disclosure of information regarding procedure. Among patients who have primary education background, mean change in anxiety of patient undergoing 3rd molar surgery and after disclosure of information regarding procedure was 35.10 ± 2.01 and 45.16 ± 2.82 respectively with mean change 10.35 ± 2.04 , as shown in Table. No.3.




Figure No. 1: Pie Chart of Gender Distribution.

Table No.2: Descriptive statistics of STAI-S scale (before and after).

	N	Min.	Max.	Mean	Std. Deviation
STAI scale (before)	50	30.20	40.20	35.06	2.25
STAI scale (after)	50	38.00	49.00	45.60	2.50
Anxiety (STAI scale score)	before	after	mean change	p-value	
	35.06 ± 2.25	45.60 ± 2.50	10.53 ± 0.25	0.000	

Table No.3: Demographic data stratification with comparison of mean change in anxiety of a patient undergoing 3rd molar surgery before and after disclosure of information regarding procedure.

Variable		Anxiety (STAI scale score)			p-value
		Before	After	Mean change	
Age	18-30years	35.09 ± 1.97	45.44 ± 2.13	10.35 ± 2.11	0.000
	31-45years	35.03 ± 2.59	45.78 ± 2.92	10.74 ± 2.14	0.000
Gender	Male	35.03 ± 2.59	45.78 ± 2.92	10.74 ± 2.14	0.000
	Female	35.10 ± 2.25	45.51 ± 2.39	10.41 ± 2.07	0.000
Education level	Uneducated	35.48 ± 2.56	45.00 ± 3.16	9.51 ± 2.56	0.000
	Primary	35.10 ± 2.01	45.16 ± 2.82	10.05 ± 2.04	0.000
	Middle	34.17 ± 2.70	45.70 ± 1.88	11.53 ± 2.31	0.000
	Matric	35.10 ± 1.47	4.37 ± 1.40	11.27 ± 1.08	0.000
	Graduation	35.91 ± 2.17	46.75 ± 2.12	10.83 ± 1.64	0.000
	Post-graduation	35.82 ± 2.02	45.21 ± 2.30	9.39 ± 0.28	0.000
	Socio-Economic status	Low	35.24 ± 2.15	45.86 ± 2.28	10.61 ± 1.84
Middle	34.33 ± 2.53	44.66 ± 3.08	10.33 ± 2.94	0.000	
Upper	34.29 ± 2.24	44.59 ± 3.12	10.30 ± 0.88	0.000	

DISCUSSION

There is a belief that providing patients with detailed information about their treatment could exacerbate their anxiety and lead to a reluctance to undergo the procedure. Conversely, studies have shown that doctor-patient communication fosters a collaborative relationship and can help alleviate anxiety. However, it is important to note that this communication can have a reciprocal effect, where the doctor can be influenced by the patient's anxiety and vice versa. Thus, the purpose of this study was to assess how an informed consent protocol impacts patient anxiety levels.⁹

Informed consent is crucial for clinical practice. It is still primarily a legal and ethical concept. The key consideration in any informed consent is its substance. By content, we mean the informed consent's material, which must always transmit two sorts of information. The first type is the therapy or operation that the patient will get. In other words, the patient is educated about the various stages of the procedure: pre, intra, and postoperative. The second piece of information is on the feelings that the patient is likely to have: pain, somnolence, stiffness, and so on. The importance of risks disclosure in the informed consent process cannot be overstated, as it is necessary to highlight potential adverse effects and avoid medical malpractice litigation cases.¹⁰

It has been shown that the efficacy of informative procedures is heavily influenced by patients' attitudes. The information provided has been shown to have positive effects on those who try to overcome stressful situations by gathering as much information as possible about them, but it may have negative effects on "avoidant" patients (those who reject all information in order to overcome anxiety by not thinking about the problem).¹¹

Patients appear to be more interested in information related to benefits and post-procedure complications than risks, which are more important from an ethical standpoint. The study's findings showed a mean change in anxiety levels of 35.06 ± 2.25 and 45.60 ± 2.50 before and after disclosing information regarding the procedure. Casap et al.¹² the influence of informed consent on stress levels linked with the removal of impacted mandibular third molars was evaluated. They discovered that presenting excessively thorough lists and disclosures prior to excision of impacted mandibular third teeth might enhance patient anxiety. Another study reported a mean change in anxiety levels of 36.6 and 42.4 before and after information disclosure regarding the procedure.¹³⁻¹⁵

Overall, the informed consent process is a critical component of any surgical procedure. However, it is essential to recognize the potential anxiety-inducing nature of this process and to take steps to alleviate patient anxiety in the hours and days leading up to the procedure. Vigilant patients who seek out information benefit more from the informed consent process. The

timing of information disclosure may not be as critical as once thought and patients appear to be more interested in benefits-related information than risks.¹⁴⁻²⁰

This study has several limitations. First, the sample was selected from a single local hospital. The patients included in this research were relatively young, which may have caused a selection bias. The representativeness of the sample might be restricted, and our results may have poor generalizability. Secondly, the exposure and outcome variables were collected through self-completed questionnaires, which may not accurately reflect the situation. For sample selection, it would have been ideal to conduct the study in a multicenter setting. Finally, no postoperative pain evaluation technique was intended for this study, and grading surgical difficulties in connection to anxiety level was not addressed.

CONCLUSION

The study concluded that patient experienced anxiety when underwent molar surgery. So there is a need of patient counseling about the procedural risks and associated complications regarding 3rd molar surgery in a sequential manner. Indeed, the informed consent form itself was a major contributor to elevate patient anxiety. The presentation of detailed information and nonsurgical treatment options might dissuade patients from undergoing extractions.

Acknowledgement: we are thankful to Dr. Mor Khan for his contribution in drafting of this study for the final publication. We are also thankful to our juniors at the Oral and Maxillofacial Department of FUCD for helping in data collection.

Author's Contribution:

Concept & Design of Study:	Kiran Bashir, Sadia Paiker
Drafting:	Erum Riaz, Wajiha Walayat
Data Analysis:	Syed Muhammad Zaki Mehdi, Fatima Khattak
Revisiting Critically:	Kiran Bashir, Sadia Paiker
Final Approval of version:	Kiran Bashir, Sadia Paiker

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.FF/FUMC/215-371-PHY/18 dated 13.11.2018.

REFERENCES

1. Sancak KT, Akal ÜK. Effect of verbal and written information and previous surgical experience on anxiety during third molar extraction. JOMS

- 2019;77(9):1769-e1.
2. Saincher RR, Pentapati KC, Gadicherla S. Effect of audio-visual treatment information on hemodynamic parameters during the transalveolar extraction of mandibular third molars: A randomized clinical trial. *J Int Society Preventive Comm Dentistry* 2019;9(1):21.
 3. Lakshman K, Mamatha H, Rachana N, Sumitha C, Ranganath N, Gowda V, et al. Effectiveness of Preoperative Multimedia Video-Based Education on Anxiety and Haemodynamic Stability of Oncosurgery Patients Undergoing Spinal Anaesthesia-A Randomised Controlled Trial. *J Clin Diagnostic Res* 2022;16(1).
 4. Navarro-Gastón D, Munuera-Martínez PV. Prevalence of preoperative anxiety and its relationship with postoperative pain in foot nail surgery: a cross-sectional study. *Int J Environmental Research Public Health* 2020; 17(12):4481.
 5. Laskin DM, Priest JH, Alfaqih S, Carrico CK. Does viewing a third molar informed consent video decrease patients' anxiety? *JOMS* 2018;76(12): 2515-7.
 6. Erbasar GN, Sancak KT. Should Preoperative Information Before Impacted Third Molar Extraction Be Visual, Verbal, or Both? *JOMS*. 2023;81(5):632-40.
 7. Schoenfeld EM, Kanzaria HK, Quigley DD, Marie PS, Nayyar N, Sabbagh SH, et al. Patient preferences regarding shared decision making in the emergency department: findings from a multisite survey. *Academic Emerg Med* 2018; 25(10):1118-28.
 8. Qiao F, Zhang M, Zhang T, Zhu D. Dental anxiety is related to postoperative symptoms in third molar surgery. *Frontiers Psychiatr* 2022;13:956566.
 9. GocmenG, Atali O, Gonul O, Goker K. Impact of informed consent on patient's decision regarding 3rd molar surgery. *Nigerian Int J Clin Pract* 2017; 158-162.
 10. Omezli MM, Torul D, Kahveci K. Does watching videos increase the perioperative anxiety in patients undergoing third molar surgery? A randomized trial. *JOMS* 2020;78(2):216-e1.
 11. Lagares D, Meseguer M, Velazquez F, Iglesias P, Hernandez G, Pacheco E, et al. The effects of informed consent Format on preoperative anxiety in patients undergoing inferior third molar surgery. *Med Oral Cir Bucal* 2014;19(3):e270-273.
 12. Wong NS, Yeung AW, Li KY, McGrath CP, Leung YY. Non-pharmacological interventions for reducing fear and anxiety in patients undergoing third molar extraction under local anesthesia: Systematic review and meta-analysis. *Int J Environmental Research Public Health* 2022; 19(18):11162.
 13. Torres-LagaresD, Heras-MeseguerM, Azcárate-Velázquez F, Hita-Iglesias P, Ruiz-de-León-Hernández G, et al. The effects of informed consent format on preoperative anxiety in patients undergoing inferior third molar surgery. *Med Oral Patol Oral Cir Bucal* 2014;19:e270-3.
 14. Moaddabi A, Hasheminia D, Bagheri S, Soltani P, Patini R. Effect of opaque eye coverage on anxiety in candidates for surgical removal of impacted third molars: a randomized clinical trial *Oral Surg. Oral Med. Oral Radiol* 2021;132(3):267-72.
 15. Gheisari R, Resalati F, Mahmoudi S, Golkari A, Mosaddad SA. Do different modes of delivering postoperative instructions to patients help reduce the side effects of tooth extraction? A randomized clinical trial. *J Oral Maxillofacial Surg* 2018; 76(8):1652-e1.
 16. Rajala M, Kaakinen P, Fordell M, Kääriäinen M. The quality of patient education in day surgery by adult patients. *J Peri Anesthesia Nursing* 2018; 33(2):177-87.
 17. Myoken Y, Kawamoto T, Fujita Y, Tsubahara Y, Toratani S, Yanamoto S, et al. Use of an educational video with mobile technology for the informed consent process: A randomised controlled trial on patients undergoing surgery for an impacted lower third molar. *Br J Oral Maxillofacial Surg* 2023;61(7):497-502.
 18. Etim S, Nzomiwu C, Eigbobo J. The practice of obtaining consents for dental care among dental practitioners in Nigeria. *Afr J Med Med Sci* 2020;49:61-5.
 19. Göçmen GÖ, Atalı O, Gonul O, Goker K. Impact of informed consent on patient decisions regarding third molar removal. *Nigerian J Clin Practice* 2017;20(2):158-62.
 20. Britton D, Burns A, Hudyba S, Nugent M. Third molar consent forms: how comprehensive should they be? *Br J Oral Maxillofacial Surg* 2016; 54(7):788-90.

The Influence of Surgical Volume on Outcomes in Radical Cystectomy: A Population-Based Analysis

Shahjehan¹, Syed Ameroon Shah², Maham Arooj Zaidi³, Khalid Islam², Muhammad Zohair² and Naseeb Dad²

ABSTRACT

Objective: The purpose of this research was to look at the effect of surgical volume on postoperative outcomes in patients receiving radical cystectomy for bladder cancer in a population-based cohort of 113 patients.

Study Design: A prospective analysis design study

Place and Duration of Study: This study was conducted at the Department of Urology IKD, Peshawar between August 2021 to August 2023.

Methods: Radical cystectomy was performed on 113 patients between August 2021 and August 2023. Data on demographics, comorbidities, surgery volume, and outcomes was collected. Based on operations conducted by each surgical practitioner, surgical volume was low, midrange, or high. Postoperative complications, hospital stay, and 30-day mortality were key outcomes.

Results: The study consisted 113 patients: 23 low volume, 34 middle volume, and 56 high volume. All participants had a mean age of 65.2 ± 7.1 years. The majority of patients (52.17%) were female. 82.61% had radical cystectomy, 17.39% partial. The majority of procedures were open (100%). Most patients reported infection (5.31%), followed by hemorrhage (3.54%) and serious problems (7.08%). Surgical volume did not significantly affect complications or LOS >7 days (adjusted OR 0.75, 95% CI 0.53-1.06, $p=0.105$). These data imply that surgical volume may affect significant complications but not overall complications, LOS, or mortality.

Conclusion: Our study revealed a link between surgical treatment volume and significant side effects in cystectomy patients.

Key Words: Radical Cystectomy, Surgical Volume, Bladder Cancer, Population-Based Analysis, Postoperative

Citation of article: Shahjehan, Shah SA, Zaidi MA, Islam K, Zohair M, Dad N. The Influence of Surgical Volume on Outcomes in Radical Cystectomy: A Population-Based Analysis. Med Forum 2024;35(1):68-72. doi:10.60110/medforum.350115.

INTRODUCTION

Bladder cancer is a major worldwide health issue, and radical cystectomy is a crucial aspect of its treatment^[1]. The results of radical cystectomy, a highly intricate surgical technique, may be affected by several variables, such as the number of surgeries conducted by the surgical care providers^[2,3]. Extensive study has been conducted on the correlation between the number of surgeries performed and the results after surgery.

¹. Department of Urology, MMCGRH, Peshawar.

². Department of Urology, Institute of Kidney Diseases, Peshawar.

³. Department of Surgery, Khyber Teaching Hospital, Peshawar.

Correspondence: Syed Ameroon Shah, Specialist Registrar, Institute of Kidney Diseases, Peshawar.

Contact No: 03339712477

Email: shahameroon@gmail.com

Received: October, 2023

Accepted: November, 2023

Printed: January, 2024

Evidence indicates that a greater number of surgeries may be linked to better patient outcomes^[4]. This study, based on a population, seeks to add to the current knowledge by investigating how the number of surgeries performed affects the outcomes after radical cystectomy in a group of 113 patients.

Bladder cancer poses a significant challenge to healthcare systems globally, requiring efficient solutions to improve surgical treatment and better patient outcomes^[5,6]. The concept of surgical volume, which refers to the quantity of operations carried out by a surgical care provider, has been identified as a possible factor influencing the quality of different surgical interventions^[7]. Previous research has shown a positive association between increased surgical volumes and enhanced outcomes in operations such as coronary artery bypass grafting and pancreatotomy^[8]. However, the influence of the number of surgeries performed on the results of radical cystectomy in the setting of bladder cancer is still a subject that needs further research.

This study used a population-based methodology, using a heterogeneous sample of 113 individuals who had radical cystectomy within a certain era and healthcare system. The aim is to evaluate whether there is a

correlation between the number of cystectomy surgeries conducted by surgical care providers and differences in postoperative complications, duration of hospitalization, and mortality within 30 days. Our objective is to analyze surgery volume and classify it as low, the middle, or high in order to uncover any discernible trends that may be used to shape healthcare policy and clinical decision-making.

Gaining insight into the correlation between the number of surgeries performed and the resulting results in radical cystectomy is essential for maximizing patient care and efficiently allocating resources. The results of this investigation might have significant ramifications for healthcare professionals, governments, and patients, highlighting the potential advantages of centralizing cystectomy services. The ultimate objective is to provide significant insights that may improve the quality of treatment for persons who are having radical cystectomy for bladder cancer.

METHODS

The study included a total of 113 patients. Data about demographic factors, coexisting medical conditions, and clinical attributes were gathered. Patient identifiers were anonymized to guarantee confidentiality. The surgical volume was classified according to the quantity of radical cystectomy operations conducted by each surgical professional. Centers were categorized as low, middle, or high volume based on specified criteria identified by the distribution of cases in the dataset. The primary variables assessed were surgical complications, duration of hospitalization, and mortality within 30 days. The postoperative complications were classified based on predefined criteria and their severity was assessed.

Statistical Analysis: Patient demographics and clinical features were summarized using descriptive statistics. The correlation between the number of surgeries performed and the resulting results was evaluated using suitable statistical procedures, such as chi-square for categorical variables and analysis of variance (ANOVA) for continuous data. A multivariate analysis was performed to account for any confounding factors.

Ethical Considerations: The work complied with ethical criteria and obtained permission by the IKD, Peshawar Review Board, and Ethics Committee. Strict adherence to patient confidentiality was ensured, and data were managed in compliance with applicable privacy legislation.

RESULTS

The study had 113 patients in total, distributed as follows: 23 in the low volume group, 34 in the intermediate volume group, and 56 in the high volume group. The average age of all participants was 65.2 ± 7.1 years. Females constituted the majority of patients, at 52.17% of the overall patient population. Regarding

comorbidities, hypertension was the prevailing disease, reported by 26.09% of patients. Among the patients, 21.74% reported having diabetes, making it the second most common comorbidity. Upon comparing the three volume groups, no substantial disparities were seen in terms of age or gender distribution. However, there were discrepancies in the occurrence of comorbidities, as a greater percentage of patients in the low volume category disclosed hypertension and diabetes in comparison to the intermediate and high volume categories. The research sample exhibited diversity in age, gender, and comorbidities, making it a representative sample for the study (Table-1).

The surgical characteristics of study participants are shown in Table 2. 17.39% of the 113 individuals had partial cystectomy, whereas 82.61% had radical cystectomy. The majority of procedures were open (100%). All subjects had a mean operational time of 240 minutes and a standard variation of 30 minutes. Cystectomy type and surgical method did not vary across the three volume groups. The high volume group had a mean operating time of 200 minutes, compared to 220 minutes in the intermediate volume group and 240 minutes in the low volume group. These surgical features reveal what was done on study participants and may affect their results.

The study participants' postoperative complications are shown in Table 3. Out of 113 patients, 29.20% had post-surgery complications. Most individuals reported infection (5.31%), followed by hemorrhage (3.54%) and serious problems (7.08%). Complication rates were similar among the three volume groups. The low volume group (13.04%) had more serious problems than the intermediate (8.82%) and high volume (3.57%) groups. These data imply that all volume groups have postoperative difficulties, although low volume centers may have a greater risk of significant complications.

Table 4 shows the study participants' LOS and mortality rates. All individuals' mean LOS was 6.5 days, with a 2.0-day standard deviation. The mean LOS was similar among the three volume groups. However, low volume (7.0 days) had a tendency towards longer LOS than intermediate (6.5 days) and high volume (6.0 days). A larger percentage of low volume participants (34.78%) experienced a LOS of more than 7 days than intermediate (11.76%) and high volume (7.14%) groups. The 30-day mortality rate was 3.54% for all individuals, with the low volume group (8.70%) having a greater risk than the intermediate (2.94%) and high volume (1.79%) groups. These data show that low-volume centers may have longer hospital stays and greater fatality rates than intermediate and high-volume centers.

Table 5 shows multivariate surgery volume and outcome statistics. After controlling for possible confounding variables, surgical volume did not predict

any complication (OR 0.75, 95% CI 0.53-1.06, p=0.105) or LOS >7 days (OR 1.32, 95% CI 0.88-1.97, p=0.186). However, surgical volume was associated with severe complications (adjusted OR 0.61, 95% CI 0.37-1.00, p=0.048), with intermediate and high volume centers having lower chances than low volume centers.

Surgical volume did not affect 30-day mortality (adjusted OR 0.92, 95% CI 0.47-1.78, p=0.796). These data imply that surgery volume may affect significant complications but not overall complications, LOS, or mortality.

Table No. 1: Demographic Characteristics of Study Participants

Demographic Factor	Total (n=113)	Low Volume (n=23)	Intermediate Volume (n=34)	High Volume (n=56)
Age (years)	Mean (SD)	65.2±7.1	64.8±6.5	66.4±7.8
Gender	Female	12 (52.17%)	19 (55.88%)	26 (46.43%)
	Male	11 (47.83%)	15 (44.12%)	30 (54.56%)
Comorbidities	Hypertension	6 (26.09%)	10 (29.41%)	11 (19.64%)
	Diabetes	5 (21.74%)	7 (20.59%)	6 (10.71%)

Table No. 2: Surgical Characteristics

Surgical Variable	Total (n=113)	Low Volume (n=23)	Intermediate Volume (n=34)	High Volume (n=56)
Type of Cystectomy	Radical	19 (82.61%)	27 (79.41%)	51 (91.07%)
	Partial	4 (17.39%)	7 (21.59%)	5 (8.93%)
Surgical Approach	Open	23 (100%)	34 (100%)	56 (100%)
Operative Time (min)	Mean (SD)	240 (30)	220 (25)	200 (20)

Table No. 3: Distribution of Postoperative Complications

Complication Type	Total (n=113)	Low Volume (n=23)	Intermediate Volume (n=34)	High Volume (n=56)
Any Complication	33 (29.20%)	9 (39.13%)	12 (35.29%)	12 (21.43%)
Major Complication	8 (7.08%)	3 (13.04%)	3 (8.82%)	2 (3.57%)
Infection	6 (5.31%)	2 (8.70%)	1 (2.94%)	3 (5.36%)
Hemorrhage	4 (3.54%)	2 (8.70%)	1 (2.94%)	1 (1.79%)

Table No. 4: Length of Hospital Stay (LOS) and Mortality Rate.

LOS Variable	Total (n=113)	Low Volume (n=23)	Intermediate Volume (n=34)	High Volume (n=56)
Mean LOS (days)	6.5 (2.0)	7.0 (1.5)	6.5 (1.8)	6.0 (2.2)
LOS >7 days	16 (14.16%)	8 (34.78%)	4 (11.76%)	4 (7.14%)
30-Day Mortality	4 (3.54%)	2 (8.70%)	1 (2.94%)	1 (1.79%)

Table No. 5: Multivariate Analysis of Surgical Volume and Outcomes

Outcome Measure	Adjusted Odds Ratio (95% CI)	P-value
Any Complication	0.75 (0.53-1.06)	0.105
Major Complication	0.61 (0.37-1.00)	0.048
LOS >7 days	1.32 (0.88-1.97)	0.186
30-Day Mortality	0.92 (0.47-1.78)	0.796

DISCUSSION

The findings of this study indicate that the number of surgeries performed may influence the probability of significant complications in individuals receiving cystectomy. This finding aligns with other research that has similarly shown a substantial correlation between the number of surgeries performed and the results after different surgical procedures^[9]. In a research conducted by Finlayson EV et al.^[10] in 2003, it was shown that there is a correlation between a larger number of

surgeries performed and a decreased occurrence of postoperative complications and mortality in patients having major cancer surgery. Similarly, a research conducted by Hanchate AD et al.^[11] in 2010 discovered that performing a larger number of surgeries was linked to reduced occurrences of postoperative complications and death in patients following coronary artery bypass graft surgery.

Our study revealed that the total complication rate was 29.20%, with significant problems seen in 7.08% of patients. These results align with the findings of a research conducted in 2010, which indicated that patients who had cystectomy experienced an overall complication rate of 30.5% and a serious complication rate of 8.3%^[12]. In addition, our research revealed a 30-day mortality rate of 3.54%, which is slightly lower than the 5.1% reported by Wolters M, et al (2017)^[13]. The resemblance in complexity and fatality rates suggests our research sample accurately reflects the whole patient population receiving cystectomy.

Our analysis revealed a noteworthy correlation between surgical volume and major complications, indicating that intermediate and high volume centers had a lower risk of major problems compared to low volume centers. The earlier result aligns with the findings of a research conducted by Konety et al.^[14] in 2007. This study likewise observed a noteworthy correlation between the number of surgeries performed and the occurrence of serious problems in patients having cystectomy. However, our research did not discover a substantial correlation between the number of surgeries performed and the occurrence of complications, length of hospital stay, or mortality rates. This contrasts with the results reported by Konety et al. (2005). The observed disagreement might be attributed to variations in the research's design and sample size. Our study, with a lower sample size, may have lacked sufficient statistical power to identify significant relationships.¹⁵

Study Limitation: Our study has various limitations that should be addressed when interpreting outcomes. First, our prospective research may have biased participant selection and data collection. Second, our investigation was done at one center, which may restrict its applicability. Finally, our small sample size may have prevented us from finding substantial relationships between surgery volume and outcomes. Despite these limitations, this analysis sheds light on how surgical volume affects cystectomy outcomes.

CONCLUSION

The results of our research revealed a significant correlation between the number of surgeries performed and the occurrence of severe complications in individuals having cystectomy. These findings align with other research and emphasize the significance of surgical volume in affecting postoperative results. Healthcare organizations should contemplate the consolidation of certain surgical procedures in high-volume centers to enhance patient outcomes. Additional study is required to have a deeper understanding of the processes behind this correlation and to pinpoint measures for enhancing outcomes in centers with low patient volumes.

Acknowledgment: We extend our gratitude to all the individuals who participated in this research endeavor. We would like to express our gratitude for the support and help given by the staff at the study center. Their inputs were important for the completion of this investigation.

Author's Contribution:

Concept & Design of Study:	Shahjehan
Drafting:	Syed Ameroon Shah, Maham Arooj Zaidi
Data Analysis:	Khalid Islam, Muhammad Zohair, Naseeb Dad

Revisiting Critically:	Shahjehan, Syed Ameroon Shah
Final Approval of version:	Shahjehan

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.213 dated 20.02.2020

REFERENCES

1. Griffiths TL. Action on Bladder Cancer. Current perspectives in bladder cancer management. *Int J Clin Practice* 2013;67(5):435-48.
2. Bruins HM, Veskimäe E, Hernandez V, Neuzillet Y, Cathomas R, Comperat EM, et al. The importance of hospital and surgeon volume as major determinants of morbidity and mortality after radical cystectomy for bladder cancer: a systematic review and recommendations by the European Association of Urology Muscle-invasive and Metastatic Bladder Cancer Guideline Panel. *Eur Urol Oncol* 2020;3(2):131-44.
3. Scarberry K, Berger NG, Scarberry KB, Agrawal S, Francis JJ, Yih JM, et al. Improved surgical outcomes following radical cystectomy at high-volume centers influence overall survival. *InUrologic Oncology: Seminars and Original Investigations*. Elsevier 2018;36(6):308-e11.
4. Goossens-Laan CA, Gooiker GA, van Gijn W, Post PN, Bosch JR, Kil PJ, et al. A systematic review and meta-analysis of the relationship between hospital/surgeon volume and outcome for radical cystectomy: an update for the ongoing debate. *Eur Urol* 2011;59(5):775-83.
5. Cheung G, Sahai A, Billia M, Dasgupta P, Khan MS. Recent advances in the diagnosis and treatment of bladder cancer. *BMC Med* 2013;11(1):13.
6. Witjes JA, Bruins HM, Cathomas R, Compérat EM, Cowan NC, Gakis G, et al. European Association of Urology guidelines on muscle-invasive and metastatic bladder cancer: summary of the 2020 guidelines. *Eur Urol* 2021;79(1):82-104.
7. Lorenz K, Raffaeli M, Barczyński M, Lorente-Poch L, Sancho J. Volume, outcomes, and quality standards in thyroid surgery: an evidence-based analysis—European Society of Endocrine Surgeons (ESES) positional statement. *Langenbeck's Archives Surg* 2020;405:401-25.
8. Shahian DM, Normand SL. Low-volume coronary artery bypass surgery: measuring and optimizing performance. *J Thoracic Cardiovascular Surg* 2008;135(6):1202-9.
9. Schmidt CM, Turrini O, Parikh P, House MG, Zyromski NJ, Nakeeb A, Howard TJ, Pitt HA,

- Lillemoe KD. Effect of hospital volume, surgeon experience, and surgeon volume on patient outcomes after pancreaticoduodenectomy: a single-institution experience. *Archives Surg* 2010; 145(7):634-40.
10. Finlayson EV, Goodney PP, Birkmeyer JD. Hospital volume and operative mortality in cancer surgery: a national study. *Archives Surg* 2003; 138(7):721-5.
 11. Hanchate AD, Stukel TA, Birkmeyer JD, Ash AS. Surgery volume, quality of care and operative mortality in coronary artery bypass graft surgery: a re-examination using fixed-effects regression. *Health Services and Outcomes Res Methodol* 2010;10:16-32.
 12. Liedberg F. Early complications and morbidity of radical cystectomy. *Eur Urol Supplements* 2010; 9(1):25-30.
 13. Wolters M, Oelke M, Lutze B, Weingart M, Kuczyk MA, Chaberny IF, et al. Deep surgical site infections after open radical cystectomy and urinary diversion significantly increase hospitalisation time and total treatment costs. *Urologia Internationalis* 2017;98(3):268-73.
 14. Konety BR, Allareddy V. Influence of post-cystectomy complications on cost and subsequent outcome. *J Urol* 2007;177(1):280-7.
 15. Konety BR, Dhawan V, Allareddy V, Joslyn SA. Impact of hospital and surgeon volume on in-hospital mortality from radical cystectomy: data from the health care utilization project. *J Urol* 2005;173(5):1695-700.

Frequency of Peptic Ulcer Disease in Patients with Chronic Use of Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)

Shakeel Ahmad¹, Muhammad Naeem², Hamid Ullah³ and Liaqat Ali¹

ABSTRACT

Objective: To ascertain how often individuals on chronic nonsteroidal anti-inflammatory Drugs (NSAIDs) develop peptic ulcer disease.

Study Design: A Cross Sectional Study

Place and Duration of Study: This study was conducted at the Department of Gastroenterology, Qazi Hussain Ahmad Medical Complex Nowshera, from June 2022 to June 2023.

Methods: During the study period all patients with endoscopic findings of peptic ulcer and chronic use of NSAIDs were enrolled for the study. Age, gender, current indications for NSAIDs use, and period of NSAID use were noted. Patients were also assessed for other associated risk factors.

Results: During the study period 184 patients had peptic ulcers on endoscopy. In 15.8% (29/184) of individuals, peptic ulcers was caused by chronic NSAIDs use. Arthritis and malignancy related pain were the most common indications for chronic NSAIDs use and diclofenac was the most commonly used NSAIDs. H. pylori-infection was the most common associated risk factor (48%).

Conclusion: Our study showed the significant association of the chronic NSAIDs use and multiple other risk factors with the peptic ulcer disease. It emphasizes cautious use of NSAIDs in high risk patients, particularly who have multiple other risk factors.

Key Words: Peptic Ulcer Disease, Chronic Use, Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)

Citation of article: Ahmad S, Naeem M, Hamid Ullah, Ali L. Frequency of Peptic Ulcer Disease in Patients with Chronic Use of Nonsteroidal Anti-Inflammatory Drugs (NSAIDs). *Med Forum* 2024;35(1):73-75. doi:10.60110/medforum.350116.

INTRODUCTION

Peptic ulcer is one of the most common diseases of the gastrointestinal tract with estimated lifetime prevalence of approximately 5–10% in the general population.¹ It is defined as a breach in the mucosa of the alimentary tract. It is produced when there is imbalance between the aggressive factors like gastric acid, and proteolytic enzyme and protective factors^{2,3} like mucus and bicarbonate secretion and prostaglandins production, combined with superimposed injury from environmental or immunologic agents.⁴

H. pylori infection, cigarette smoking, alcoholism, psychological stress, ischemia, certain medications like

chronic NSAIDs and corticosteroid use, antiplatelet and anticoagulants are the common risk factors for peptic ulcer disease. Nonsteroidal anti-inflammatory drugs (NSAIDs) are widely prescribed and marketed to treat pain and inflammation, and are therefore, the leading cause of non-H—Pylori peptic ulcers.⁵

Chronic use of NSAIDs causes suppression of mucosal prostaglandin (PGE2) and direct irritative topical effect. Prostaglandins stimulate mucus and bicarbonate production, and regulates blood flow to the GI tract.⁶ Its inhibition by NSAIDs leads to loss of alimentary protective factors and in combination with multiple environmental risk factors and H. pylori infection leads to peptic ulcer disease.⁷

Peptic ulcer disease predominantly presents with dyspeptic symptoms like epigastric burning sensation, epigastric pain, pyrosis, nausea, vomiting, bloating and belching.⁸ It may sometimes present with complications like anemia, hematemesis, melena, perforation, gastric or duodenal outlet obstruction or peptic stricture. Malignant formation is rare and related to underlying gastritis.^{9,10}

Since H. pylori infection is decreasing, NSAID-related peptic ulcers are rising because of the injudicious use of the NSAIDs without a prescription^{11,12}. The aim of our study is to find out the frequency of Peptic ulcer disease in our population because of chronic NSAIDs use so

¹. Department of Gastroenterology, DHQ Hospital, Mardan.

². Department of Gastroenterology, Mardan Medical Complex, Mardan.

³. Department of Gastroenterology, Qazi Hussain Ahmad Medical Complex, Nowshera.

Correspondence: Hamid Ullah, Assistant Professor, Qazi Hussain Ahmad Medical complex, Nowshera.

Contact No: 0312 9565134

Email: drhamidullah222@gmail.com

Received: August, 2023

Accepted: September, 2023

Printed: January, 2024

that the issue is highlighted and appropriate steps taken for its prevention in the future.

METHODS

This is a cross sectional study conducted in the Gastroenterology department of Qazi Hussain Ahmad Medical Complex, Nowshera from June 2022 to June 2023. All patients with endoscopic findings of peptic ulcer and chronic use of NSAIDs were enrolled for the study. Chronic NSAID use is defined as if these medications are taken more than three times a week for more than three months.

Age, gender, current indications for NSAIDs use, and period of NSAID use were noted. Patients were also assessed for associated risk factors like H. pylori infection, alcoholism, smoking, tobacco chewing etc.

RESULTS

During the study period 184 patients had peptic ulcers on endoscopy. In 15.8% (29/184) of individuals, peptic ulcers was caused by chronic NSAIDs use (figure 1). Arthritis and malignancy related pain were the most common indications for chronic NSAIDs use (table 1). Diclofenac was the most commonly used NSAIDs, linked to peptic ulcer in 41.37% (12/29) of individuals (table 2). Duodenal ulcers outnumbered gastric ulcers, 41.37% (12/29) and 34.48% (10/29) respectively (table 3). H. pylori-infection was also noted in 48% (14/29) of the patients (figure 2). Other peptic ulcer risk factors were noted and also shown in figure 2.




Figure 1: Etiological pattern of PUD

Table No. 1: Common indications for chronic NSAIDs use

Indications	Percentage (N) of Patients
Arthritis	38% (11/29)
Malignancy related pain	31% (9/29)
Chronic body aches	10.3% (3/29)
Chronic Headache	10.3% (3/29)
Others	10.3% (3/29)

Table No. 2: NSAIDs Linked to Peptic Ulcers in the Study

NSAID	Percentage of Patients with Peptic Ulcers
Diclofenac	41.37% (12/29)
Neporoxen/piroxicam	20.68% (6/29)
Others (ibuprofen, mephenamic acid, aceclofenac, ketorolac etc.)	20.68% (6/29)
Combination of NSAIDs	17.24% (5/29)

Table No. 3: Location of Peptic Ulcers in Patients

Type of Ulcer	Percentage of Patients with Peptic Ulcers
Duodenal Ulcer	41.37% (12/29)
Gastric Ulcer	34.48% (10/29)
Ulcers in both stomach and duodenum	24.13% (7/29)




Figure No. 2: Associated risk factors

DISCUSSION

Peptic ulcer disease is a well-known complication of NSAID use. Our study showed comparatively high prevalence (approximately 16%) of peptic ulcer disease in patients with chronic NSAIDs use as compared to other studies which showed the prevalence of 6-11%^{13,14}.

In our study diclofenac and naproxen/piroxicam are the most common NSAIDs associated with peptic ulcer, which is similar to findings in other studies¹². However, in our study naproxen/piroxicam was the second most common NSAID used as compared to aspirin as in other studies¹². Diclofenac is the most widely prescribed analgesic in our population because of the easy availability and low cost. This finding emphasizes the need for cautious use of this drug, particularly in patients who already have multiple other risk factors for peptic ulcer disease like having a history of prior peptic ulcer or are receiving concurrent steroid treatment.

In our study like most of the other studies, NSAIDs induced duodenal ulcer outnumbered gastric ulcer. According to our data the correlation between NSAID-

induced peptic ulcers and *H. pylori* infection exists in approximately half of the patients with peptic ulcer¹⁵. It emphasizes the importance of *H. pylori* testing and eradication prior to initiation of NSAIDs use¹⁶.

Finally, the research emphasizes how important it is to take into account various other risk variables like drinking alcohol and smoking when discussing NSAID-related peptic ulcers. In summary, this research offers important new information on the prevalence and risk factors of NSAID-induced peptic ulcer disease in local population. With this knowledge, medical professionals may prescribe NSAIDs more cautiously and create individualized treatment plans for individuals who are susceptible to peptic ulcers because of the associated risk factors¹⁷.

CONCLUSION

Our study showed the significant association of the chronic NSAIDs use particularly diclofenac and aspirin with the peptic ulcer disease. It emphasizes cautious use of NSAIDs in high risk individuals, particularly who have multiple other risk factors like *H. pylori* infection.

Author's Contribution:

Concept & Design of Study:	Shakeel Ahmad
Drafting:	Muhammad Naem, Hamid Ullah
Data Analysis:	Liaqat Ali
Revisiting Critically:	Shakeel Ahmad, Muhammad Naem
Final Approval of version:	Shakeel Ahmad

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.122/BKMC dated 15.06.2022

REFERENCES

- Perez-Aisa MA, Del Pino D, Siles M, Lanas A. Clinical trends in ulcer diagnosis in a population with a high prevalence of *Helicobacter pylori* infection. *Alimentary Pharmacol Therapeutics* 2005;21(1):65-72.
- Phillips WJ, Currier BL. Analgesic pharmacology: II. Specific analgesics. *J Am Acad Orthop Surg* 2004;12(4):221-33.
- Narayanan M, Reddy KM, Marsicano E. Peptic Ulcer Disease and *Helicobacter pylori* infection. *Mo Med* 2018;115(3):219-224.
- Lanas Á, Carrera-Lasfuentes P, Arguedas Y, García S, Bujanda L, Calvet X, et al. Risk of upper and lower gastrointestinal bleeding in patients taking nonsteroidal anti-inflammatory, anti-inflammatory drugs, antiplatelet agents, or anticoagulants. *Clin Gastroenterol Hepatol* 2015;13(5):906-12.e2
- Lau JY, Sung JJ, Metz DC, Howden CW. A systematic review of the epidemiology of

complicated peptic ulcer: incidence, recurrence, risk factors, and mortality. *Gastroenterol* 2008;134(Suppl. 1): A32.

- Ehlin AG, Montgomery SM, Ekblom A, Pounder RE, Wakefield AJ. Prevalence of gastrointestinal diseases in two British national birth cohorts. *Gut* 2003;52:1117-21.
- Rosenstock S, Jorgensen T, Bonnevie O, Andersen L. Risk factors for peptic ulcer disease: a population-based prospective cohort study comprising 2416 Danish adults. *Gut* 2003;52: 186-93.
- Konturek SJ, Bielanski W, Plonka M, et al. *Helicobacter pylori*, nonsteroidal anti-inflammatory, anti-inflammatory drugs, and smoking in risk patterns of gastroduodenal ulcers. *Scand J Gastroenterol* 2003;38: 923-30.
- Lassen A, Hallas J, Schaffalitzky de Muckadell OB. Complicated and uncomplicated peptic ulcers in a Danish county 1993-2002: a population-based cohort study. *Am J Gastroenterol* 2006;101:945-53.
- Garcia Rodriguez LA, Hernandez-Diaz S. Risk of uncomplicated peptic ulcer among users of aspirin and nonaspirin nonsteroidal anti-inflammatory drugs. *Am J Epidemiol* 2004;159:23-31.
- Bartholomeeusen S, Vandenbroucke J, Truyers C, Buntinx F. Time trends in the incidence of peptic ulcers and oesophagitis between 1994 and 2003. *Br J Gen Pract* 2007;57:497-9.
- Hamid S, Yakoob J, Jafri W, Islam S, Abid S, Islam M. Frequency of NSAID induced peptic ulcer disease. *J Pak Med Assoc* 2006;56(5):218.
- Liang CM, Yang SC, Wu CK, Li YC, Yeh WS, Tai WC, et al. Risk of Recurrent Peptic Ulcer Disease in Patients Receiving Cumulative Defined Daily Dose of Nonsteroidal Anti-Inflammatory Drugs. *J Clin Med* 2019;8(10):1722.
- Tang CL, Ye F, Liu W, Pan XL, Qian J, Zhang GX. Eradication of *Helicobacter pylori* infection reduces the incidence of peptic ulcer disease in patients using nonsteroidal anti-inflammatory drugs: A meta-analysis. *Helicobacter* 2012;17:286-296.
- Laine L, Curtis SP, Cryer B, Kaur A, Cannon CP. Assessment of upper gastrointestinal safety of etoricoxib and diclofenac in patients with osteoarthritis and rheumatoid arthritis in the Multinational Etoricoxib and Diclofenac Arthritis Long-term (MEDAL) program: a randomized comparison. *The Lancet* 2007;369(9560):465-73.
- Quan C, Talley NJ. Management of peptic ulcer disease unrelated to *Helicobacter pylori* or (NSAIDs). *The Am J Gastroenterol* 2002;97(12): 2950-61.
- Kuna L, Jakab J, Smolic R, Raguz-Lucic N, Vcev A, Smolic M. Peptic ulcer disease: a brief review of conventional therapy and herbal treatment options. *J Clin Med* 2019;8(2):179.

Comparison of the Skin Closures Using Staples Versus Prolene Sutures in Patients Undergoing Clean Elective Abdominal Surgeries.

Skin Closures Using Different Sutures in Abdominal Surgery

Shumaila Naseer¹, Tariq Hayat Khan², Ayaz Gul³ and Nida Mumtaz²

ABSTRACT

Objective: To assess the surgical site infection and skin closure time of individuals receiving clean elective abdominal operations with prolene sutures vs staples.

Study Design: Randomized Controlled Trial study

Place and Duration of Study: This study was conducted at the Department of Surgery, Lady Reading hospital, Peshawar from 1st Jan 2021 to July 2021.

Methods: This Randomized Controlled Trial study was conducted with the necessary approvals from the ethical board and research committee of the CPSP at the Department of Surgery, Lady Reading hospital, Peshawar. A total of 124 patients, of various genders, underwent clean elective abdominal surgery and were included in the study. Patients in group A underwent skin closure using the staple method, while patients in group B underwent skin closure using the prolene suture method after surgery.

Results: Participants in this research ranged in age from 18 to 65, with Group A averaging 45.048±7.83 years and Group B 43.451±9.27 years. SSI rates differed significantly between groups A and B. SSI occurred in 19 (30.6%) of group A patients and 37 (59.7%) of group B patients (P= 0.001). The mean skin closure time in group A was substantially lower than group B (p=0.000). Group A had an average closure time of 126.774±32.78 seconds, whereas Group B had 459.677±60.43 seconds.

Conclusion: Our study results indicate that skin staples have been found to result in lower rates of wound infection compared to sutures in clean elective surgeries.

Key Words: Clean procedures, Skin closures, Staples, Prolene sutures, Surgical Site Infection, Closure time

Citation of article: Naseer S, Khan TH, Gul A, Mumtaz N. Comparison of the Skin Closures Using Staples Versus Prolene Sutures in Patients Undergoing Clean Elective Abdominal Surgeries. Med Forum 2024;35(1):76-79.doi:10.60110/medforum.350117.

INTRODUCTION

A wound closure technique should be easy to use, have similar strength across the incision, ensure skin apposition till healing, avoid wound infection, and be aesthetically pleasing. A good wound closure material is cheap, non-allergic, and easy to create and use.¹ Any skin approximation method must keep the margins in place long enough for healing². The wound closure technique and material contribute to wound infection because the substance acts as a foreign entity and

causes a variable inflammatory response that limits tissue blood flow and causes ischemia.³ Braided suture gaps may harbor pathogens. The surgeon should employ less-traumatic sutures with enough mechanical strength. To minimize scarring, remove sutures promptly⁴. Carefully suture incisions and wounds, using suitable closure methods⁵. Surgical wounds are usually sutured. The surgeon may use continuous or interrupted, natural or synthetic, absorbable or non-absorbable, single filament or braided sutures, depending on wound length and location⁶. Staples may be better for surgical wound closure because to their low tissue reactivity. Contaminated wounds are more resistant to infection because foreign material cannot enter and damage the local immune response. Staples may reduce incision diameter, wound healing time, local inflammation, and cross marks.⁷⁻⁹

Inert polypropylene monofilament sutures are one type. They reduce infection risk. Silk and other coated sutures induce infection more often. Metal skin staples made of stainless steel are easy to install. They may close skin 80% faster than subcuticular or interrupted suturing¹⁰. Previous research found substantial differences in skin closure time and surgical site

¹. Department of General Surgery, Jinnah Medical College, Peshawar.

². Department of General Surgery / Surgery³, Lady Reading hospital, Peshawar.

Correspondence: Dr. Ayaz Gul, Assistant Prof. of General Surgery, Lady Reading Hospital, Peshawar.

Contact No: 03339179481

Email: drayazgul333@gmail.com

Received: July, 2023

Accepted: September, 2023

Printed: January, 2024

infection (SSI) incidence between prolene suture and staples groups: 251.07 ± 28.61 vs. 87.28 ± 17.20 seconds ($p < 0.0001$) and 61.2% vs. 38.8% ($p 0.024$). Current study will test skin staples and prolene sutures for SSI and skin closure time following elective abdominal surgery.

METHODS

With CPSP ethics board and research council approval, Lady Reading Hospital's Department of Surgery in Peshawar, Pakistan, conducted this Randomized Controlled Trial from 1st Jan 2021 to July 2021. The study comprised all eligible out-of-department patients. The patients were told about the research's purpose, benefits, and surgical method before signing their informed consent. Every patient was examined and historyd. Each patient was randomly assigned to two groups using a blocking approach. Group A patients received staples for skin closure after surgery, whereas group B got prolene sutures. A single, qualified general surgeon with at least five years of experience conducted each treatment.

Using a timer, the student tracked the time between the first and last skin sutures throughout the surgery. All patients received standard post-op care.

Diabetes, liver cirrhosis, using steroids within six months, and severe renal sickness were eliminated from the study owing to their potential to bias the results. The next 30 days, all patients were followed for SSI. Data was analyzed using SPSS 20. Chi square was utilized to compare SSI across groups, and independent samples measured skin closure time. A significance level of < 0.05 was used for the T Test.

RESULTS

The study has 18–65-year-old participants. In Group A, the average age was 43.45 years $\pm 9.27SD$, weight was 69.983 Kg $\pm 10.94SD$, height was 1.538 meters $\pm 0.11SD$, BMI was 29.837 Kg/m² $\pm 5.29SD$, and skin closure time was 459.677 sec $\pm 60.43SD$. Ages ranged from 18 to 65 in the research. at Group B. Mean scan time is 0.7463, however age, weight, height, and BMI p-values are not significant. A 0.000 p-value showed that both groups closed differently.

Group A included 43 males (69.4%) and 47 women (75.8%), whereas group B had 19 women (30.6%) and 15 men (24.2%). Males dominated both groups but were statistically insignificant with a p-value of 0.5463. Table III shows surgery distribution by group within each group. Group A had 23 exploratory laparotomies (37.1%), 25 open appendices (40.3%), and 14 hernia repairs (22.6%).

Surgical site infection (SSI) occurred in 19 (30.6%) patients in group A and similar pattern was found in group B, although neither group was significantly different (-v a l u e = 0.7673 Fig. 1 SSI was substantially greater in Group B than Group A

($p=0.001$). Compared to prolene suture, staple is more practicable. Table 1




Figure No. 1: Indication of surgery in both groups

Table No. 1: SSI in both groups n=124

SSI	N=62		P-valve
	Group-A	Group-B	
Yes	19 (30.6%)	37 (59.7%)	0.001
No	43 (69.4%)	25 (40.3%)	
Total	62 (100%)	62 (100%)	

DISCUSSION

NICE's latest surgical site infection prevention and treatment guidelines addresses suturing and surgical site infections for the first time. This is the first time that sutures are recommended over staples for closing the skin post-caesarean section to prevent wound dehiscence in postpartum mothers. However, sutures have little evidence to support use over staples in other surgeries. Different closure procedures affect surgical site infections differently based on the anatomical location. In craniofacial surgery, suture and staple groups had comparable infection rates (2 and 3%)¹². Two meta-analyses comparing staples and sutures in orthopedic procedures found conflicting results on surgical site infection risks¹³. An revised meta-analysis indicated no significant difference in SSI risk between staples and sutures¹⁴. Cochrane review data on coronary artery bypass surgery showed comparable findings¹⁵. In addition to skin staples, drape fusion, hernia mesh fusion, and intestinal anastomosis may be employed in surgery.^{16,17} Nowadays, most skin staples are stainless steel, although absorbable ones were used sometimes. Animal studies showed low inflammation with absorbable staples.¹⁸ Another author found that skin staplers are only faster in elective breast and abdominal surgeries¹⁹. In the staples and suture groups, the mean closure time was 80 and 242 seconds, respectively. Suturing and stapling cause comparable wound infections. In the beginning and during removal, staples hurt more.²⁰ Some research found that staples are less uncomfortable than sutures after six weeks.²¹ We found that the suture group had a closure time nearly three

times longer than the staples group (126.774 ± 32.78 seconds vs. 459.677 ± 60.43 seconds, p value < 0.0001). Stapling may be seven times faster than stitching.^{22,23} Prolene or nylon sutures scar face better than staples in cosmetic surgery.²⁴ For abdominal procedures, sutures were more cosmetically pleasing.²⁵ Cosmetic appearance and patient satisfaction with suture and staples may be the same at six weeks in elective cesarean sections.¹⁹ Sutures were more cosmetic than staples for emergency cesarean sections. Staples lengthen hospital stays.^{25,26} According to a metaanalysis by Smith et al., skin staples in orthopedic surgery are linked to greater infection rates. In hip and knee surgery, staples are not advised.¹³ Another research found comparable findings for orthopedic surgery wound infection. A research found SSI in 19.6% of patients in group A and 37.7% in group B ($P = 0.001$). In one research, prolene suture and staples groups had 61.2% and 38.8% (p value 0.024) surgical site infections (SSIs)¹¹. Since sutures and staples have pros and downsides, the contradictory evidence may be justified. Metal staples may be less irritative and more infection-resistant than least reactogenic sutures²⁷. Staples are recommended in emergencies because they close skin quicker, saving 5.5 to 8 minutes. Staples may cause staple track development, hair follicle damage, perspiration and sebaceous gland damage, bacterial migration into the wound bed, and pain during removal. Tight skin closure that retains dermal structure may avoid surgical site infection, since the patient's fetoplasm is the main source of infection. Intracutaneous sutures tighten the skin without harming it. Patients may find absorbable sutures more pleasant since they may remain in the wound without removal. Additionally, sutures cost just 20% of staples²⁸. All suture materials are alien to human tissue and may promote inflammation, compromising wound healing and increasing infection risk²⁹. Surgeons choose staples for midline incision closure, despite potential hazards and benefits.

CONCLUSION

We found that skin staples cause less wound infection than sutures in clean elective surgeries. Staples resemble skin quicker than stitches. Well-designed randomized controlled studies with large sample numbers are required to corroborate this since surgeons need better evidence for decision-making. Future studies should address obesity and other postoperative complications risk factors.

Author's Contribution:

Concept & Design of Study: Shumaila Naseer
 Drafting: Tariq Hayat Khan, Ayaz Gul
 Data Analysis: Nida Mumtaz
 Revisiting Critically: Shumaila Naseer, Tariq

Hayat Khan

Final Approval of version: Shumaila Naseer

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.998/LRH/MTI dated 02.11.2020

REFERENCES

1. Krishnan R, MacNeil SD, Malvankar-Mehta MS. Comparing sutures versus staples for skin closure after orthopedic surgery: systematic review and meta-analysis. *BMJ Open* 2016;6(1)23- 9.
2. Imamura K, Adachi K, Sasaki R, Monma S, Shioiri S, Seyama Y, et al. Randomized comparison of subcuticular sutures versus staples for skin closure after open abdominal surgery: a multicenter open-label randomized controlled trial. *J Gastrointest Surg* 2016;20(12):2083-92.
3. Henriksen NA, Deerenberg EB, Venclauskas L, Fortelny RH, Miserez M, Muysoms FE. Meta-analysis on materials and techniques for laparotomy closure: the MATCH review. *World J Surg* 2018;42(6):1666- 78.
4. Kumar R, Hastir A, Goyal S, Walia RS. Sutures versus staplers for skin closure of midline incision in laparotomy patients and their outcome. *Pain* 2017;5:12-5.
5. Varghese F, Gamaliel J, Kurien JS. Skin stapler versus sutures in abdominal wound closure. *Int Surg J* 2017;4(9):3062-6.
6. Cirocchi R, Randolph JJ, Montedori A, Arezzo A, Mearini EE, Abraha I, et al. Staples versus sutures for surgical wound closure in adults (Protocol). *Cochrane Database of Systematic Reviews* 2014(8):1-3.
7. Henriksen NA, Deerenberg EB, Venclauskas L, Fortelny RH, Miserez M, Muysoms FE. Meta-analysis on materials and techniques for laparotomy closure: The MATCH Review. *World J Surg* 2018:1-3.
8. Kuroki LM, Mullen MM, Massad LS, Wu N, Liu J, Mutch DG, et al. Wound complication rates after staples or suture for midline vertical skin closure in obese women: a randomized controlled trial. *Obstet Gynecol* 2017;130(1):91-9.
9. Oswal S, Borle R, Bhola N, Jadhav A, Surana S, Oswal R. Surgical staples: a superior alternative to sutures for skin closure after neck dissection-a single-blinded prospective randomized clinical study. *J Oral Maxillofac Surg* 2017;75(12):2707 e1- e6.
10. Slade Shantz JA, Vernon J, Morshed S, Leiter J, Stranges G. Sutures Versus staples for wound closure in orthopedic surgery: a pilot randomized controlled trial. *Patient Saf Surg* 2013;7(1):6- 10.

11. Basit A, Abbasi SH, Haider S, Kiani YM, Shah FH. To compare outcomes of stainless skin staples and polypropylene sutures for skin closure in clean elective surgeries. *Isra Med J* 2018;10(1):32-5.
12. Sidebottom AJ, Grogan J, May P, Richardson D. Prospective comparison of methods of closure of the coronal flap after craniofacial surgery in children. *Br J Oral Maxillofac Surg* 2003;41:309.
13. Smith TO, Sexton D, Mann C, Donell S. Sutures versus staples for skin closure in orthopedic surgery: meta-analysis. *BMJ* 2010;340:c1199.
14. Krishnan RJ, Crawford EJ, Syed I. Is the Risk of Infection Lower with Sutures than with Staples for Skin Closure After Orthopaedic Surgery? A Meta-analysis of Randomized Trials. *Clin Orthop Relat Res* 2019; 477:922.
15. Biancari F, Tiozzo V. Staples versus sutures for closing leg wounds after vein graft harvesting for coronary artery bypass surgery. *Cochrane Database Syst Rev* 2010:D8057.
16. Amin N, Fu B, Rutka J, Das P. Stapled double head and neck drape for otological procedures. *J Laryngol Otol* 2013;127(11):1139-40.
17. Khan AA, Majeed S, Shahzadi M, Hussain SM, Ali MZ, Siddique K. Polypropylene suture versus skin staples for securing mesh in lichtenstein inguinal hernioplasty. *J Coll Physic & Surg Pak* 2014;24(2):88-90.
18. Malard O, Duteille F, Darnis E, Espitalier F, Perrot P, Ferron C, et al. A novel absorbable stapler provides patient-reported outcomes and cost-effectiveness noninferior to subcuticular skin closure: a prospective, single-blind, randomized clinical trial. *Plastic Reconstructive Surg* 2020;146(6):777e-89e.
19. Ku D, Koo DH, Bae DS. A prospective randomized control study comparing the effects of dermal staples and intradermal sutures on postoperative scarring after thyroidectomy. *J Surgical Res* 2020;256:413-21.
20. Tseng TH, Jiang CC, Fu SH, Lee TL, Chuang YH, Chiang H. Topical anesthesia for staple removal from surgical wounds on the knee: a prospective, double-blind, randomized trial. *J Surg Res* 2017;215:167-72.
21. Rousseau JA, Girard K, Turcot-Lemay L, Thomas N. A randomized study comparing skin closure in cesarean sections: staples vs subcuticular sutures. *Am J Obstet Gynecol* 2019;200(3):265.
22. O'Neill JK, Robinson P, Giddins GE. Staples for intraoperative skin retraction in hand surgery. *J hand Microsurg* 2014;6(2):100-101.
23. La Rosa M, Omere C, Redfern T, Abdelwahab M, Spencer N, Villarreal J, et al. The impact of low-dose versus high-dose antibiotic prophylaxis regimens on surgical site infection rates after cesarean delivery. *Archives Gynecol Obstet* 2020;301:69-73.
24. Simcock JW, Armitage J, Dixon L, MacFarlane K, Robertson GM, Frizelle FA. Skin closure after laparotomy with staples or sutures: a study of the mature scar. *ANZ J Surg* 2014;84(9):656-9.
25. Sharma C, Verma A, Soni A, Thusoo M, Mahajan VK, Verma S. A randomized controlled trial comparing cosmetic outcome after skin closure with staples' or subcuticular sutures' in emergency cesarean section. *Arch Gynecol Obstet* 2014;290(4):655-59.
26. Syed KA, Gandhi R, Davey JR, Mahomed NN. Risk of wound infection is greater after skin closure with staples than with sutures in orthopaedic surgery. *J Bone Joint Surg Am* 2014;92(16):2732.
27. Radhae R, Pallavi A, Prathibha A, Susnata C. Abdominal incisions and sutures in obstetrics and gynaecology. *The Obstet Gynaecol* 2014:13.
28. Hochberg J, Meyer KM, Marion MD. Suture choice and other methods of skin closure. *Surg Clin North Am* 2009;89:627.
29. Rahbari NN, Knebel P, Diener MK. Current practice of abdominal wall closure in elective surgery —Is there any consensus? *BMC Surg* 2009;9:8.

The Status of Hepatitis B Vaccination Among Hemodialysis Patients at a Tertiary Care Hospital

Status of
Hepatitis B
Vaccination
Among
Hemodialysis

Shad Muhammad, Arbab Muhammad Ali and Muhammad Ikram

ABSTRACT

Objective: To ascertain the immunization status for hepatitis B in patients with chronic renal disease who are reliant on dialysis.

Study Design: A cross-sectional study

Place and Duration of Study: This study was conducted at the Nephrology, department. Peshawar's Lady Reading Hospital, from July 1, 2021, to December 31, 2021.

Methods: Those receiving hemodialysis for chronic renal disease were included. The anti-HBS antibody titer in the patients' blood after three or four doses of the 20 microgram hepatitis B vaccine was used to validate the patients' hepatitis B vaccination status. A cut-off value of ≥ 10 IU/L was applied to the anti-HBS antibody titer.

Results: 109 patients in all were enrolled. The patients' ages varied from 20 to 60. The patients' average age was 49.80 ± 5.245 years. The male to female ratio was 1.6:1. Of the patients, 81 (74.3%) got three doses of the vaccine, while 28 (25.7%) received four doses. 56 patients (51.4%) had vaccination records.

Conclusion: CKD patients are less likely to develop ≥ 10 IU/L of anti-HBS antibodies after hepatitis vaccination. Patients who receive vaccination in the early course of the disease are more likely to develop better response.

Key Words: Chronic Kidney Disease, HBV vaccination, Hemodialysis

Citation of article: Muhammad S, Ali AM, Ikram M. The Status of Hepatitis B Vaccination Among Hemodialysis Patients at a Tertiary Care Hospital. Med Forum 2024;35(1):80-82. doi:10.60110/medforum.350118.

INTRODUCTION

Because the kidneys play a critical role in eliminating harmful metabolic products—the buildup of which might have detrimental consequences on human health—maintaining renal function is essential to overall wellbeing.^[1] A patient with chronic kidney disease (CKD) has a glomerular filtration rate (GFR) of less than 60 mL/min per 1.73 m², a structural change in the renal parenchyma, and a loss of renal function that lasts longer than three months.^[2] Renal transplantation and peritoneal dialysis are additional options, even though hemodialysis is mostly provided in clinical settings as renal replacement treatment for patients with chronic kidney disease.^[3] Hepatitis B virus (HBV) infection is a major worldwide health hazard, with 150 million people living with its chronic carrier condition. Due to the fact that dialyzed patients are more

susceptible to blood and its products than the general population is, as well as the risk of contaminated hemodialysis equipment and supplies, the incidence of HBV is significantly higher in this group.^[4] HBV prevalence in CKD patients on hemodialysis has been shown to vary from 1.2 to 6.6%. According to current standards, hepatitis B immunization is advised for all patients with chronic kidney disease who are reliant on dialysis since it not only protects against hepatitis B but also improves patient survival. Despite the fact that chronic kidney disease (CKD) is an immunocompromised condition, there is currently a lack of good immunization against HBV among CKD patients, especially in underdeveloped nations.^[5] Amjad et al. observed that 19.9% of CKD patients who were on dialysis had received an HBV vaccination.^[6] A different research by Guimaraes et al, found that 59.2% of dialysis-dependent CKD patients had received an HBV vaccination.^[7] Patients with chronic renal disease have a relatively high prevalence of HBV infection. There is relatively little information on the frequency of HBV vaccinations in CKD patients receiving dialysis, despite current recommendations recommending immunization against HBV in all CKD patients. Furthermore, as no local research has been done on this topic recently, the findings of studies done on other groups cannot be extrapolated to our community. Consequently, I intended to ascertain the hepatitis B vaccination status among dialysis-dependent chronic kidney disease (CKD) patients based on the prevalence of HBV

Department of Nephrology, MTI, Lady Reading Hospital, Peshawar.

Correspondence: Muhammad Ikram, Assistant Professor, Nephrology, Lady Reading Hospital, Peshawar.

Contact No: 03339354462

Email: mikramakhunzada@gmail.com

Received: July, 2023

Accepted: September, 2023

Printed: January, 2024

vaccinations in our community. The findings of my research will provide our local health officials up-to-date data on HBV vaccination rates among CKD patients, which will be useful for future planning.

METHODS

A cross-sectional study conducted at Peshawar's Lady Reading Hospital's nephrology department from July 1 to December 31, 2021. Sample size was calculated using the WHO sample size computation and the estimated frequency of hepatitis B vaccinations (p = 19.9%). 7.5% error, 95% confidence. Sequential sampling without probability. This study covered all 20–60-year-old dialysis patients. Patients with HIV, immunosuppressive medicines, blood dyscrasias, liver or renal illness, or congenital or acquired immunodeficiency problems. Cancer and transplants were excluded.

Both the LRH research department and ethics committee approved the recruitment of 109 eligible patients. Each patient provided informed consent, ensuring anonymity and no risk from this investigation. Age, gender, stadiometer height, and weighing scale weight in kilograms were recorded. After a comprehensive abdomen and general physical exam, the patient's medical history, dialysis dependence, CKD cause, and duration were noted. The patient's medical history was examined to ascertain their hepatitis B vaccination status, including dosages and time since immunization. Patients who got an HBV vaccination or had clear records were documented. The serum anti-HBs antibody titer was tested using ELISA to determine hepatitis B vaccination status. ELISA analysis was performed at the hospital laboratory within

30 minutes of sample collection to detect anti-HBs antibody titer. The operational criteria (serum anti-HBs antibody titer > 10 IU/L) were used to gather data on hepatitis B vaccination status. IBM-SPSS 22 was used to analyze the data.

RESULTS

Patients in this study had a mean age of 49.80 ± 5.245 years, weight of 45.22 ± 7.101 kg, BMI of 21.381 ± 1.020, and disease duration of 22.061 ± 2.511. A total of 56 (51.3%) patients with anti-HBs antibody titers ≥10micrograms were confirmed vaccinated, as shown in Table 1. Table 2 shows immunization status by gender, age, BMI, vaccine doses, and sickness duration.

Table No.1: Patients Demographics

		Frequency	%tage
Gender	Male	68	62.4
	Female	41	37.6
Age	30Y or >	74	67.9
	<30 Y	35	32.1
BMI (kg/m2)	20 kg/m2 or below	48	44
	More than 20 kg/m2	61	56
VACCINE DOSES	03 Doses	81	74.3
	04 Doses	28	25.7
DISEASE DURATION	24 months or more	77	70.6
	Less than 24 months	32	29.4
VACCINATION STATUS	Vaccinated	56	51.4
	Unvaccinated	53	48.6

Table No. 2: Results according to gender, age, BMI, vaccination dose and disease duration

	Gender		Age (yrs)		BMI (kg/m2)		Vaccine Doses		Disease Duration (Months)	
	Male(68)	Female (41)	≥ 40 years (74)	< 40 years (109)	≥ 20 (48)	< 20 (61)	03 doses (81)	04 doses (28)	≥ 24 months (77)	< 24 months (32)
N=109										
Vaccinated (56, 51.4%)	37 (54.4%)	19 (46.3%)	35 (47.3%)	21 (60%)	27 (56.2%)	29 (47.5%)	39 (48.1%)	17 (60.7%)	32 (41.5%)	24 (75.0%)
Unvaccinated (53, 48.6%)	31 (45.6%)	22 (53.7%)	39 (52.7%)	14 (40%)	21 (43.8%)	32 (52.5%)	42 (51.9%)	11 (39.3%)	45 (58.5%)	08 (25.0%)
p value	0.414		0.215		0.366		0.251		0.001	

DISCUSSION

Among CKD patients who had HBV immunization, 56 (51.3%) had anti-HBS antibody titers ≥10 micrograms. A minimum of 10 micrograms of anti-HBS antibody is suggested for hepatitis B prevention. No significant connection was found between HBV vaccination response and gender, age, BMI, or vaccine doses. Though more individuals who got 04 doses of anti-HBV vaccination developed antibodies than those who

received 03 doses, this response was not statistically significant (p = 251). Duration of sickness correlated with vaccination response (p = 0.001). Low response may be linked to immune system weakening as sickness duration increases.^[8] Hepatitis B vaccination rates in our country are low despite nephrology groups and the CDC's advice. Patients' low socioeconomic position may explain this. CKD patients' hepatitis B virus vaccination status is little investigated, however several studies have examined vaccine response in CKD

patients. An early 1990s UK survey found that just 5% of dialysis units consistently immunized patients.⁹ Vaccination rates in the US rose from 47% to 56% between 1997 and 2002.^[10] Our findings indicated 51.3% HBV-vaccinated patients. No local data was available to compare our findings. Compared to our findings, several emerging nations have superior immunization levels. Brazil had over 60% of CKD patients immunized against HBV, with 15% of them incompletely.^[11] Our findings are concerning since CKD patients are high-risk populations and vaccination is the best strategy to avoid HBV infection, coupled with segregation of HBV patients and their equipment and general infection management.^[5] This low immunization rate suggests that nephrologists and dialysis clinics seldom follow guidelines. The US Renal Data System 2011 Annual Data Report found that men started hemodialysis more than females in 2009. Maric also found that diabetic males are more likely to acquire CKD. We found similar results. Gender did not affect CKD patients' hepatitis B vaccination. As in earlier trials, most of our CKD patients were over 40. The mean age of CKD patients in India was 51 years, whereas in China it was 63.6 years.^[12, 13] We found no significant connection between vaccination status and patient age or CKD duration. While older age has been linked to reduced vaccination rates in the general population, we did not find any data on CKD patients' hepatitis B vaccination status.

In the UK, author found that most CKD patients are poor^[14]. HBV vaccination rate was substantially linked with socioeconomic status in our research; lower socioeconomic class patients had lower immunization rates. Author found comparable findings in the overall population.^[15] Lower- and lower-middle-class patients may not be able to afford hepatitis B vaccine or understand its importance.

CONCLUSION

Vaccination status (in terms of titer response to HBV vaccine, i.e. ≥ 10 IU/L) among chronic kidney disease patients is low. Compared to the conventional 03 doses, the response is better in patients who receive 04 doses. Patients in the early course of the disease are likely to show better response to HBV vaccines.

Author's Contribution:

Concept & Design of Study:	Shad Muhammad
Drafting:	Arbab Muhammad Ali, Muhammad Ikram
Data Analysis:	Muhammad Ikram
Revisiting Critically:	Shad Muhammad, Arbab Muhammad Ali
Final Approval of version:	Shad Muhammad

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.1012/LRH/MTI dated 15.09.2020

REFERENCES

- Fraser SD, Blakeman T. Chronic kidney disease: identification and management in primary care. *Pragmat Obs Res* 2016;7:21-32.
- Kakitapalli Y, et al. Detailed Review of Chronic Kidney Disease. *Kidney Dis (Basel)* 2020;6(2): 85-91.
- Um e K, Khan S, Ahmad L. Impact of hemodialysis on the wellbeing of chronic kidney diseases patients: a pre-post analysis. *Middle East Current Psychiatr* 2020;27(1):54.
- Gasim GI, Bella A, Adam I. Immune response to hepatitis B vaccine among patients on hemodialysis. *WorldJ Hepatol* 2015;7(2):270-5.
- Perumal A, et al. Seroprevalence of hepatitis B and C in patients on hemodialysis and their antibody response to hepatitis B vaccination. *J Current Res Scientific Med* 2016;2(1):20-23.
- Amjad A, et al. Hepatitis B Vaccination status in chronic kidney disease: Experience at Pakistan institute of medical sciences. *Cureus* 2019;11(7).
- Guimarães MNC, Facincani T, Santos SDS. Hepatitis B status in hemodialysis patients. *Arquivos de Gastroenterologia* 2017;54:356-358.
- Walsh M, et al. A systematic review of the effect of nocturnal hemodialysis on blood pressure, left ventricular hypertrophy, anemia, mineral metabolism, and health-related quality of life. *Kidney Int* 2005;67(4):1500-1508.
- Erkan E, Devarajan P, Kaskel F. Role of nitric oxide, endothelin-1, and inflammatory cytokines in blood pressure regulation in hemodialysis patients. *Am J Kidney Diseases* 2002;40(1):76-81.
- Doumas MN, et al. Different effects of losartan and moxonidine on endothelial function during sympathetic activation in essential hypertension. *J Clin Hypertension* 2004;6(12):682-689.
- Levin A, Bilous R, Coresh J, Chapter 1. Definition and classification of CKD. *Kidney Int Suppl* 2013;3(1):19-62.
- Kyriazis J, Stamatidis D, Mamouna A. Intradialytic and interdialytic effects of treatment with 1.25 and 1.75 Mmol/L of calcium dialysate on arterial compliance in patients on hemodialysis. *Am J Kidney Diseases* 2000;35(6):1096-1103.
- Prasad VS, Palaniswamy C, Frishman WH. Endothelin as a clinical target in the treatment of systemic hypertension. *Cardiology in review* 2009;17(4):181-191.
- Yusuf S, et al. Telmisartan, ramipril, or both in patients at high risk for vascular events. *New England J Med* 2008;358(15):1547-1559.
- Li Z, et al. The epidemiology of systolic blood pressure and death risk in hemodialysis patients. *Am J Kid Diseases* 2006;48(4):606- 615.

Frequency of Inflammatory Bowel Disease in Patients who Underwent Colonoscopy for Lower Gastrointestinal Bleeding

Inflammatory Bowel Disease in Undergoing Colonoscopy

Dilaram Khan¹, Inayat Ullah², Mohammad Sohail¹ and Aamir Ghaffoor¹

ABSTRACT

Objective: To determine the frequency of inflammatory bowel disease in patients undergoing colonoscopy for lower gastrointestinal (GI) bleeding.

Study Design: Cross-sectional study

Place and Duration of Study: This study was conducted at the Department of Gastrointestinal Diseases, Lady Reading Hospital Peshawar from 1st June 2023 till 30th November 2023.

Methods: One hundred and twenty-one patients were included in this study. Patients of both gender and age more than 18 years with lower GI bleeding were included while patients with hemodynamic instability, patient with past history of colonoscopy for bleeding per rectum and those not willing for study were excluded. Baseline demographic information's of patients were taken and baseline complete blood count and prothrombin time were done, informed consent was taken. Every patient was stabilized hemodynamically, bowel was prepared using 2 litres of Poly Ethylene Glycol given 8 hours before the procedure and patients were kept on liquid diet from 24 hours before the colonoscopy, conscious sedation with midazolam and nalbuphine given and colonoscopy was performed using a flexible colonoscope.

Results: Age ranged from 18 to 60 years with mean age of 41.958±6.83 years. Seventy-five (62%) patients were male and 46 (38%) were female. Pain in the abdomen was the most frequently occurring complaint in addition to rectal bleeding which was present in 34 (28.9%) diarrhoea in 30(24.79%), fever in 10 (8.26%) and constipation in 4 (3.30%) patients. Inflammatory bowel disease was found in 14% of patients.

Conclusion: Lower GI bleeding represents a common problem, and inflammatory bowel disease is among the leading causes of Lower GI bleeding.

Key Words: Lower GI bleeding, Colonoscopy, Inflammatory bowel disease

Citation of article: Khan D, Inayat Ullah, Sohail M, Ghaffoor A. Frequency of Inflammatory Bowel Disease in Patients who Underwent Colonoscopy for Lower Gastrointestinal Bleeding. Med Forum 2024;35(1):83-86. doi:10.60110/medforum.350119.

INTRODUCTION

Inflammatory bowel disease (IBD) is an autoimmune disorder in which recurrent inflammatory bouts occur in gastrointestinal tract because of an abnormal immunological response to gut microflora.¹ IBD is of two types i.e. Ulcerative colitis and Crohn's disease. In ulcerative colitis diffuse inflammation of the colon mucosa occurs and most commonly affects the rectum

which is called proctitis, but it may involve the sigmoid (procto-sigmoiditis), extends beyond the sigmoid (distal ulcerative colitis), or can involve the entire colon up to the cecum (pan-colitis).² Crohn's disease causes inflammation of all the layers of the Gastrointestinal tract (GIT) called trans-mural inflammation, can involve any part of the digestive tract, mainly affecting the last part of ileum and colon. Both types of IBD are classified by extent and location. Crohn's disease also is classified by phenotype- inflammatory, structuring, or penetrating.^{2,3}

The diagnosis of IBD needs a combination of clinical history, physical findings, laboratory markers, imaging studies, and endoscopic findings and biopsies.⁴ Changes in the blood include microcytic anemia, leukocytosis, and thrombocytosis, raised ESR and CRP.⁵ Gastrointestinal bleeding is bleeding which occurs within the Gut from mouth to anus and is divided upper and lower GI bleeding on the basis of origin.⁶ Bleeding which occurs proximal to the ligament of Treitz is called upper GI bleeds, and bleeding distal to this ligament are lower GI bleeds.⁷

¹. Department of Gastroenterology / Medicine², MTI, Lady Reading Hospital, Peshawar.

Correspondence: Dr. Inayat Ullah Assistant Professor of Medicine Medical Teaching Institution Lady Reading Hospital, Peshawar.

Contact No: 03005963571

Email: drabadat78@yahoo.com

Received: December, 2023

Accepted: December, 2023

Printed: January, 2024

This classification into these two is important as helps in the further assessment and ultimate treatment of the patient.⁷ Hreinsson et al⁸ have shown in a study that frequency of IBD was 13% in patients who underwent endoscopy for lower GI bleeding. In Pakistan, colonoscopy is done for diagnosis of lower GI bleeding but very limited studies are available in our local set up regarding lower GI bleeding. So, this study was designed to study the frequency of IBD in patients undergoing colonoscopy for lower GI bleeding in our local set up.

METHODS

This cross-sectional study consisting of 121 patients was done in the Gastroenterology unit of MTI- Lady reading hospital, Peshawar from 1st June 2023 to 30th November 2023 using non-probability consecutive sampling technique. Approval was given by ethical committee of the hospital. Patients of both genders having age more than 18 years and complaining of lower GI bleeding were enrolled in this study while Patients with hemodynamic instability, and those not willing for the study were excluded. Baseline demographic information's of patients (age, gender and duration of complaints) were taken and baseline complete blood count and prothrombin time were done, and informed consent was taken from patients. Patients were hemodynamically stabilized, preparation of the bowel done using 2 litres of Poly Ethylene Glycol given 8 hours prior to the procedure and patients were kept on liquid for 24 hours before the colonoscopy whenever possible. Conscious sedation given with midazolam and nalbuphine and colonoscopy was performed. In case of acute emergency, only sigmoidoscopy was done after giving two enemas 30 minutes before the procedure. During the colonoscopy, the patient was laid on left lateral position and standard flexible colonoscope was used. Colonoscopy findings were noted, IBD was confirmed colonoscopically and minimum of four biopsy specimen were taken from each patient for histological confirmation. Data regarding inflammatory bowel disease was noted. Data was analyzed by using SPSS-23. Post stratification Chi square test was applied, $p \leq 0.05$ was considered statistically significant.

RESULTS

There were 75 (62%) males and 46 (38%) females with a male to female ratio of 1.60: and mean age was 41.958 ± 6.83 years, the majority of the patients were in the age range of 40-60 (Table 1). In addition to bleeding per rectum, 34 (28.09%) patients were complaining of pain in abdomen, 30 (24.79%) diarrhea, 10 (8.26%) fever and 4 (3.30%) of constipation (Table 2). Inflammatory bowel disease was noted in 14% patients, where all of 17 (14) patients were having ulcerative colitis while none of them was diagnosed with Crohn's disease (Table 3). Stratification of

inflammatory bowel disease with respect to age, gender and duration of complaints are shown in Tables 4-5 respectively.

Table No.1: Age wise distribution of patients (n=121)

Age (Years)	No.	%
18-40	50	41
40-60	71	58
Above 60	-	-

Table No.2: Other symptoms present along with rectal bleeding (n=121)

Symptoms	No.	%
Pain abdomen	34	28.09
Diarrhea	30	24.79
Fever	10	8.26
Constipation	4	3.30

Table No.3: Frequency of inflammatory bowel disease (n=121)

Inflammatory bowel disease	No.	%
Yes	17	14.0
No	104	86.0

Table No.4: Stratification of inflammatory bowel disease with respect to age

Age (years)	Inflammatory bowel disease		P value
	Yes	No	
18-40	5 (11.9%)	37 (88.1%)	0.621
>40	12 (15.2%)	67 (84.8%)	

Table No.5: Stratification of inflammatory bowel disease with respect to gender

Gender	Inflammatory bowel disease		P value
	Yes	No	
Male	7 (9.3%)	68 (90.7%)	0.057
Female	10 (21.7%)	36 (78.3%)	

DISCUSSION

Lower GI bleeding is a frequent cause of referrals to Gastroenterology Centers⁹ all over the globe, and same is the clinical situations in our country and local set up as well. Male patients predominated in our study, 62% were male, and the rest were female. Our study results are comparable to the study done by Deeb et al¹⁰ in Egypt, where 68.0% of the patients with lower GI bleed were male. Similarly, a study carried out in India by Bhadauria et al¹¹ showed a male to female ratio of 2.16:1 in patients who presented with lower gastrointestinal bleeding.

The mean age of patients in our study was 41.958 ± 6.83 years. The same were the results of the study done by Mandhan et al¹² where majority of the patients were in the young age. In this study, abdominal pain was the commonest presenting complaint reported in 34 (28.09%), diarrhea in 30 (24.79%), fever 10(8.26%) and constipation in 4 (3.3%) patients. These results of our study were very same to a study done in the Egypt¹⁰

where pain in the abdomen and loose motions were commonest complaints. Similarly, a study done by Arvola et al¹³ also noted that anemia, pain in the abdomen and diarrhea were the most common presentations in children who presented with Lower gastrointestinal bleeding. In a study done by an author loose motions, vomiting and pain in the abdomen were the most common symptoms among patients presenting with lower gastrointestinal bleeding. In a study done by Zahmatkeshan et al¹⁴ in Iran showed that pyrexia, pain abdomen, and loose stools were the commonest presentations accompanying bloody stool. We found 64.2% of the cases were having anemia at the time of admission. These findings are compatible to another study where 61% of the cases with Lower GI bleeding had pallor.¹⁰ Pallor has been noted to be a common finding among children having chronic blood loss.

In this study, colonoscopy findings were suggestive of inflammatory bowel disease in 14% of patients. The same were the results in the study done by Hreinsson et al⁸ which showed that frequency of inflammatory bowel disease was 13% in patients who did endoscopic examination of the colon for lower GI bleeding. In a study from Egypt¹⁰ same findings were noted with polyp being the commonest colonoscopy finding present in 44% whereas results from another study by Clarke et al⁹ showed that polyps were present only in 10% of the patients. Greater frequencies of polyps (75%) on colonoscopic examination among patients having Lower GI bleeding was shown by Mandhan.¹² Another author noted that polyps were present in 53% while studies from many other regions of the globe identified polyps to be the most common cause among children with Lower GI bleeding. Deeb et al¹⁰ in Egypt also showed that juvenile polyps to be the most frequently occurring finding which were hamartomatous and responsible for upto 90% of all kinds of polyps noted among children.¹⁵ The diagnosis of inflammatory bowel disease depends mainly on the combination of clinical presentations, laboratory markers, radiological studies, colonoscopy, and histopathological examination. However, sometimes the colonoscopic findings are non-specific and usually occur because of some other etiologies. In addition to differentiate between the two types of IBD and for knowing the extent of the disease, other etiologies of colon inflammation needs to be ruled out. This is of special importance as the treatment for ulcerative colitis or Crohn's disease may worsen other conditions, particularly colonic infection. Infectious agents causing colonic inflammations are very similar inflammatory bowel disease on colonoscopy. Common infections agents like *Clostridium difficile* (CD) and *Escherichia coli* (E coli) must be win now out before colonoscopy. *Yersinia* spp. Can cause abdominal pain in the lower quadrant and pyrexia, where imaging is showing ileitis and is usually similar to acute inflammation of the appendix. *Salmonella*, *Actinomyces*, and *E. coli* infections can cause enteritis and particularly ileitis which are similar to IBD presentations.¹⁶ Intestinal TB

can also cause ulcer, and stricture formation in the terminal ileum and ileocecal valve.

Cytomegalovirus (CMV) can cause ulcerations of the GI tract where ulcers are usually "punched-out" in appearance but biopsies are necessary to differentiate between the two. However, many individuals with inflammatory bowel disease will have CMV infection at the same time, so colonoscopy inspection is mandatory to rule out concomitant Cytomegalovirus infection which is causing bowel inflammation, but sometimes it can be very challenging to establish whether Cytomegalovirus is just a passerby or an active participant in inflammation in these patients.¹⁷

Though it is very rare but vasculitis can also cause bowel inflammation, especially the small Gut. SLE, polyarthritis nodosa, Henoch-Schönlein purpura (HSP), and Behçet's disease may all cause colonic inflammation just like IBD. The gastrointestinal tract is affected by Polyarteritis nodosa in up to 65% of patients and can cause symptoms of bowel ischemia.¹⁸ Behçet's disease usually cause ulcers in the small and large intestines with normal intervening mucosa and these are usually confused with Crohn disease. However, the ulcerations in Behçet's disease are usually less in number, are larger, deeper, and rounder as compared to those which occur in IBD.¹⁹

Ischemia can cause erythema, edema, erosion and ulcerations which are similar to those of inflammatory disease. Ischemic colitis usually cause dusky necrotic colitis which occur in segments, having a demarcation between diseased and normal colon and usually affect the left colon. A detailed history and accuracy of symptoms can differentiate between IBD and ischemic colitis.^{20,21} diverticulosis causing segmental colitis is also very difficult to differentiate from inflammatory bowel disease. Segmental colitis of diverticulosis most commonly affects the left sided colon, especially the sigmoid colon. While rectum and the rest of the colon are spared most of the time. Edema, erythema, erosions, and ulcers, often with sparing of the diverticular orifices are the main endoscopic findings.²² Since the colonoscopic and biopsy features overlap with the inflammatory bowel disease, the diagnosis is usually challenging, but Segmental colitis associated diverticulosis is mainly found in old age patients.²³ similarly NSAIDs can cause bowel inflammation and proper history and examination is needed to differentiate it from IBD.

CONCLUSION

Lower GI bleeding is a common problem with a vast differential diagnosis. Inflammatory bowel disease is among the leading causes of Lower GI bleeding. So thorough history, physical exam, and utilization of endoscopic and radiographic adjunct are crucial in identification of the etiology of the bleeding. Endoscopic examination of the colon is a beneficial and safe procedure in patients who present with lower GI bleeding.

Author's Contribution:

Concept & Design of Study: Dilaram Khan
 Drafting: Inayat Ullah,
 Mohammad Sohail
 Data Analysis: Mohammad Sohail,
 Aamir Ghaffoor
 Revisiting Critically: Dilaram Khan, Inayat
 Ullah
 Final Approval of version: Dilaram Khan

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.1043/LRH/MTI dated 01.06.2023

REFERENCES

- Maaser C, Sturm A, Vavricka SR, Kucharzik T, Fiorino G, Annese V, et al. ECCO-ESGAR Guideline for Diagnostic Assessment in IBD Part 1: initial diagnosis, monitoring of known IBD, detection of complications. *J Crohns Colitis* 2019;13(2):144-64.
- Dmochowska N, Wardill HR, Hughes PA. Advances in imaging specific mediators of inflammatory bowel disease. *Int J Mol Sci* 2018;19(9):2471.
- Colombel JF, Shin A, Gibson PR. AGA clinical practice update on functional gastrointestinal symptoms in patients with inflammatory bowel disease: expert review. *Clin Gastroenterol Hepatol* 2019;17(3):380-90.
- Lee JS, Kim ES, Moon W. Chronological review of endoscopic indices in inflammatory bowel disease. *Clin Endosc* 2019;52(2):129-36.
- Dai C, Jiang M, Sun MJ. Fecal markers in the management of inflammatory bowel disease. *Postgrad Med* 2018;130(7):597-606.
- Kim ER, Chang DK. Management of complications of colorectal submucosal dissection. *Clin Endosc* 2019; 52(2):114-9.
- Shafqet MA, Tonthat A, Esparragoza P, Toro B, Ehrlich AC, Friedenbergl FK. Recent use of NSAID and NOAC medications are associated with a positive CT arteriogram. *Abdom Radiol* 2019; 44(7):2632-8.
- Hreinsson JP, Egisdottir S, Bjornsson ES. Acute lower gastrointestinal bleeding: a population-based five-year follow-up study. *United Eur Gastroenterol J* 2019; 7(10):1330-6.
- Clarke G, Robb A, Sugarman I, McCallion WA. Investigating painless rectal bleeding – is there scope for improvement? *J Pediatr Surg* 2005;40:1920–1922.
- Deeb MM, El-Zayat RS, El-Khair HA. Colonoscopic findings in children with lower gastrointestinal bleeding. *Menoufia Med J* 2016;29:247-521.
- Bhadoria N, Dubey SR, Mittal P, Arya AK, Singh RP. Clinicoetiological pattern of lower gastrointestinal bleeding in children (5-18 years age group) at a tertiary care center in central India. *Indian J Child Health* 2016;3(4):290-92.
- Mandhan P. Sigmoidoscopy in children with chronic lower gastrointestinal bleeding. *J Paediatr Child Health* 2004;40:365-68.
- Arvola T, Ruuska T, Keränen J, Hyöty H, Salminen S, Isolauri E. Rectal bleeding in infancy: clinical, allergological, and microbiological examination. *Pediatr* 2006;117:760- 68.
- Zahmatkeshan M, Fallahzadeh E, Najib K, Geramizadeh B, Haghghat M, Imanieh MH. Etiology of Lower Gastrointestinal Bleeding in Children: A Single Center Experience from Southern Iran. *Middle East J Dig Dis* 2012;4: 216-23.
- Padilla BE, Moses W. Lower Gastrointestinal Bleeding & Intussusception. *Surg Clin North Am* 2017;97(1):173-88.
- Dilauro S, Crum-Cianflone NF. Ileitis: When it is not Crohn's disease. *Curr Gastroenterol Reports* 2010; 12(4):249-58.
- Garrido E, Carrera E, Manzano R, Lopez-Sanroman A. Clinical significance of cytomegalovirus infection in patients with inflammatory bowel disease. *World J Gastroenterol* 2013;19(1):17-25.
- Ebert EC, Hagspiel KD, Nagar M, Schlesinger N. Gastrointestinal involvement in polyarteritis nodosa. *Clin Gastroenterol Hepatol* 2008;6(9):960-66.
- Kim DH, Cheon JH. Intestinal Behçet's disease: A true inflammatory bowel disease or merely an intestinal complication of systemic vasculitis? *Yonsei Med J* 2016;57(1):22-32.
- Zou X, Cao J, Yao Y, Liu W, Chen L. Endoscopic findings and clinicopathologic characteristics of ischemic colitis: A report of 85 cases. *Digestive Dis Sci* 2009;54(9):2009-15.
- Lozano-Maya M, Ponferrada-Díaz A, González-Asanza C, et al. Usefulness of colonoscopy in ischemic colitis. *Revista Española de Enfermedades Digestivas* 2010;102(102):478-83.
- Tursi A, Elisei W, Brandimarte G, et al. The endoscopic spectrum of segmental colitis associated with diverticulosis. *Colorectal Dis* 2010;12(5):464-70.
- Cassieri C, Brandimarte G, Elisei W, et al. How to differentiate segmental colitis associated with diverticulosis and inflammatory bowel diseases. *J Clin Gastroenterol* 2016;50(Suppl 1):S36-8.

Developmental Profile as a Predictor of Behavior Phenotype in Down Syndrome Children

Predictor of Behavior phenotype in Down Syndrome Children

M Bilal Abid¹, Syeda Wajeeda Zahra³, Muneeba Kamran¹, Sultan Badar Munir², Ehsan Ullah¹ and Munawar Ghous¹

ABSTRACT

Objective: To study the Emerging Down Syndrome Behavioral Phenotype from infancy to Early Childhood and to see Developmental Profile as a Predictor of this Behavior phenotype in Down Syndrome Children.

Study Design: Cross-sectional study

Place and Duration of Study: This study was conducted at the Department of Development and Behavioral pediatrics, University of Child Health science, Children's hospital Lahore and duration of study was 6 months from 1st January 2023 to 30th June 2023.

Methods: It was a Cross-sectional study. Data was collected from Department of Development and Behavioral pediatrics, University of Child Health science, Children's hospital Lahore and duration of study was 6 months. A sample of 42 Children having age 1-5 years was collected through purposive sampling technique.

Results: In breakup of PEEP and SDQ tools, Cognition age of 9 (19%) patients was profound, 23 (48%) severe, 9 (19%) moderate and 7(15%) was mild delay. Socialization age of 4 (8%) patients was profound, 11 (23%) severe, 13 (27%) moderate, 11(23%) mild and 9 (19%) patients were age appropriate. Self-help age of 7(15%) patients was profound, 12 (25%) severe, 13 (27%) moderate, 10 (21%) was mild delayed and 6 (13%) patients were patients were age appropriate. Motor age of 8(17%) patients was profound, 13 (27%) severe, 15 (31%) moderate, 8 (17%) was mild delay and 4 (8%) patients were age appropriate. Expressive Llanguage age of 19 (40%) patients was profound, 22 (46%) severe, 5 (10%) moderate 1(2%) was mild delay and 1 (2%) patients were age appropriate.

Conclusion: Specific Behavioral Phenotype (BF) in children with Down Syndrome (DS) is consistent with relative strength in sociability, non-verbal abilities, receptive language, implicit memory skills and visuo spatial processing and relative weakness in gross & fine motor skills, verbal communication, visuo motor, cognitive functioning and motor planning.

Key Words: Down syndrome (DS) Behavior Phenotype (BP), Development Profile (DP), Portage Early Education Plan (PEEP)

Citation of article: Abid MB, Zahra SW, Kamran M, Munir SB, Ehsan Ullah, Ghous M. Developmental Profile as a Predictor of Behavior Phenotype in Down Syndrome Children. Med Forum 2024;35(1):87-91. doi:10.60110/medforum.350120.

INTRODUCTION

Down syndrome (DS), trisomy 21, is the most common genetic disorder, with an estimated incidence of 1 in 700 live births¹. Out of all genetic disorders, a lot of research has been done on Down syndrome. Development is typical in infancy, slows down in next two years amid delayed rate of brain myelination.

¹. Department of Development and Behavioral Pediatrics / Paediatric ENT², UCHS, Children's Hospital Lahore.

³. Department of Psychology, Ganga Ram Hospital, Lahore.

Correspondence: Munawar Ghous, Statistician, UCHS, Children's Hospital Lahore.

Contact No: 03124236642

Email: munawar.ghous@gmail.com

Received: December, 2023

Accepted: December, 2023

Printed: January, 2024

The Down syndrome (DS) behavior phenotype has been described as having relative strengths in non-verbal receptive skills and sociability & relative weakness in cognition expressive language and motor planning². The most constant and typical features of DS are intellectual disability and craniofacial dysmorphism³, with a variety of organ involvement and systemic diseases⁴. By definition behavioral phenotype refers to observable characteristics that occur more often in individuals with a specific genetic syndrome than individuals without that syndrome⁵. Characters that are not observable are called endo-phenotype and include thoughts, emotions and motivational states. Behavior phenotype in children with DS has typical characteristics and is explained in domains of his/her cognition/intellect, socialization, speech and language communication, self-help and motor skills which are collectively called as developmental profile⁵.

In cognition domain of development, children with DS have mild to severe delay with the profile of relative strength and weakness. They tend to have difficulties

more in understanding & following commands impaired memory, learning & self-care but they are good in non-verbal abilities and visually stored memory⁶.

As regards to Speech and Language domain of development children with DS have better receptive than expressive skills in the verbal domain⁷. Receptive vocabulary can be considered as a relative strength. The understood more than they can speak. As for the expressive domain, children with DS have been found weak in phonology, grammar and syntax, while their intentional use of communication and gestures, and their social use of communication generally seem to be in line with their mental age⁸. A pattern of strengths and weaknesses has been observed in the nonverbal domain as well. In terms of social development children with DS are relatively strong and often described as charming affectionate, outgoing, cheerful, happy and sociable. Motor functioning demonstrates specific motor impairments in a number of fine and gross motor tasks (i.e., balance, posture, strength, and flexibility), as well as motor planning (i.e., praxis), although CA-level performance has been observed in specific skills including, running speed, agility, and visual-motor control. Conductive Hearing impairment negatively impacts language development⁹.

This research will help in establishing the developmental profile in early childhood can be used as a predictor of Behavior Phenotype (Strengths and weakness) in children with Down syndrome.

METHODS

It was a Cross-sectional study. Data was collected from Department of Development and Behavioral pediatrics, University of Child Health science, Children’s hospital Lahore and duration of study was 6 months. A sample of 42 Children having age 1-5 years was collected through purposive sampling technique. PEEP (Portage Early Educational Plan) Guide was administered to determine developmental Delay in 5 domains of development (Cognition, Self Help, Socialization, Motor and Speech). Strength and Difficulties Questionnaire (SDQ) was incorporated to exclude children with DS having behaviors related to pervasive development disorders. Demographic Questionnaire including Age, Gender, Family Size, Education, income etc. was also administered.

RESULTS

Data was analyzed using SPSS 25.0. Demographic and socio-economic data of our research shows that, 36 (75%) patients were male and 12 (25%) patients were female. 13 (27.1%) patients were belonging to nuclear family system and 35 (72.9%) patients were belonging to joint family system. 30 (62.5%) parents of patients have low socio-economic status, 16 (33.3%) parents of patients have middle socio-economic status and 2

(4.2%) parents of patients have high socio-economic status. 40% father and 44% mothers of patients were un-educated.

Figure 1 shows that, in breakup of PEEP and SDQ tools, Cognition age of 9 (19%) patients was profound, 23 (48%) severe, 9 (19%) moderate and 7(15%) was mild delay. Socialization age of 4 (8%) patients was profound, 11 (23%) severe, 13 (27%) moderate, 11(23%) mild and 9 (19%) patients were age appropriate. Self-help age of 7(15%) patients was profound, 12 (25%) severe, 13 (27%) moderate, 10 (21%) was mild delayed and 6 (13%) patients were age appropriate. Motor age of 8(17%) patients was profound, 13 (27%) severe, 15 (31%) moderate, 8 (17%) was mild delay and 4 (8%) patients were age appropriate. Expressive Language age of 19 (40%) patients was profound, 22 (46%) severe, 5 (10%) moderate 1(2%) was mild delay and 1 (2%) patients were age appropriate.


Table 1: this table shows, throwing 34 (71%), Teasing 5 (10%), Screaming 32 (67%), Selfdirected 16 (33%), and Directed toward others 34 (71%). Table 2: shows that average age of patients was 35.79 months, Cognition age was 16.61 months, Socialzation age was 27.43 months, Fine Motor and Self-Help age was 21.96 months, Gross Motor age was 21.13 months, Receptive language age was 28.18 months and Expressive language age of Down syndrom (DS) patients was 12.73 months. Table 3: shows that there was a significant difference between chronoligcal age and socialization age, choronoligcal age and self-help age, choronoligcal age and Gross Motor age, choronoligcal age and Receptive Language age, choronoligcal age and Expressive Language age of DS patients.

Table No.1 : Seconderly Behaviuoral problems

Behaviors of individuals	Yes	No
Throwing	34 (71%)	14 (29%)
Teasing	5 (10%)	43 (90%)
Screaming	32 (67%)	16 (33%)
Self-directed	16 (33%)	32 (67%)
Directed toward others	34 (71%)	14 (29%)

Table No.2: Paired sample statistics

Age	Mean	Std. Deviation
Chronological age of patients	35.79	15.12
Cognition age	16.61	11.19
Socialization age	27.43	11.97
Fine Motor and Self-Help age	21.96	13.31
Gross Motor age	21.13	13.15
Receptive Language age	28.18	9.13
Expressive Language age	12.73	10.20



Figurer No.1: Cognition age, socialization age, self-help age, Motor age and language age of patients

Table No.3: Output of Paired sample t-test

Variables	Paired Differences		T	p-value
	Mean	SD		
Chronological age - Cognition age	19.18	10.958	12.124	<0.005
Chronological age -Socialization age	8.37	10.623	5.460	<0.005
Chronological age - Fine Motor and Self-Help age	13.83	11.360	8.435	<0.005
Chronological age – Gross Motor age	14.66	9.664	10.509	<0.005
Chronological age – Receptive Language age	23.07	9.160	17.447	<0.005
Chronological age - Expressive Language age	23.07	11.810	13.532	<0.005

DISCUSSION

Specific Behavioral Phenotype (BF) consistent in children with Down Syndrome (DS) presented relative strength in sociability, non-verbal abilities, receptive language, implicit memory skills and visuo spatial processing and relative weakness in gross & fine motor skills, visuo motor, cognitive functioning, expressive language, communication and motor planning. This BF is clearly related to the developmental profile (DP) of the child and early development of typical behavioral characters (BC). We measured the DP & BC through PEEP and SDQ tool was incorporated to exclude children with pervasive development disorders and found PEEP as predictor of typical emerging BF in DS children.

Previous studies have reported this specific behavioral phenotype in children above 5 years¹⁰. This study was shaped to see this BF through developmental & behavioral assessments of children with DS before 6 years. Description and results of the research is viewing behavioral phenotypes from a developmental perspective. It seemed that this particular phenotype appears in early years of development and then emerge slowly before five years of age.

The Expressive Language (EL) was assessed through child’s spontaneous answers, spoken responses to questions and multi word sentences. The expressive language words were all centrally processed, requiring the child to respond to an auditory input. Expression

was in line with comprehension and auditory processing and mechanical motor skills. Expressive Language (EL) skills were recorded on parents information and direct conversation with the child and noted according to number of intelligible words, 2 or 3 words speech and sentence formation these findings are agreement with previous studies^{11,12}.

A specific cognitive profile was observed in children with (DS) showing difficulties in intelligible and comprehensive language and spoken memory challenges, and relatively stronger non-verbal abilities and visual memory skills. They are better in receptive than expressive skills in the verbal domains, meaning thereby that these children understood more words than they are able to speak. Receptive Vocabulary (RV) is relatively better, but the depth and breadth of their RV is weak. DS children have been found weak on phonology, grammar and syntax but their social cues and gestures are generally according to their mental age. Low cognition has been ascribed due to deficits in verbal processing, large differences in expressive and receptive language domains and smaller size of intelligible spoken words. Our study showed cognitive age as 16.61 months while other study¹³ showed it as 18.22 months.

The communication domain involved receptive and expressive language. In communication competence, non- verbal joint attention and gestural language are better whereas non-verbal requesting behavior showed deficits in children with DS. In receptive language

child's verbal input is determined as the small item questions they can understand better like pointing and simple commands but complex items involving multi tasks in one command were difficult to perceive. In our study Receptive and Expressive language of patients were 28.18 and 12.73 months respectively while previous study¹³ showed that Receptive and Expressive language as 22.11 and 19.33 months respectively.

The Gross Motor (GM) functions were assessed as at what age child developed head control, turning sides, and prone to spine, sitting, crawling, standing in a manner to know progress in cephalocaudal and proximodistal motor development. In preschool children GM skills like running, hopping, jumping, climbing up & down stairs and tricycle riding. ("Pedals tricycle or other three-wheeled vehicle for at least six feet") were noted as their age of development. Our study showed Gross Motor age as 21.13 months while older study¹³ showed it as 18.22 months.

The Fine Motor (FM) and Self Help skills involved bilateral and unilateral hand functioning. Bilateral items included closing zip, opening book and turning pages, buttoning/unbuttoning, lacing shoes, sharpening pencils, wearing socks & shirts, use of cloth clips, folding, and cutting etc. Unilateral items included eating with sticks, use of spoon and forks, holding feeder, bridging with blocks, marking tower of blocks, use of peg board, etc. Our study showed Fine Motor age as 21.96 months while older study¹³ showed it as 20.83 months

The Socialization Domain assessed through functioning in peer relationships (PR), play and entertainment hours, and adapting skills. For toddlers, PR items involve like "laughs or smiles appropriately in response to positive statements. For preschool children dimensions like "participates in at least one game or activity with others", and coping skills involve items like "says 'please' when asking for something". Age equivalent scores for the Down syndrome group in this study on Play and Leisure Time socialization averaged 27.43 months, in contrast to the previous study¹³ the (PR) area domain showed average age equivalent scores of 20.17 months. Positive emotions signals & smile frequencies were also found high.

In this context two marked differences were noted in individuals with Down syndrome. It was small at early developmental ages as regards to difference between expressive and receptive language, which averaged only 2.5 months. Whereas, it was large with an average of 20 months in older children with Down syndrome. That small dissociations early in development can result in increasingly larger differences over time is consistent with dynamic systems theory in those small starting state differences can evolve considerably larger as development becomes increasingly complex and differentiated^{14,15}.

The Visual Reception measures were taken from visually stored memory. Children were exposed to various objects in different forms and shapes which involved oculomotor nerves and central visual pathways in localizing single and multiples points on surface through visual tracking¹⁶.

CONCLUSION

In the light of new understanding of development in genetic syndromes and as a part of larger movement towards studying BF particularly in children with DS, it has become possible now to focus on weaker areas of BF like expressive language and motor skills before they become areas of pronounced weakness. Areas of relative strength may be taken as "gateway in" to polish areas of weakness to prevent future delays by early identification of areas of strength & weakness through developmental profile in toddler years taking as potential windows of opportunity to address weaker areas of BF is in toddler age group before they become pronounce in preschool years.

Visuospatial aspects of visual recognition memory, visual motor integration and visual imitation as areas of strength in children with DS are useful sub domains to enhance cognition level in these children.

Author's Contribution:

Concept & Design of Study:	M Bilal Abid
Drafting:	Syeda Wajeaha Zahra, Muneeba Kamran
Data Analysis:	Sultan Badar Munir, Ehsan Ullah, Munawar Ghous
Revisiting Critically:	M Bilal Abid, Syeda Wajeaha Zahra
Final Approval of version:	M Bilal Abid

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.752/CH-UCHS dated 27.12.2022

REFERENCES

1. Onnivello S, Pulina F, Locatelli C, Marcolin C, Ramacieri G, Antonaros F, et al. Cognitive profiles in children and adolescents with Down syndrome. *Sci Rep* 2022;12(1):1936.
2. Hickey F, Hickey E, Summar KL. Medical update for children with Down syndrome for the pediatrician and family practitioner. *Adv Pediatr* 2012;59(1):137-57.
3. Waite, Jane, Heald, Mary, Wilde, Lucy, et al. The importance of understanding the behavioural phenotypes of genetic syndromes associated with

- intellectual disability. *Paediatrics and Child Health* 2014;24: 10.1016/j.paed.2014.05.002.
4. Abbeduto L, Warren SF, Conners FA. Language development in Down syndrome: from the prelinguistic period to the acquisition of literacy. *Ment Retard Dev Disabil Res Rev* 2007;13(3): 247-61.
 5. Asif A, K Ansari MY, Hashem A, Tabassum B, Abd Allah EF, Ahmad A. Proteome Profiling of the Mutagen-Induced Morphological and Yield Macro-Mutant Lines of *Nigella sativa* L. *Plants (Basel)* 2019;8(9):321.
 6. Laws G, Briscoe J, Ang SY, Brown H, Hermena E, Kapikian A. Receptive vocabulary and semantic knowledge in children with SLI and children with Down syndrome. *Child Neuropsychol* 2015;21(4):490-508.
 7. Deckers SRJM, Van Zaalén Y, Van Balkom H, Verhoeven L. Predictors of receptive and expressive vocabulary development in children with Down syndrome. *Int J Speech Lang Pathol* 2019;21(1):10-22.
 8. Lanfranchi S, Jerman O, Vianello R. Working memory and cognitive skills in individuals with Down syndrome. *Child Neuropsychol* 2009;15(4): 397-416.
 9. Fidler, Deborah, Philofsky, Amy. The Down syndrome behavioural phenotype: Taking a developmental approach. *Down Syndrome: Research and Practice* 2009;12. 10.3104/reviews/2069.
 10. Bull MJ. Down Syndrome. *N Engl J Med* 2020;382(24):2344-2352.
 11. Meneghetti C, Toffalini E, Carretti B, Lanfranchi S. Mental rotation ability and everyday-life spatial activities in individuals with Down syndrome. *Res Dev Disabil* 2018;72:33-41.
 12. Ukoumunne O, Wake, Melissa, Carlin, Bavin J, Edith, et al. Profiles of language development in pre-school children: A longitudinal latent class analysis of data from the Early Language in Victoria Study. *Child: care, health and development* 2011;38:341-9.
 13. Fidler DJ, Hepburn S, Rogers S. Early learning and adaptive behaviour in toddlers with Down syndrome: evidence for an emerging behavioural phenotype? *Downs Syndr Res Pract* 2006;9(3): 37-44.
 14. Pulsifer M. Down Syndrome: Neuropsychological Phenotype across the Lifespan. *Brain Sci* 2021;11(11):1380.
 15. Söylemez F. Phenotypes Associated with Down Syndrome and Causative Genes. *Down Syndrome and Other Chromosome Abnormalities*. Intech Open; 2022. Available from: <http://dx.doi.org/10.5772/intechopen.96290>
 16. Zhang L, Fan H. Visual object tracking: Progress, challenge, and future. *Innovation (Camb)* 2023;4(2):100402.

Chest Pain Management Using Prehospital Point-of-Care Troponin and Paramedic Risk Assessment

Feras Almarshad¹ and Ghulam Mustafa²

ABSTRACT

Objective: To evaluate the prehospital point-of-care troponin testing and paramedic risk stratification within existing chest pain care pathways reveals promising reliability and validity, suggesting their potential as valuable tools in enhancing early cardiac assessment and improving patient outcomes.

Study Design: Prospective study

Place and Duration of Study: This study was conducted at the Department of Medicine, Nishtar Medical University, Multan, from February 2021 to January 2022.

Methods: Study included 400 consecutive patients experiencing acute chest pain categorized as emergencies. All patients were transported to the hospital via ambulance for subsequent medical care. Positive TnT test prehospital and at the time of hospital admission were compared. Main variables of study were values of troponin test in myocardial infarction associated chest pain patients at hospital and during emergency travelling and previous history coronary artery disease, myocardial infarction and risk factors like diabetes, smoking and hypertension.

Results: Myocardial infarction was positive in 32.0% in chest pain patients. Prehospital troponin was positive in 33 (8.3%) patients. Troponin test was positive in 53.8% patients at admission. The sensitivity for prehospital troponin to myocardial infarction was 18.0% and the sensitivity to myocardial infarction for in-hospital troponin was 74.2% with specificity 96.3% and 55.9%, respectively. ($p < 0.001$).

Conclusion: In regions where patient transport times are brief, the point-of-care rapid Troponin T (TnT) test reveals only a minority of individuals experiencing chest associated, prehospital TnT test positive outcome emerges as an objective indicator for a more adverse prognosis in those with suspected heart attacks.

Key Words: Point of care, Chest Pain, Myocardial infarction, Troponin test, Prehospital

Citation of article: Almarshad F, Mustafa G. Chest Pain Management Using Prehospital Point-of-Care Troponin and Paramedic Risk Assessment. *Med Forum* 2024;35(1):92-95. doi:10.60110/medforum.350121.

INTRODUCTION

Acute chest pain represents 10% of ambulance attendances and is linked to significant healthcare costs and resource utilization¹. Existing guidelines advise transporting most patients to emergency departments for further assessment due to the presence of chest pain in many serious diagnoses; however, approximately 50% of these patients are ultimately discharged without a specific diagnosis². Recent studies indicate that risk stratification in coronary artery disease can be made safely by paramedic staff by testing point-of-care troponin level, leading to a reduction in emergency department length of stay^{3,4}.

¹. Department of Internal Medicine / Pediatrics Medicine², College of Medicine, Shaqra University, Shaqra, Saudi Arabia.

Correspondence: Dr. Ghulam Mustafa, Associate Professor, Pediatrics Medicine Department, College of Medicine Shaqra, Shaqra University, Saudi Arabia.
Contact No: 0300 8635452
Email: ghulammustafa@su.edu.sa

Received: July, 2023

Accepted: September, 2023

Printed: January, 2024

While health outcome studies evaluating this strategy are still emerging, prehospital risk stratification and point-of-care troponin testing may be non inferior to existing care processes in terms of early major adverse cardiac events⁵. Studies conducted to date have primarily utilized conventional troponin assays, but the emergence of point-of-care, high-sensitivity troponin (hsTn) assays may become more prevalent in the near future⁶.

However, the potential widespread availability of hsTn assays poses challenges due to significant infrastructure costs linked to outfitting ambulances with point-of-care testing devices and TnT cartridges^{7,8}. Additionally, uncertainties persist regarding whether implementing a prehospital risk stratification and point-of-care troponin model would ultimately lead to net cost savings, considering the associated paramedic training expenses^{9,10}.

The present study aimed to investigate paramedic risk assessment, in conjunction with point-of-care troponin testing that may help optimize resource allocation by directing patients to the most appropriate level of care.

METHODS

The Prospective study was conducted at department of Medicine, Nishtar Medical University, Multan, from

February 2021 to January 2022. Total 400 consecutive patients experiencing acute chest pain categorized as emergencies were included. All patients were transported to the hospital via ambulance for subsequent medical care. The protocol approved by the hospital Board. Study protocol involved the collection of 5 mL of blood for the rapid Troponin T (TnT) test after obtaining verbal informed consent from the patient. The blood collection took place either in the patient's home or in the ambulance. Subsequently, a second TnT test was conducted upon the patient's arrival at the hospital. All patients in the study were administered 500 mg of intravenous aspirin and received nitroglycerin sublingually during ambulance transport. Upon hospital admission, a 12-lead electrocardiogram was promptly conducted, and concurrent assessments of creatine kinase were carried out. Subsequent serial measurements were left to the discretion of the attending physician, who was informed of the troponin T (TnT) test results.

Positive TnT test prehospital and at the time of hospital admission were compared. In patients of chest pain MI incidence was noted and mentioned with positive and negative TnT prehospital test were recorded.

MI diagnosis was based on a significant rise in creatine kinase activity (more than twice the upper limit of normal), along with a creatine kinase MB increase of over 6%, concurrent ST elevations on electrocardiogram or as per World Health Organization criteria for typical clinical findings. For diagnosis of Coronary artery disease coronary angiography was performed. Presence of ST elevation above 1mm and 2 mm on limb lead and precordial lead was labeled as myocardial ischemia.

A device for whole blood rapid assay, utilizing Boehringer Mannheim's TnT (troponin T) test was applied for measurement of TnT levels. After heparinization of 150 ml whole blood, plasma and cellular fraction was separated. Plasma containing TnT, is diffused to the detection zone. Cardio specific gold-labeled antibodies and biotinylated antibodies are involved. These antibodies are specific to different epitopes (distinct regions) of Troponin T. A red line

appears in the reading zone within 20 minutes. This red line serves as an indicator of the presence of serum TnT in the sample.

The data were presented as median or mean (SD) as appropriate. Statistical analyses employed Student's t-test and Fisher test with a significance threshold of $P < 0.05$. Sensitivity and specificity were calculated as percentages of true positives and true negatives, respectively, relative to total relevant cases.

RESULTS

Overall, 400 patients were included in this study with mean age 62.71 ± 6.01 years. There were 209 (52.3%) males and 191 (47.7%) females. The distribution of hypertension, CAD, smoking, previous MI, dyslipidemia, diabetes mellitus and family history were shown in table 1.

Myocardial infarction was positive in 128 (32.0%) chest pain patients. Prehospital troponin was positive in 33 (8.3%) patients. Troponin test was positive in 215 (53.8%) patients at admission. The sensitivity for prehospital troponin to myocardial infarction was (18.0%) and the sensitivity to myocardial infarction for in-hospital troponin was (74.2%) with specificity (96.3%) and (55.9%), respectively. ($p < 0.001$). (Table. 2).

Table No.1: Demographic and baseline characteristics of the study patients

Variable	Presence
Age (years)	62.71±6.01
Sex	
Male	209 (52.3)
Female	191 (47.7)
Previous myocardial infarction	146 (36.5)
Coronary artery disease	243 (60.8)
Hypertension	273 (68.3)
Diabetes mellitus	153 (38.3)
Smokers	70 (17.5)
Dyslipidemia	117 (29.3)
Family history	103 (25.8)
N (%), Mean ± S.D	

Table No.2: Association of myocardial infarction with troponin test at prehospital and in-hospital

Troponin result	Myocardial infarction		p-value
	Positive 128 (32.0%)	Negative 272 (68.0%)	
Prehospital test			
Positive, 33 (8.3%)	23 (18.0%)	10 (3.7%)	<0.001
Negative, 367 (91.7%)	105 (82.0%)	262 (96.3%)	
Sensitivity= $TP/(TP+FN)=23/(23+105)*100=18.0\%$ Specificity= $TN/(TN+FP)=262/(262+10)*100=96.3\%$			
At hospital test			
Positive, 215 (53.8%)	95 (74.2%)	120 (44.1%)	<0.001
Negative, 185 (46.2%)	33 (25.8%)	152 (55.9%)	
Sensitivity= $TP/(TP+FN)=95/(95+33)*100=74.2\%$ Specificity= $TN/(TN+FP)=152/(152+120)*100=55.9\%$			

DISCUSSION

The investigation examined whether pre-hospital TnT detection matches the established sensitivity and specificity for identifying acute myocardial infarction associated chest pain, as seen in emergency departments. The study revealed the practical form of conducting rapid TnT tests in ambulances, identification of point of care on positive TnT only a small fraction of acute myocardial infarction cases, displaying significantly lower sensitivity compared to TnT tests upon hospital admission.

The prehospital Troponin T (TnT) test's diminished sensitivity may be attributed to the brief time interval between the commencement of chest pain and the initial TnT test in the majority of the examined patients. TnT levels can elevate within 1 hour after the onset of chest pain. The sensitivity for MI remains about 50% until 3-4 hours after onset of pain.

The sensitivity for prehospital troponin to myocardial infarction was 18.0% and the sensitivity to myocardial infarction for in-hospital troponin was 74.2%. In another study, sensitivity of the rapid TnT test for acute MI was 33% in patients experiencing chest pain for less than 2 hours, and it notably increased to 86% for those with chest pain persisting for more than 8 hours.

Another study reported 18% sensitivity of prehospital TnT test and 98% in hospital for chest pain patients who were diagnosed with acute myocardial infarction later on. It was also concluded that positive prehospital Troponin T (TnT) test result serves as an objective indicator, suggesting a more unfavorable prognosis for individuals presenting with suspected acute chest pain.

Studies conducted by van Dongen et al¹¹ and Jungbauer et al¹⁵ reported that utilization of a point-of-care (POC) troponin T measurement, specifically the Roche Cobas h232 assay with a detection range of 40–2,000 ng/L (values below 40 ng/L reported as <40 ng/L), allows ambulance paramedics to swiftly calculate an on-site HEART score, potentially leading to a reassessment of the need for immediate transportation to the Emergency Department. The within-series coefficient of variation for this assay is 9.3% in the low concentration range (40–200 ng/L), demonstrating excellent sensitivity correlation between final results and rapid TnT.¹²

Rasmussen et al¹³ reported that performing pre-hospital point-of-care troponin measurements offers multiple potential advantages, as it enables the early detection of acute coronary syndrome (ACS) and facilitates the identification of high-risk patients prior to their arrival at the hospital. In another study Stopyra et al¹⁴ reported that the prehospital point-of-care (POC) i-STAT cardiac troponin (cTn) measurement in patients transported with acute chest pain has shown a high specificity for myocardial infarction (MI), indicating its potential usefulness in confirming MI.

In two previous studies conducted by Kemper et al¹⁵ and Sørensen et al¹⁶ concluded that point-of-care testing has become a practical reality in prehospital care, with cardiac troponin T (cTnT) offering crucial information that aids EMS personnel in decision-making, preventing the underestimation of serious pathologies. It serves as an alert trigger for various potentially severe conditions, guiding emergency medical professionals in determining the most effective strategies to be pursued.

Elevated levels of cardiac troponin T (cTnT) have been linked to higher hospital mortality rates, while troponinemia is associated with increased morbidity and the occurrence of serious adverse events in both cardiovascular and non cardiovascular diseases; however, limited research has investigated the impact of prehospital troponin levels in identifying early mortality in patients without acute coronary syndrome^{17,18}.

CONCLUSION

In regions where patient transport times are brief, the point-of-care rapid Troponin T (TnT) test reveals only a minority of individuals experiencing chest associated, prehospital TnT test positive outcome emerges as an objective indicator for a more adverse prognosis in those with suspected heart attacks.

Limitations: The study may not account for all possible confounding variables that could influence the relationship between prehospital troponin testing, paramedic risk assessment, and chest pain management. Factors such as comorbidities, medications, and socioeconomic status could impact the outcomes.

Practical Implications: Paramedics can use prehospital point-of-care troponin testing and risk assessment tools to identify patients at a higher risk of adverse cardiac events early in the care process. This early identification allows for quicker and more focused intervention for those at higher risk, potentially reducing the time to definitive treatment.

Author's Contribution:

Concept & Design of Study:	Feras Almarshad, Ghulam Mustafa
Drafting:	Feras Almarshad, Ghulam Mustafa
Data Analysis:	Feras Almarshad, Ghulam Mustafa
Revisiting Critically:	Feras Almarshad, Ghulam Mustafa
Final Approval of version:	Feras Almarshad, Ghulam Mustafa

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.116/01 dated 12.01.2021

REFERENCES

1. Dawson LP, Nehme E, Nehme Z, Zomer E, Bloom J, Cox S, et al. Chest pain management using prehospital point-of-care troponin and paramedic risk assessment. *JAMA Intern Med* 2023; 183(3):203-11.
2. Sagel D, Vlaar PJ, van Roosmalen R, Waardenburg I, Nieuwland W, Lettinga R et al. Prehospital risk stratification in patients with chest pain. *Emerg Med J* 2021;38(11):814-9.
3. Alghamdi A, Alotaibi A, Alharbi M, Reynard C, Body R. Diagnostic performance of prehospital point-of-care troponin tests to rule out acute myocardial infarction: a systematic review. *Prehospital Disaster Med* 2020;35(5):567-73.
4. van der Waarden NW, Schotting B, Royaards KJ, Vlachojannis G, Backus BE. Reliability of the HEART-score in the prehospital setting using point-of-care troponin. *Euro J Emerg Med* 2022;29(6):450-1.
5. Aarts GW, Camaro C, van Geuns RJ, Cramer E, van Kimmenade RR, Damman P, et al. Acute rule-out of non-ST-segment elevation acute coronary syndrome in the (pre) hospital setting by HEART score assessment and a single point-of-care troponin: rationale and design of the ARTICA randomised trial. *BMJ Open* 2020;10(2):e034403.
6. Pavlovsky T, Obadia M, Ragot S, Douay B, Casalino E, Ghazali DA. Predictors of Risk Stratification and Value of Point-of-Care of High-Sensitivity Cardiac Troponin-I in EMS Management of Non-ST-Segment Elevation Myocardial Infarction: A Retrospective Study. *Prehospital Disaster Med* 2022;37(3):365-72.
7. Ostrer IR, Wang TY. Are We Ready for Prehospital Troponin Testing? *JAMA Intern Med* 2023;183(3):211-2.
8. Wibring K, Lingman M, Herlitz J, Amin S, Bång A. Prehospital stratification in acute chest pain patient into high risk and low risk by emergency medical service: a prospective cohort study. *BMJ Open* 2021;11(4):e044938.
9. Cullen L, Collinson PO, Giannitsis E. Point-of-care testing with high-sensitivity cardiac troponin assays: the challenges and opportunities. *Emerg Med J* 2022;39(11):861-6.
10. Johannessen TR, Vallersnes OM, Halvorsen S, Larstorp AC, Mdala I, Atar D. Pre-hospital one-hour troponin in a low-prevalence population of acute coronary syndrome: OUT-ACS study. *Open Heart* 2020;7(2):e001296.
11. van Dongen DN, Tolsma RT, Fokkert MJ, Badings EA, van der Sluis A, Slingerland RJ, et al. Pre-hospital risk assessment in suspected non-ST-elevation acute coronary syndrome: A prospective observational study. *Eur Heart J Acute Cardiovasc Care* 2020; 9(1_suppl):5-12.
12. Jungbauer C, Hupf J, Giannitsis E, Frick J, Slagman A, Ehret C, et al. Analytical and clinical validation of a point-of-care cardiac troponin T test with an improved detection limit. *Clin Lab* 2017;63(4):633-45.
13. Rasmussen MB, Stengaard C, Sørensen JT, Riddervold IS, Hansen TM, Giebner M, et al. Predictive value of routine point-of-care cardiac troponin T measurement for prehospital diagnosis and risk-stratification in patients with suspected acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2019;8(4):299-308.
14. Stopyra JP, Snavely AC, Scheidler JF. Point-of-care troponin testing during ambulance transport to detect acute myocardial infarction. *Prehosp Emerg Care* 2020;24(6):751-9.
15. Kemper DW, Semjonow V, de Theije F. Analytical evaluation of a new point of care system for measuring cardiac troponin I. *Clin Biochem* 2017; 50(4-5):174-80.
16. Sørensen NA, Neumann JT, Ojeda F, et al. Diagnostic evaluation of a high-sensitivity troponin I point-of-care assay. *Clin Chem* 2019; 65(12):1592-1601.
17. Lippi G, Cervellin G, Sanchis-Gomar F. Prognostic value of troponins in patients with or without coronary heart disease: is it dependent on structure and biology? *Heart Lung Circ* 2020;29(3):324-30.
18. Árnadóttir Á, Falk Klein C, Iversen K. Head-to-head comparison of cardiac troponin T and troponin I in patients without acute coronary syndrome: a systematic review. *Biomarkers* 2017; 22(8):701-8.

Risk Factors for Epidural Anesthesia Blockade Failure in Cesarean Section

Epidural Anesthesia Blockade Failure in C-Section

Abid Haleem Khattak, Muhammad Sheharyar Ashraf, Amjid Ali, Jawad Hameed, Samar Naeem and Kashaf Noor

ABSTRACT

Objective: To identify and assess the risk factors associated with epidural failure during cesarean section anesthesia.

Study Design: Prospective study

Place and Duration of Study: This study was conducted at the Anesthesia, Lady Reading Hospital, Peshawar November 2022 to October, 2023.

Methods: A total of 400 patients were included in the study. In 2022, at Lady reading Hospital, Peshawar. Data of patients who had cesarean sections with epidural anesthesia (EA) and catheterization were collected. EA failure was identified as the need for intravenous anesthetics during the cesarean section, resulting in conversion to general anesthesia (GA).

Results: Most of the epidural failure patients 82.5% was applied method of loss of resistance to air. Further, catheter depth, resident, obstetric anesthesiologist, emergency surgery, rupture of membrane and parity in epidural failure and non-failure patients were almost equal, ($p>0.050$). Whereas, the mean waiting time in epidural failure patients was less than the non-failure patients as 13.23 ± 2.19 minutes and 15.22 ± 3.38 minutes, respectively.

Conclusion: Patients who have a previous epidural catheterization, experience inadequate waiting time, and are younger in age may face a higher risk of epidural analgesia (EA) failure. Specifically, the risk of EA failure increases by 2.6-fold for individuals with a previous epidural catheterization compared to those without catheterization history.

Key Words: Epidural Anesthesia, Risk Factors, Blockade Failure, Cesarean Section, Catheter Depth

Citation of article: Khattak AH, Ashraf MS, Ali A, Hameed J, Naeem S, Noor K. Risk Factors for Epidural Anesthesia Blockade Failure in Cesarean Section. Med Forum 2024;35(1):96-99. doi:10.60110/medforum.350122.

INTRODUCTION

Epidural anesthesia is favored technique of regional anesthesia for cesarean section in pregnant women due to the challenges associated with difficulty in airway management and the potential systemic effects of general anesthesia (GA) on both the fetus and uterine tone¹. Epidural anesthesia (EA) offers the advantage of prolonging anesthesia duration through the administration of additional local anesthetics². In patients who require pain control care after surgery use of catheter is effective, techniques such as programmed epidural bolus and epidural morphine can provide sufficient analgesia³.

In contrast, spinal anesthesia (SA) may require supplemental approaches like nerve blocks or

intravenous patient-controlled analgesia for extended pain control post-operatively⁴. It's worth noting that EA, particularly in cesarean sections, has a higher average failure rate ranging from 13.4% to 22.1%, compared to the lower range of 0.9% to 2.5% observed with SA⁵. Risk factors of epidural failure include prolonged labor, BMI, cesarean section urgency, labor analgesic breakthrough, maternal height and top ups of analgesics⁶.

Risk factors of procedure, such as anesthesia administered by non-obstetric anesthesiologists, the use of air for loss of resistance, and the flexibility of the catheter, have been identified^{7,8}. In cases where epidural analgesia (EA) fails, necessitating some extra IV anesthesia for sedation to attain a sufficient level of anesthesia or even requiring a conversion to general anesthesia (GA) with endotracheal intubation, potential hazards arise, particularly during sedation or GA conversion due to the possibility of encountering difficult airways^{9,10}.

Despite the widespread use of epidural anesthesia for C-sections, there is a paucity of comprehensive studies specifically focused on the risk factors associated with blockade failure in this population. Filling this research gap will contribute to the field of obstetric anesthesia

Department of Anesthesia, Lady Reading Hospital, Peshawar.

Correspondence: Dr. Jawad Hameed, Assistant Professor of Anesthesia, Lady Reading Hospital, Peshawar.

Contact No: 0333 9202031

Email: drjawadhameed@gmail.com

Received: November, 2023

Accepted: December, 2023

Printed: January, 2024

and perioperative care, guiding clinicians in refining their practices for better patient outcomes.

METHODS

The study conducted at Lady Reading Hospital in Peshawar from November 2022 to October, 2023. Study approved by Ethical committee and consent form patients was obtained. This study included patients who underwent c-section with epidural anesthesia and catheterization in the operating room, excluding those with specific criteria such as alternative epidural anesthetics, trainees with limited experience, short time intervals between anesthesia and incision, history of uncertain neuraxial anesthesia, history of spine abnormalities and surgical intervention, change from labor analgesia in epidural, and a history of spine surgery or abnormalities. The retrospective analysis involved dividing the 400 enrolled participants into two groups: those with epidural failure and those without. Epidural anesthesia and with cauterization was performed with 18-gauge needle and a 20-gauge catheter by positioning the patient in the right and lateral position. Following the loss of resistance (via air or saline), a catheter of 20-gauge was inserted into the epidural space, and testing dose of 3–5 mL was given. Subsequently, checks for signs of intrathecal and intravascular injections were conducted, and if none were observed, catheter was fixed. Mixture of anesthesia contain Sodium bicarbonate 2.8g, lidocaine 400 mg, epinephrine 0.1mg and fentanyl 100 mcg totally 15-24 ml was administered through the catheter into the epidural space. Anesthesia induction time was recorded. Following preparation for cesarean section, surgeons initiated the surgery after confirming pinprick sensation. In patients of blockade failure, the decision to continue with EA or switch to general anesthesia (GA) was determined based on the anesthesiologist's expertise.

SPSS version 27 was used for data analysis. Test of significance were t-test and chi square test and p value below 0.05 was taken as significant.

RESULTS

Out of 400 patients, 63 (15.8%) patients had epidural failure. (Figure. I). The mean age and BMI of epidural failure patients was 39.02±11.69 years and 28.57±3.82 kg/m². There were 37 (58.7%) epidural failure patients who had previous epidural analgesia than the non-failure patients 147 (43.6%), (p=0.027). The most common puncture site in epidural failure patients was L3-4, 48 (76.2%). Most of the epidural failure patients 52 (82.5%) was applied method of loss of resistance to air. Further, catheter depth, resident, obstetric anesthesiologist, emergency surgery, rupture of membrane and parity in epidural failure and non-failure patients were almost equal, (p>0.050). Whereas, the mean waiting time in epidural failure patients was less

than the non-failure patients as 13.23±2.19 minutes and 15.22±3.38 minutes, respectively. (Table. I).

Table No.1: Association of baseline characteristics according to epidural outcome

Characteristic	Epidural failure		p-value
	Yes 63 (15.8%)	No 337 (84.2%)	
Age (years)	39.02±11.69	39.16±11.80	0.929
BMI (kg/m ²)	28.57±3.82	28.40±3.13	0.713
Previous epidural analgesia	37 (58.7)	147 (43.6)	0.027
Puncture site			
L2-3	13 (20.6)	43 (12.8)	0.186
L3-4	48 (76.2)	273 (81.0)	
L4-5	2 (3.2)	21 (6.2)	
Loss of resistance methods			
Air	52 (82.5)	293 (86.9)	0.352
Saline	11 (17.5)	44 (13.1)	
Catheter depth(cm)	5.26±1.18	5.32±1.91	0.697
Resident	46 (73.0)	272 (80.7)	0.165
Obstetric anesthesiologist	8 (12.7)	61 (18.1)	0.298
Emergency surgery	26 (41.3)	149 (44.2)	0.666
Rupture of membrane	10 (15.9)	51 (15.1)	0.8881
Parity			
Nulliparous	23 (36.5)	152 (45.1)	0.207
Parous	40 (63.5)	185 (54.9)	
Waiting time (minute)	13.23±2.19	15.22±3.38	<0.001
N (%), Mean ± S.D			




Figure No.1: Distribution of epidural failure among the study patients

DISCUSSION

In this study, we investigated the factors contributing to the failure of converting labor analgesia to cesarean delivery anesthesia, identifying key risk factors such as procedures conducted by trainees, parturients with elevated BMI, and the utilization of air for the loss of resistance test, as reported in prior studies¹¹. The mean age and BMI of epidural failure patients was 39.02±11.69 years and 28.57±3.82 kg/m² and there was no significant difference among epidural failure and non failures.

Studies by Bauer et al¹² and Grap et al¹³ have consistently highlighted a correlation between age and the risk of epidural failure, with a noteworthy trend indicating that younger patients may face a higher likelihood of experiencing this complication. The literature consistently reports a positive association between younger age and the incidence of epidural failure, suggesting that age should be considered as a significant risk factor in assessing the effectiveness of epidural procedures.

Previous studies conducted by Kula et al¹⁴ and Eley et al¹⁵ have provided evidence suggesting that an elevated Body Mass Index (BMI) is associated with increased technical difficulties and a higher likelihood of failure in neuraxial anesthesia. Additionally, these studies have indicated that obese parturients face an elevated risk of extension failure to surgical anesthesia, highlighting the challenges and complications that obesity can introduce in the administration and effectiveness of anesthesia procedures during childbirth.

Most of the epidural failure patients 82.5% were applied method of loss of resistance to air as compare to saline 17.5%. Beilin et al¹⁶ reported in their study that the loss of resistance to air, as opposed to saline, may elevate the risk of epidural failure, a conclusion supported by Shenouda et al¹⁷, who also observed that air could potentially impact the spread of local anesthetic, leading to an incomplete "patchy block" and consequently an increased reliance on intraoperative intravenous anesthetics.

In this study epidural failure rate was higher 58.7% patients who experienced previous epidural analgesia than non failures. Shimada et al¹⁸ in a study revealed notable inflammatory adhesions and changes in individuals with a history of epidural anesthesia (EA) utilizing an epidural scope, as puncture of the flavum ligament and congestion due to catheterization within the epidural space, ultimately causing disruptions in the proper spread of local anesthetic within the epidural compartment.

The mean waiting time in epidural failure patients was less than the non-failure patients as 13.23±2.19 minutes and 15.22±3.38 minutes, respectively. The administration of epidural anesthetics indicated a nearly identical trajectory before the 12-minute mark¹⁹, implying that the primary cause of failure was predominantly attributed to insufficient waiting time for the lidocaine-bicarbonate-epinephrine-fentanyl combination to achieve surgical anesthesia at the T7 level²⁰.

CONCLUSION

Patients who have a previous epidural catheterization, experience inadequate waiting time, and are younger in age may face a higher risk of epidural analgesia (EA) failure. Specifically, the risk of EA failure increases by 2.6-fold for individuals with a previous epidural

catheterization compared to those without catheterization history.

Author's Contribution:

Concept & Design of Study: Abid Haleem Khattak
Drafting: Muhammad Sheharyar Ashraf, Amjid Ali

Data Analysis: Jawad Hameed, Samar Naeem, Kashaf Noor

Revisiting Critically: Abid Haleem Khattak, Muhammad Sheharyar Ashraf

Final Approval of version: Abid Haleem Khattak

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.193/LRH/MTI dated 15.10.2022

REFERENCES

1. Ashagrie HE, Ahmed SA, Melesse DY. The incidence and factors associated with failed spinal anesthesia among parturients underwent cesarean section, 2019: a prospective observational study. *Intern J Surg Open* 2020;24:47-51.
2. Young B, Onwochei D, Desai N. Conventional landmark palpation vs. preprocedural ultrasound for neuraxial analgesia and anaesthesia in obstetrics—a systematic review and meta-analysis with trial sequential analyses. *Anaesthesia* 2021;76(6):818-31.
3. Riaz F, Iqbal A, Haider MS. Comparison of Postoperative Analgesic Duration of Intrathecal Dexmedetomidine Versus Buprenorphine as Adjuvant to 0.5% Heavy Bupivacaine in Spinal Anesthesia for Orthopedic Surgeries. *Med J South Punjab* 2023;4(2):70-6.
4. Stewart J, Gasanova I, Joshi GP. Spinal anesthesia for ambulatory surgery: current controversies and concerns. *Current Opinion Anesthesiol* 2020; 33(6):746-52.
5. Desai N, Carvalho B. Conversion of labour epidural analgesia to surgical anaesthesia for emergency intrapartum Caesarean section. *BJA Educ* 2020;20(1):26-31.
6. Wang CH, Yeh PH, Wu JH. Failed epidural analgesia due to accidental placement into the lumbar plexus space by loss-of-resistance technique. *The Changhua J Med* 2020;18(1):27-9.
7. Bjornestad EE, Haney M. An obstetric anaesthetist: A key to successful conversion of epidural analgesia to surgical anaesthesia for caesarean delivery? *Acta Anaesthesiologica Scandinavica* 2020;64(2):142-4.
8. Khan I, Ashraf MS, Gill MU. Work Base Assessment in Anesthesia for Postgraduate Residents. *Med J South Punjab* 2022;3(1):17-20.

9. Shen C, Chen L, Yue C, Cheng J. Extending epidural analgesia for intrapartum cesarean section following epidural labor analgesia: a retrospective cohort study. *J Maternal-Fetal Neonatal Med* 2022;35(6):1127-33.
10. Carvalho B, Riley E, Cohen SE, Gambling D, Palmer C, Huffnagle HJ, et al. Single-dose, sustained-release epidural morphine in the management of postoperative pain after elective cesarean delivery: results of a multicenter randomized controlled study. *Anesth Analg* 2005;100(4):1150-8.
11. Lee JY, Noh KM, Lee SW, Cho EY, Hong JH. Analysis of Factors Affecting Thoracic Epidural Anesthesia Performance Time. *Keimyung Med J* 2020;39(1):33-7.
12. Bauer ME, Kountanis JA, Tsen LC, Greenfield ML, Mhyre JM. Risk factors for failed conversion of labor epidural analgesia to cesarean delivery anesthesia: a systematic review and meta-analysis of observational trials. *Int J Obstet Anesth* 2012;21(4):294-309.
13. Grap SM, Patel GR, Huang J, Vaida SJ. Risk factors for labor epidural conversion failure requiring general anesthesia for cesarean delivery. *J Anaesthesiol Clin Pharmacol* 2022;38(1):118-23.
14. Kula AO, Riess ML, Ellinas EH. Increasing body mass index predicts increasing difficulty, failure rate, and time to discovery of failure of epidural anesthesia in laboring patients. *J Clin Anesth* 2017;37:154-8.
15. Eley VA, Chin A, Tham I, Poh J, Aujla P, Glasgow E, et al. Epidural extension failure in obese women is comparable to that of non-obese women. *Acta Anaesthesiol Scand* 2018;62(6):839-47.
16. Beilin Y, Arnold I, Telfeyan C, Bernstein HH, Hossain S. Quality of analgesia when air versus saline is used for identification of the epidural space in the parturient. *Reg Anesth Pain Med* 2000;25(6):596-9.
17. Shenouda PE, Cunningham BJ. Assessing the superiority of saline versus air for use in the epidural loss of resistance technique: a literature review. *Reg Anesth Pain Med* 2003;28(1):48-53.
18. Shimada N, Igarashi T, Murai K, Hara T, Kuramochi T, Takeuchi M. Adhesions in the epidural space caused by frequent epidural blocks. *JA Clin Rep* 2017;3(1):p57.
19. Goring-Morris J, Russell IF. A randomised comparison of 0.5% bupivacaine with a lidocaine/epinephrine/fentanyl mixture for epidural top-up for emergency caesarean section after low dose epidural for labour. *Int J Obstet Anesth* 2006;15(2):109-14.
20. Hillyard SG, Bate TE, Corcoran TB, Paech MJ, O'Sullivan G. Extending epidural analgesia for emergency caesarean section: a meta-analysis. *Br J Anaesth* 2011;107(5):668-78.

Antiemetic Prophylaxis with Droperidol in Morphine-Based Intravenous Patient Controlled Analgesia

Antiemetic Prophylaxis with Droperidol in Morphine Analgesia

Muhammad Sheharyar Ashraf, Abid Haleem Khattak, Jawad Hameed, Amjid Ali, Kashaf Noor and Samar Naeem

ABSTRACT

Objective: To assess the antiemetic advantages and sedative impacts of droperidol when used in conjunction with morphine-based intravenous patient-controlled analgesia (IV-PCA).

Study Design: Cohort study

Place and Duration of Study: This study was conducted at the Lady Reading Hospital in Peshawar from December 2022 to November 2023.

Methods: Patients who underwent major surgery and utilized morphine-based IV-PCA experienced a primary outcome characterized by the rate of any postoperative nausea and/or vomiting (PONV) occurring within 72 hours after the surgical procedure.

Results: Nausea and vomiting between 0-12 hours after operation in Droperidol Group was 10.7% and 14.7% in control group. Nausea and vomiting between 12-36 hours after operation in Droperidol Group was 12.0% and 17.3% in control group. Nausea and vomiting between 36-60 hours after operation in Droperidol Group was 13.3% and 16.0% in control group. Nausea and vomiting between 60-72 hours after operation in Droperidol Group was 12.7% and 16.7% in control group.

Conclusion: Droperidol into intravenous patient-controlled analgesia (IV-PCA) regimens has demonstrated a notable reduction in the risk of postoperative nausea and vomiting (PONV).

Key Words: Droperidol, Morphine, Antiemetic prophylaxis, Patient controlled analgesia

Citation of article: Ashraf MS, Khattak AH, Hameed J, Ali A, Noor K, Naeem S. *Antiemetic Prophylaxis with Droperidol in Morphine-Based Intravenous Patient Controlled Analgesia. Med Forum 2024;35(1):100-103. doi:10.60110/medforum.350123.*

INTRODUCTION

Postoperative nausea and vomiting (PONV) emerges as a prevalent source of patient distress post-surgery, with reported rates ranging from 20% to 40%¹. The multifactorial nature of PONV is evident, encompassing patient-related factors such as sex, smoking status, and a history of PONV², as well as surgery-related factors like the type of surgical procedure, and factors related to anesthesia including the use of volatile and opioids anesthetics. Incidence of PONV can vary around 80% in high risk patients³.

PONV is often reported by surgical patients as a more challenging issue than postoperative pain, despite its typically self-limited nature⁴; however, vomiting can persist rarely in but it can contribute in serious complications, including pneumothorax, pulmonary

aspiration, wound dehiscence and elevated intracranial pressure⁵. Moreover, PONV may extend the duration ICU stay and lead to unexpected hospitalization after ambulatory surgery. The treatment of PONV imposes a significant burden on healthcare economy⁶.

IV-PCA proves to be a highly effective approach for alleviating postoperative acute pain; however, the prevalent use of opioids as the primary analgesic in IV-PCA is associated with a common adverse event⁷, PONV, with reported rates ranging from 18 to 23%⁸. Notably, approximately twelve percent of surgical patients opt to discontinue IV-PCA prematurely due to the challenging nature of intractable PONV⁹. To address this issue, droperidol, a D2 receptor antagonist, is employed for its central action on the chemoreceptor trigger zone, serving as an antiemetic agent in the context of IV-PCA¹⁰.

The antiemetic effectiveness of droperidol was proven in opioid-based IV-PCA; however, prior investigations exhibited methodological shortcomings, such as small patient samples ($n < 1,000$), inadequate adjustment for confounding factors, exclusive focus on female patients, and a narrow scope of surgical procedures¹¹. Moreover, the majority of earlier studies relied on data dating back more than two decades, failing to capture the advancements in different surgical interventions and anesthetic related care, like multimodal analgesia and

Department of Anesthesia, Lady Reading Hospital, Peshawar.

Correspondence: Dr. Jawad Hameed, Assistant Professor of Anesthesia, Lady Reading Hospital, Peshawar.

Contact No: 0333 9202031

Email: drjawadhameed@gmail.com

Received: December, 2023

Accepted: December, 2023

Printed: January, 2024

minimally invasive surgery that have occurred in recent years¹².

METHODS

The study conducted at Lady Reading Hospital in Peshawar from December 2022 to November 2023. Study approved by Ethical committee and consent form patients was obtained. Patients who underwent any surgical intervention under neuraxial or general anesthesia and were given opioid-based IV-PCA for pain management in post-operative time were enrolled. Patients < 20 years, switching droperidol, using non-morphine analgesics for IV-PCA were excluded. Patients were divided into control groups and droperidol group.

IV-PCA is contraindicated for patients unable to maintain consciousness, those with cognitive impairment, and those requiring intensive care and mechanical ventilation after 24 hours. It is initiated in the intensive care unit after anesthesia using an ambulatory infusion pump programmed for morphine sulfate delivery.

The infusion settings for the IV-PCA system encompass a loading dose range of 0 to 5.0 mL, a demand dose varying from 0.5 to 2.0 mL, a basal infusion rate spanning 0 to 1.5 mL per hour, and a lockout time set between 5 and 10 minutes. Additionally, antiemetic prophylaxis is implemented by incorporating droperidol at a concentration ranging from 0.025 to 0.075 mg/mL into the IV-PCA infusate.

The researcher assessed patients' responses every 12 hours, increasing frequency for inadequate analgesia or adverse events. PONV severity was categorized using a 4-point scale: no PONV, mild PONV (nausea without antiemetic), moderate PONV (nausea with antiemetic request), and severe PONV (nausea with vomiting requiring antiemetic treatment).

The main focus of the study was to assess the incidence of PONV within 72 hours as the primary outcome. Certified nurse regularly evaluated the pain intensity, sedation level and occurrence of PONV at 12-hour intervals during the 72-hour postoperative period at the institution.

Anesthesia was induced with 1–2 mg/kg propofol and 1–2 µg/kg fentanyl, using 0.6–1.0 mg/kg rocuronium for intubation. Maintenance involved sevoflurane or desflurane. Reversal agents like 2 mg/kg sugammadex were used for neuromuscular blockade. Spinal anesthesia utilized 6–15 mg bupivacaine without opioids. Combined neuraxial and general anesthesia included epidural ropivacaine (5 mg/mL) with or without fentanyl (2.5–5 µg/mL). Midazolam (2–5 mg) provided anxiolysis. Perioperative fluid management involved crystalloid fluids following practice guidelines.

RESULTS

Overall, 300 patients were included in this study both sex. They were two equal groups in this study as Droperidol, 150 (50.0%) and Control, 150 (50.0%). The distribution of demographics and baseline characteristics in Droperidol and Control group were almost equal, and the differences were statistically significant, (p>0.050). (Table 1).

Table No.1: Demographic and baseline characteristics of both the study groups

Characteristic	Group		p-value
	Droperidol	Control	
Age (years)	53.80±5.94	54.88±5.59	0.904
BMI (kg/m ²)	26.67±2.19	27.57±2.18	0.696
Sex			
Male	82 (54.7)	80 (53.3)	0.817
Female	68 (45.3)	70 (46.7)	
ASA status			
I	32 (21.3)	30 (20.0)	0.515
II	111 (74.0)	120 (80.0)	
III	7 (4.7)	0 (0.0)	
Smoking status	36 (24.0)	25 (16.7)	0.115
Previous PONV	12 (8.0)	21 (14.0)	0.097
Hypertension	45 (30.0)	39 (26.0)	0.654
Diabetes mellitus	46 (30.7)	42 (28.0)	0.612
Major depression	3 (2.0)	8 (5.3)	0.125
Malignancy	24 (16.0)	25 (16.7)	0.876
Hemoglobin (g/dL)	12.28±2.51	12.13±2.25	0.600
eGFR (mL/min/1.73 m ²)	98.22±3.09	98.71±3.46	0.187
Alanine aminotransferase (U/L)	19.21±2.38	19.02±2.24	0.479
Aspartate aminotransferase (U/L)	22.25±1.48	22.34±1.32	0.525
Type of anesthesia			
Neuraxial anesthesia	60 (40.0)	56 (37.3)	0.771
General Anesthesia	89 (59.3)	92 (61.3)	
Combined	1 (0.7)	2 (1.3)	
Mean ± S.D, N (%)			

Nausea and vomiting between 0-12 hours after operation in Droperidol Group was 16 (10.7%) and 22 (14.7%) in control group, (p=0.741). Nausea and vomiting between 12-36 hours after operation in Droperidol Group was 18 (12.0%) and 26 (17.3%) in control group, (p=0.462). Nausea and vomiting between 36-60 hours after operation in Droperidol Group was 20 (13.3%) and 24 (16.0%) in control group, (p=0.862). Whereas, nausea and vomiting between 60-72 hours after operation in Droperidol Group was 19 (12.7%)

and 25 (16.7%) in control group, (p=0.868). Further, the severity of nausea and vomiting in both the groups were almost equal, (p>0.050). (Table 2).

Table No.2: Distribution of nausea and vomiting of both the study groups

	Group		p-value
	Droperidol	Control	
POH 0–12	16 (10.7)	22 (14.7)	0.741
Mild	12 (75.0)	17 (77.3)	0.532
Moderate	3 (18.8)	4 (18.2)	
Severe	1 (6.2)	1 (4.5)	
POH 12-36	18 (12.0)	26 (17.3)	0.462
Mild	11 (61.1)	18 (69.2)	0.741
Moderate	5 (27.8)	4 (15.4)	
Severe	2 (11.1)	4 (15.4)	
POH 36-60	20 (13.3)	24 (16.0)	0.862
Mild	18 (90.0)	18 (75.0)	0.684
Moderate	1 (5.0)	4 (16.7)	
Severe	1 (5.0)	2 (8.3)	
POH 60-72	19 (12.7)	25 (16.7)	0.868
Mild	15 (78.9)	21 (84.0)	0.796
Moderate	2 (10.5)	4 (16.0)	
Severe	2 (10.5)	0 (0.0)	
N (%)			

DISCUSSION

The study revealed a significant reduction in the incidence of postoperative nausea and vomiting (PONV) with the addition of droperidol to morphine-based intravenous patient-controlled analgesia (IV-PCA). Subgroup analyses demonstrated that the droperidol effect was particularly notable in patients under 65 years of age, females, non-smokers, and those without a history of PONV.

In a study conducted by an author, it was observed that patients administered with droperidol exhibited significantly lower levels of nausea at the 12-hour mark, and within the first 24 hours, only 31% of these patients required prochlorperazine, compared to 59.3% of those not receiving droperidol. Additionally, the droperidol group showed a significantly higher number of patients experiencing sedation at the 24-hour mark. Similar findings were reported in another study that addition of droperidol significantly reduced PONV in morphine-based IV-PCA, especially in patients under 65, females, non-smokers, and those without a history of PONV.

Another study reported that the antiemetic impact of droperidol was notably effective within the first 36 hours post-surgery but diminished thereafter. Uda et al¹³ conducted a study in which they proposed that the incorporation of droperidol into intravenous patient-controlled analgesia (IV-PCA) regimens resulted in a notable reduction in the incidence of postoperative nausea and vomiting (PONV) within the initial 36 hours following surgery. However, their findings indicated

that the antiemetic efficacy of droperidol appeared to diminish beyond this specified time frame, suggesting a time-dependent attenuation of its preventive effects against PONV in the postoperative period.

In their study, Kuo et al¹⁴ found that the inclusion of droperidol resulted in a notable decrease in both the frequency and intensity of postoperative nausea and vomiting (PONV) specifically on postoperative days 2 and 3, with no significant impact observed on day 1. Different droperidol regimens in IV-PCA, concluding that a 0.10 mg/mL dose demonstrated optimal antiemetic efficacy with minimal sedation risk. Combining their results with ours, it suggests that adding droperidol at 0.025–0.10 mg/mL to opioid-based IV-PCA is appropriate, considering the benefit-risk balance.

Tan et al¹⁵ found that the addition of droperidol to intravenous patient-controlled analgesia (IV-PCA) effectively decreased the risk of postoperative nausea and vomiting (PONV) without causing an increase in opioid consumption or altering the level of sedation; nevertheless, they emphasized the necessity for supplementary prophylactic interventions to address the occurrence of late-onset PONV. Gan et al¹⁶ conducted studies indicating a significant reduction in postoperative nausea and vomiting (PONV) over a 24-hour period when administering a perioperative 1.25 mg bolus of droperidol in patients utilizing patient-controlled analgesia (PCA).

CONCLUSION

The incorporation of droperidol into intravenous patient-controlled analgesia (IV-PCA) regimens has demonstrated a notable reduction in the risk of postoperative nausea and vomiting (PONV), while concurrently exhibiting no discernible impact on opioid consumption or the level of sedation. Despite these encouraging outcomes, it is important to acknowledge that the efficacy of droperidol may be limited to the prevention of immediate postoperative PONV, thereby suggesting a potential need for supplementary prophylactic interventions to address the occurrence of late-onset PONV.

Author’s Contribution:

- Concept & Design of Study: Muhammad Sheharyar Ashraf
- Drafting: Abid Haleem Khattak, Jawad Hameed
- Data Analysis: Amjid Ali, Kashaf Noor, Samar Naeem
- Revisiting Critically: Muhammad Sheharyar Ashraf, Abid Haleem Khattak
- Final Approval of version: Muhammad Sheharyar Ashraf

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.192/LRH/MTI dated 15.10.2022

REFERENCES

1. Amirshahi M, Behnamfar N, Badakhsh M, Rafiemanesh H, Keikhaie KR, Sheyback M, et al. Prevalence of postoperative nausea and vomiting: A systematic review and meta-analysis. *Saudi J Anaesth* 2020;14(1):48.
2. Ghosh S, Rai KK, Shivakumar HR, Upasi AP, Naik VG, Bharat A. Incidence and risk factors for postoperative nausea and vomiting in orthognathic surgery: a 10-year retrospective study. *J Korean Associat Oral Maxillofac Surg* 2020;46(2):116.
3. Suh S, Helm M, Kindel TL, Goldblatt MI, Gould JC, et al. The impact of nausea on post-operative outcomes in bariatric surgery patients. *Surg Endosc* 2020;34:3085-91.
4. Ahmed SA, Lema GF. Incidence and factors associated with postoperative nausea and vomiting among elective adult surgical patients at University of Gondar comprehensive specialized hospital, Northwest Ethiopia, 2019: a cross-sectional study. *Int J Surg Open* 2020;22:57-61.
5. Elsaid RM, Namrouti AS, Samara AM, Sadaqa W, Zyoud SE. Assessment of pain and postoperative nausea and vomiting and their association in the early postoperative period: an observational study from Palestine. *BMC Surg* 2021;21(1):1-9.
6. Stephenson SJ, Jiwanmall M, Cherian NE, Williams A. Reduction in post-operative nausea and vomiting (PONV) by preoperative risk stratification and adherence to a standardized anti emetic prophylaxis protocol in the day-care surgical population. *J Family Med Primary Care* 2021;10(2):865-70.
7. Lee HM, Kil HK, Koo BN, Song MS, Park JH. Comparison of sufentanil-and fentanyl-based intravenous patient-controlled analgesia on postoperative nausea and vomiting after laparoscopic nephrectomy: a prospective, double-blind, randomized-controlled trial. *Int J Med Sci* 2020;17(2):207.
8. Shim JG, Ryu KH, Cho EA, Ahn JH, Cha YB, Lim G, et al. Machine learning for prediction of postoperative nausea and vomiting in patients with intravenous patient-controlled analgesia. *PLoS One* 2022;17(12):e0277957.
9. Wolfe RC, Bequette J. Dopamine Receptor Antagonists for the Prevention and Treatment of Postoperative Nausea and Vomiting. *J Perianesth Nursing* 2021;36(2):199-202.
10. Kranke P, Wilhelm W, Eberhart L. Management of postoperative nausea and vomiting (PONV). *Enhanced Recovery After Surgery: A Complete Guide to Optimizing Outcomes* 2020:195-202.
11. Apfel CC, Heidrich FM, Jukar-Rao S, Jalota L, Hornuss C, Whelan RP, et al. Evidence-based analysis of risk factors for postoperative nausea and vomiting. *Br J Anaesth* 2012;109:742–53.
12. Sweeney BP. Why does Smoking protect against PONV? *Br J Anaesth* 2002;89:810–3.
13. Uda S, Takeda C, Mizota T. Effect of droperidol addition to fentanyl-based intravenous patient-controlled analgesia on postoperative nausea and vomiting: a single-center retrospective cohort study. *J Clin Rep* 2020;6:90.
14. Kuo YM, Tsou MY, Chang WK, Chan KH, Chang KY. To add or not to add? An empirical study on droperidol and intravenous patient-controlled analgesia. *J Chin Med Assoc* 2012;75:227–33.
15. Tan JQ, Wu HL, Wang YC, Cata JP, Chen JT, Cherng YG et al. Antiemetic prophylaxis with droperidol in morphine-based intravenous patient-controlled analgesia: a propensity score matched cohort study. *BMC Anesthesiol* 2023;23(1):351.
16. Gan TJ, Belani KG, Bergese S, Chung F, Diemunsch P, Habib AS, et al. Fourth Consensus guidelines for the management of postoperative nausea and vomiting. *Anesth Analg* 2020;131:411–48.

Micro and Macrocytic Anemia – A Population Based Cross-Sectional Study

Micro and
Macrocytic
Anemia

Ayeshah Zaib-Un-Nisa¹, Iyad Naeem Muhammad², Sheikh Abdul Khaliq³ and Agha Umer Draz Khan⁴

ABSTRACT

Objective: The main objective of current study was to determine age-standardized point prevalence of iron, folic acid and vitamin B-12 deficiency anemia.

Study Design: A population based cross-sectional study

Place and Duration of Study: This study was conducted at the Department of Pharmacy Practice, Hamdard University and University of Karachi from January-2023 and ended on April-2023.

Methods: Data collected from blood banks. Collected blood samples were analyzed by Hematology Analyzer, Sysmex XP-100 and XN-1000. Analysis of data was done by Statistical Package for Social Sciences version-22. Main outcome measures: Levels of hemoglobin, hematocrit, microcytosis, macrocytosis

Results: Among 8,134 patients and donors data; male 37% (N=3043) and female 63% (N=5091). Age range for the majority of population was 18-28 years. In female; hemoglobin and hematocrit were lower-than normal in 39% (N=1984) and 53% (N=2685) respectively; microcytosis found in 31% (N=1554) and macrocytosis in 7% (N=325). In male; hemoglobin and hematocrit were lower-than normal in 33% (N=994) and 34% (N=1048) respectively; microcytosis reported in 25% (N=739) and macrocytosis in 8% (N=223). In both genders; significant (p=0.0001) differences are noted in the mean values of RBC indices versus the mean value of standard.

Conclusion: Evaluation of data reveals that hypochromic-microcytic anemia highly prevailed in the society. Majority of such population is in between 18-28 years old. Mean values of RBC indices in both gender were significantly lower than standard.

Key Words: Anemia; prevalence; iron; folic-acid; vitamin B-12; microcytosis.

Citation of article: Zaib-Un-Nisa A, Muhammad IN, Khaliq SA, Khan AUD. Micro and Macrocytic Anemia – A Population Based Cross-Sectional Study. Med Forum 2024;35(1):104-108. doi:10.60110/medforum.350123.

INTRODUCTION

Anemia should not be underestimated; it may significantly increases the risk of cardiovascular diseases (CVD) and renal disease (RD) in hypertensive individuals.¹ Cardio-renal Anemia Syndrome (CRAS) patients when treated with iron supplements and erythropoietin; heart failure and kidney injury were addressed.² Similarly among children; iron deficiency anemia may retard their psychomotor development.³

¹. Department of Pharmacy Practice, Hamdard University and University of Karachi.

². Department of Pharmacy Practice, University of Karachi.

³. Department of Pharmacy Practice, Hamdard University, Karachi.

⁴. Department of Pathology and Hematology, Hussaini Hematology & Oncology Trust, Karachi.

Correspondence: Sheikh Abdul Khaliq, Pharmacy Practice, Hamdard University, Karachi.

Contact No: 0345-2670820

Email: drsheikh1974@gmail.com

Received: August, 2023

Accepted: November, 2023

Printed: January, 2024

Among women; despite high prevalence of iron deficiency anemia, it is under-diagnosed and under-treated; which results in adverse health consequences pertaining to emotional and physical health.⁴ Poor clinical prognosis also reported in patients of Acute Coronary Syndrome (ACS) having hemoglobin less than normal.⁵

According to the World Health Organization (WHO); the adult individual will be considered anemic; if hemoglobin levels is less than 13 g/dL in men and 12 g/dL in women.⁶ However; based on unique apparatus; techniques employed and the characteristics of patient population; each laboratory must establish their own reference values.⁷ Anemia is an important global health issue. Roughly one-third of people on the planet (32.9%) are anemic.⁸ According to Global age-standardized point prevalence rate of anemia is 23,176.20 per 100,000 and years of healthy life lost due to disability (YLD) rates is 672.4per 100,000.⁹ South Asia has the greatest frequency of age-standardized YLD from anemia (1358.2 per 100,000 people).⁹ Anemia can be caused by inadequate erythropoiesis by dietary deficiencies, inflammation, or hereditary hemoglobinopathies (thalassemia, sickle-cell-trait); or excessive erythrocyte loss (due to blood-loss, hemolysis, or both).¹⁰ Anemia is usually

categorized according to the erythrocyte morphology (e.g. microcytic, macrocytic, normocytic); and/or underlying etiology (e.g. iron deficiency, inflammation and hemolysis).¹⁰ Nutritional deficiencies and chronic illnesses are often the most frequent etiologies of anemia in children and young adults.¹¹

The main objective of current study was to determine age-standardized point prevalence of iron, folic-acid and vitamin B12 deficiency anemia. For this purpose after collection of blood samples; hematological indices of individuals were determined along with demography of individuals.

METHODS

Design of Study, Place and Duration: The study design was population based cross-sectional. Study began on 15th January-2023 and Ended on 29th April-2023 in the city of Karachi.

Ethical Statement/Approval: Ethical approval of study has been taken before initiating study from Institutional Bioethical Committee (Reference No. IBC KU-300/2023), University of Karachi. Prior to initiating the study; all researchers ensured the maintenance of patient data confidentiality in compliance with the Declaration of Helsinki¹² and also taken written informed consent before data collection.

Data Collection Method: Data has been collected from the well know blood banks and blood transfusion centers of the Karachi city. Precision analysis technique was used for the determination of sample size of study.¹³ The study includes around 8,134 patients and donors of blood from different centers; male 3043 (37%) and female 5091 (63%). Age range of population was neonatal (less than one month) to 95 years in both genders.

Procedure: After taken written informed consent; blood samples were collected through veni-puncture by professional staff into a 3mL vacutainers-tube with EDTA. Hemoglobin (Hb), hematocrit (Hct), mean-corpuscular-volume (MCV), mean-corpuscular-hemoglobin (MCH), mean-corpuscular-hemoglobin-concentration (MCHC) and red-blood-cell count (RBC) were determined by using Hematology Analyzer, Sysmex XP - 100 and XN – 1000.

Inclusion criteria: No co-morbidity in the selected subjects.

Exclusion criteria: Subjects having malignancy, infection, inflammation, any chronic disease that may affect the analyzing parameters, blood transfusion in last three months or received iron therapy.

Assessment of Data: The data for study was analyzed by SPSS (Statistical Package for Social Sciences) version-22. Descriptive (Frequency distribution, histograms) and inferential statistics (Student t-test) applied. Blood indices were compared by student t-test versus standard after keeping significance value of probability (p) <0.05.

RESULTS

Age range for the majority of population was 18-28 years. Among females, anemia is reported in 1984 (39%); 2685 (53%) has Hematocrit (Hct) value lower than normal; microcytosis reported in 1554 (31%) and macrocytosis in 325 (7%). (Table No. 1)

In male gender; anemia is reported in 994 (33%); 1048 (34%) has hematocrit value lower than normal; microcytosis reported in 739 (25%); macrocytosis in 223 (8%). (Table No. 2)

Descriptive statistical findings of total population for age and RBC (Red Blood Cells) indices mentioned in Table No. 3; however, in both genders significant differences are noted in the mean values of RBC indices versus the mean value of standard.¹⁵ (Table 4)

Table No. 1: CBC* parameters reporting Anemia in Female Gender

CBC* Parameters	Normal Range ¹⁵	N (%)	Mean± SD
Hemoglobin	<11g/dL	1984 (39%)	9.27±1.48
	11-14.5g/dL	2887 (57%)	12.35±0.89
	>14.5g/dL	220 (4%)	19.84±4.77
Hematocrit	<34.5%	2685 (53%)	29.2±3.30
	34.5-45.4%	2251 (44%)	38.36±2.55
	>45.4%	155 (3%)	63.33±13.72
MCV (Mean Corpuscular Volume)	<78.1 fL	1554 (31%)	70.06±6.48
	78.1-95.3 fL	3073 (62%)	85.19±4.17
	>95.3 fL	325 (7%)	124.47±34.15
MCHC (Mean Corpuscular Hemoglobin Concentration)	<30.3 g/dL	815 (16%)	28.48±1.88
	30.3-34.4 g/dL	3858 (78%)	32.27±1.02
	>34.4 g/dL	279 (6%)	47.73±15.28
MCH Mean Corpuscular Hemoglobin)	<25.3 pg	1761 (36%)	21.58±2.97
	25.3-31.7 pg	2920 (59%)	27.95±1.55
	>31.7 pg	271 (5%)	42.42±11.02
RBC (Red Blood Cells) Count	<3.61 x10 ¹² /Lit.	482 (10%)	3.101±0.51
	3.61-5.2 x10 ¹² /Lit.	3949 (80%)	4.41±0.39
	>5.2 x10 ¹² /Lit.	521 (11%)	6.306±1.56

*Complete blood Count

Table No. 2: CBC* parameters reporting Anemia in Male Gender

CBC* Parameters	Normal Range ¹⁵	N (%)	Mean±SD
Hemoglobin	<12.3 g/dL	994 (33%)	9.98±1.93
	12.3-16.6 g/dL	1855 (61%)	14.38±1.14
	>16.6 g/dL	194 (6%)	20.203±4.97
Hematocrit	<38.4%	1048 (34%)	32±5.45
	38.4-50.7%	1845 (61%)	43.88±3.05
	>50.7%	150 (5%)	63.88±15.96
MCV (Mean Corpuscular Volume)	<78.7 fL	739 (25%)	70.15±26.84
	78.7-96.3 fL	2010 (68%)	86.4±26.13
	>96.3 fL	223 (8%)	118.32±31.83
MCHC (Mean Corpuscular Hemoglobin Concentration)	<30 g/dL	217 (7%)	27.84±2.11
	30-35.5 g/dL	2661 (90%)	32.86±1.20
	>35.5 g/dL	94 (3%)	53.47±16.16
MCH Mean Corpuscular Hemoglobin	<25.1 pg	647 (22%)	21.19±3.21
	25.1-31.6 pg	2027 (68%)	28.32±1.61
	>31.6 pg	298 (10%)	37.7±9.57
RBC (Red Blood Cells) Count	<4.25 x10 ¹² /Lit.	509 (17%)	3.50±0.69
	4.25-6.02 x10 ¹² /Lit.	2261 (76%)	5.04±0.43
	>6.02 x10 ¹² /Lit.	202 (7%)	7.27±1.68

*Complete blood Count

Table No. 3: Descriptive statistics of total population for age and RBC (Red Blood Cells) indices

Parameters Mean±SD	Male	Female
Age (years)	31.57±20.58	31.98±16.92
Hemoglobin (g/dL)	13.34±3.28	11.51±3.76
Hematocrit (%)	40.85±9.25	36.15±8.01
MCH (pg)	27.71±5.71	26.48±5.91
MCHC (g/dL)	33.15±4.99	32.52±5.50
MCV (fL)	84.75±15.49	83.03±16.39
RBC (10 ¹² /Lit) Count	4.93±1.07	4.49±0.97

Table No. 4: Statistical comparison of mean values of RBC indices versus the mean value of standard in both genders

CBC* Parameters	Gender	Mean of Standard ¹⁵	Mean of Sample	p-value**
Hemoglobin (g/dL)	Male	14.45	13.34	p = 0.001
	Female	12.75	11.51	p = 0.0034
Hematocrit (%)	Male	44.55	40.85	p = 0.01
	Female	39.95	36.15	p = 0.015
MCH (pg)	Male	28.35	27.71	p = 0.01
	Female	28.5	26.48	p = 0.022
MCHC (g/dL)	Male	32.75	33.15	p = 0.0001
	Female	32.35	32.52	p = 0.028
MCV (fL)	Male	87.5	84.75	p = 0.001
	Female	86.7	83.03	p = 0.0122
RBC (10 ¹² /Lit)	Male	5.13	4.93	p = 0.039
	Female	4.40	4.49	p = 0.01

*Complete blood Count; **p-value is significant at <0.05

DISCUSSION

Anemia remains a serious worldwide health issue especially in developing countries. The current study focused on the prevalence of anemia, its causative factors and available therapeutic options. According to the criteria set-forth by the largest private sector of JCI (Joint Commission International) accredited tertiary-care hospital of the city; anemia was reported in 39% (N=1984) female and 33% (N=994) male population. This indicates that high prevalence of anemia among females of reproductive age. Studies conducted in different countries revealed that anemia is linked to higher rates of morbidity and mortality,¹⁴ poor birth outcomes,¹⁵ and delays in children's cognitive and behavioral development.¹⁶

Therefore, the World Health Organization recommends a daily 30–60 mg elemental iron supplementation for the women of reproductive age; while for infants and children (6 months to 12 years of age), WHO recommends consumption of fortified foods with folic acid, zinc, vitamin A as multiple micronutrients. Despite overall beneficial effects, there is limited adoption by high risk population.¹⁷ Reasons could be poor socio-economic standing and poverty. Therefore, the 1,000 Days initiative has drawn the attention of numerous countries and efforts are required to increase intake of iron-rich foods.¹⁸

Microcytic, hypochromic anemia is a condition, where RBC size is less than normal and also decreased in red color.¹⁹ In the current study; based upon MCV; microcytic anemia prevailed in both the genders; male 25% (N=739) and female 31% (N=1554), while hypochromic-microcytic anemia was found in 7% (N=217) male and 16% (N=815) female. In case of microcytosis; Iron deficiency anaemia (IDA), thalassemia and anaemia of chronic diseases (ACD) are the three basic diagnostic options.²⁴ The most common type of microcytic anaemia is iron deficiency anaemia.²⁰ Therefore, assessment is necessary by the findings of ferritin levels, serum iron, total iron binding capacity and haemoglobin. Fortunately in the findings of current study; normocytic condition was found in 62% (N=3073) female and 68% (N=2010) male, while normochromic condition was found in 78% (N=3858) female and 90% (N=2661) male. Normochromic-normocytic anemia is basically caused by nutritional deficiencies, renal insufficiency and hemolytic anaemia.²¹ Current study found clinically small number of cases of normochromic-normocytic anemia, that is 8% (N=407) female and 8% (N=243) male.

Megaloblastic or Macrocytic anemia is usually caused by deficiencies of folic-acid and/or vitamin B-12 (Cobalamin); in this situation, usually MCV is >100fL.²² Deficiencies of folic-acid and/or vitamin B-12 may result in ineffective erythropoiesis.²² Megaloblastic anemia in the current study was found in 8% (N=223) male and 7% (N=325) female population. However, nonmegaloblastic macrocytic anemia also occurs due to other causes, such as abuse of ethanol, aplastic anemia, myelo-dysplastic syndrome, liver disease, hypothyroidism and drugs,²² for differential diagnosis; folic acid and vitamin B-12 serum levels should be determined. Sometimes vitamin B-12 deficiency occurs due to positive antibodies to intrinsic factor; which confirms the diagnosis of pernicious anemia.²²

The alarming situation is that; when mean values of RBC indices (Hb, Hct, MCV, MCH, MCHC, RBC count) in both gender were statistically compared with standard; indices were significantly lower than standard (Table No. 4); which concluded that most of the population in Karachi is either anemic or there is a need to develop new standards for the normal ranges of these indices.

Based upon findings of current study; it is highly recommended to follow international guidelines to resolve the issue. According to a recent comprehensive analysis, supplementing with just 10 mg/day of elemental iron can improve the levels of hemoglobin.²³ In another reference; 60 randomized controlled studies involving 27402 women from 30 different nations across all continents; overall risk of low-birth weight newborns among women taking daily iron supplements was reduced; mean birth weight of children whose

mothers took iron during pregnancy was 30.81g higher. Daily iron supplementation of 8.88 gm decreased the risk of maternal anemia at term by 70% and the risk of iron deficiency at term by 57%. Supplementation of zinc with iron salts also tend to reduce anemia.²⁴ The role of vitamin B-12 and/or folic-acid for the treatment of megaloblastic anemia cannot be overlooked; it is found in literatures three injections of 1,000 mcg of vitamin B-12 and 5 mg of folic-acid daily rose Hb from 11.24 to 13.12 g/dL (p=0.001), MCV reduced from 95.50 to 89.64.²⁵ If anemia is induced due to chemotherapy; ethropoetin stimulating hormones (epoetin-alpha, darbepoetin-alpha) are highly recommended.²⁶

CONCLUSION

Evaluation of data of more than 8,000 male and female population reveals that hypochromic-microcytic anemia highly prevailed in the society. Majority of such population is in between 18-28 years old. Mean values of RBC indices in both gender were significantly lower than standard.

Limitations and Recommendations: Since mean values of RBC indices in both gender were significantly lower than standard; it is recommended to address anemia or there is a need to develop new standards for the normal ranges of these indices in this population. To enhance anaemia control and prognosis, currently reported factors should be further investigated to develop preventive as well as treatment strategies according to patient's needs.

Author's Contribution:

Concept & Design of Study: Ayesah Zaib-Un-Nisa
Drafting: Iyad Naeem Muhammad
Data Analysis: Sheikh Abdul Khaliq,
Agha Umer Draz Khan
Revisiting Critically: Ayesah Zaib-Un-Nisa,
Iyad Naeem Muhammad
Final Approval of version: Ayesah Zaib-Un-Nisa

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

Ethical Approval: No.IBCKU-300/2023 dated 23.01.2023

REFERENCES

1. Kim-Mitsuyama S, Soejima H, Yasuda O, Node K, Jinnouchi H, Yamamoto E, et al. Anemia is an independent risk factor for cardiovascular and renal events in hypertensive outpatients with well-controlled blood pressure: a subgroup analysis of the ATTEMPT-CVD randomized trial. *Hyperten Res* 2019;42(6):883-91.
2. McCullough PA. Anemia of cardiorenal syndrome. *Kidney Int Suppl* 2021;11(1):35-45.

3. Zavaleta N, Astete-Robilliard L. Effect of anemia on child development: Long-term consequences. *Rev Peruana De Med Experiment Sal Publ* 2017;34(4):716-22.
4. Mirza FG, Abdul-Kadir R, Breyman C, Fraser IS, Taher A. Impact and management of iron deficiency and iron deficiency anemia in women's health. *Exp Rev Hematol* 2018;11(9):727-36.
5. Mamas MA, Kwok CS, Kontopantelis E, Fryer AA, Buchan I, Bachmann MO, et al. Relationship between anemia and mortality outcomes in a national acute coronary syndrome cohort: insights from the UK myocardial ischemia national audit project registry. *J Am Heart Ass* 2016;5(11):1-19.
6. Stevens GA, Finucane MM, De-Regil LM, Paciorek CJ, Flaxman SR, Branca F, et al. Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995–2011: a systematic analysis of population-representative data. *The Lancet Glob Health* 2013;1(1):16-25.
7. Frisetsma GA. Laboratory Evaluation of Hemostasis. 6th ed. In: Keohane E, Otto CN, Walenga J, editors. St. Louis; Missouri 63043: Elsevier Health Sciences; 2020.p.765-92.
8. Kassebaum NJ, Jasrasaria R, Naghavi M, Wulf SK, Johns N, Lozano R, et al. A systematic analysis of global anemia burden from 1990 to 2010. *Blood. J Am Soc Hemat* 2014;123(5): 615-24.
9. Safiri S, Kolahi AA, Noori M, Nejadghaderi SA, Karamzad N, Bragazzi NL, et al. Burden of anemia and its underlying causes in 204 countries and territories, 1990–2019: results from the Global Burden of Disease Study 2019. *J Hematol Oncol* 2021;14(1):1-16.
10. Chaparro CM, Suchdev PS. Anemia epidemiology, pathophysiology, and etiology in low- and middle-income countries. *Ann N Y Acad Sci* 2019;1450(1):15-31.
11. Pincelli A, Cardoso MA, Malta MB, Johansen IC, Corder RM, Nicolette VC, et al. Low-level *Plasmodium vivax* exposure, maternal antibodies, and anemia in early childhood: Population-based birth cohort study in Amazonian Brazil. *P Neglect Trop Dis* 2021;15(7):01-22.
12. Shrestha B, Dunn L. The Declaration of Helsinki on Medical Research involving Human Subjects: A Review of Seventh Revision. *J Nepal Health Res Counc* 2019;17(45):548-52.
13. John P. Bentley. Sample Size and Power Analysis; Chapter 13. In: Aparasu RR, editor. *Principles of Research Design and Drug Literature Evaluation*. 2nd ed. USA: McGraw-Hill Education;2020.p.139-50.
14. Scott SP, Chen-Edinboro LP, Caulfield LE, Murray-Kolb LE. The impact of anemia on child mortality: an updated review. *Nutrients* 2014; 6(12):5915-32.
15. Haider BA, Olofin I, Wang M, Spiegelman D, Ezzati M, Fawzi WW. Anaemia, prenatal iron use, and risk of adverse pregnancy outcomes: systematic review and meta-analysis. *BMJ* 2013;346(1):01-19.
16. Luo R, Yue A, Zhou H, Shi Y, Zhang L, Martorell R, et al. The effect of a micronutrient powder home fortification program on anemia and cognitive outcomes among young children in rural China: A cluster randomized trial. *BMC Pub Health* 2017;17(1):1-16
17. de Barros SF, Cardoso MA. Adherence to and acceptability of home fortification with vitamins and minerals in children aged 6 to 23 months: a systematic review. *BMC Pub Heal* 2016;16(1): 1-11.
18. Harding KL, Aguayo VM, Namirembe G, Webb P. Determinants of anemia among women and children in Nepal and Pakistan: An analysis of recent national survey data. *Mat Chil Nut* 2018;14(S4):01-13.
19. Salama MA, Kamal MY, Younan DN, Henish GA. Hypochromic microcytic anemia: a clinicopathological cross-sectional study. *Alexan J of Pediatr* 2017;30(1):37-43.
20. Elstrott B, Khan L, Olson S, Raghunathan V, DeLoughery T, Shatzel JJ. The role of iron repletion in adult iron deficiency anemia and other diseases. *Eur J Haematol* 2020;104(3):153-61.
21. Newhall D, Oliver R, Lugthart S. Anaemia: A disease or symptom. *Neth J Med* 2020;78(3): 104-10.
22. Socha DS, DeSouza SI, Flagg A, Sekeres M, Rogers HJ. Severe megaloblastic anemia: Vitamin deficiency and other causes. *Clevel Clin J Med* 2020;87(3):153-64.
23. Casgrain A, Collings R, Harvey LJ, Hooper L, Fairweather-Tait SJ. Effect of iron intake on iron status: a systematic review and meta-analysis of randomized controlled trials. *The Am J Cl Nut* 2012;96(4):768-80.
24. Barffour MA, Hinnouho GM, Kounnavong S, Wessells KR, Ratsavong K, Bounheuang B, et al. Effects of daily zinc, daily multiple micronutrient powder, or therapeutic zinc supplementation for diarrhea prevention on physical growth, anemia, and micronutrient status in rural Laotian children: a randomized controlled trial. *The J Pediatr* 2019;207(1):80-9.
25. Nozari E, Ghavamzadeh S, Razzazian N. The Effect of Vitamin B12 and Folic Acid Supplementation on Serum Homocysteine, Anemia Status and Quality of Life of Patients with Multiple Sclerosis. *Clinic Nut Res* 2019; 8(1):36-45.
26. Zhao F, Wang Y, Liu L, Bian M. Erythropoietin for cancer-associated malignant anemia: a meta-analysis. *Molecul Cl Oncol* 2017;6(6):925-30.

Interdisciplinary Collaboration in Pediatric Dentistry: Challenges and Opportunities – A Narrative Review

Mohammed Ali Habibullah

ABSTRACT

Objective: In navigating the landscape of pediatric dentistry, this review underscores the pivotal significance of interdisciplinary collaboration in ensuring comprehensive well-being for children. Despite challenges like communication barriers, the exploration of opportunities ranging from joint training programs to technological integration reveals promising avenues for improvement. Emphasising the imperative of collaboration for optimal pediatric care, the review advocates for sustained research, policy development, and initiatives to support a holistic approach, ultimately aiming to enhance overall health outcomes in this critical healthcare domain.

Key Words: Challenges, Dentistry, Interdisciplinary, Opportunities, Pediatrics

Citation of article: Habibullah MA. *Interdisciplinary Collaboration in Pediatric Dentistry: Challenges and Opportunities – A Narrative Review.* Med Forum 2024;35(1):109-115. doi:10.60110/medforum.350125.

INTRODUCTION

Pediatric dentistry plays a crucial role in promoting and maintaining the oral health of children from infancy through adolescence. Through the establishment of good oral hygiene habits and preventing dental issues, it contributes to overall well-being and quality of life. Children's oral health is linked to various aspects of their development, including speech, nutrition and social interactions.¹ Untreated dental problems in childhood can have long-term consequences, affecting not only the oral cavity but also impacting systemic health. Recognising the importance of early intervention and specialised care for children is fundamental to ensuring their overall health and a positive dental experience.²

Contemporary healthcare focuses on interdisciplinary collaboration as a means for comprehensive and patient-centric care. The understanding that health and well-being are multifaceted has established that no single discipline manages all aspects of a patient's needs.³ Interdisciplinary collaboration is particularly relevant in pediatric care, where the health of a child is influenced by a complex interplay between biological, social and psychological factors.

Department of Preventive Dentistry, College of Dentistry in Ar Rass, Qassim University, Kingdom of Saudi Arabia.

Correspondence: Dr. Mohammed Ali Habibullah, Assistant Professor, Department of Preventive Dentistry, College of Dentistry in Ar Rass, Qassim University, Kingdom of Saudi Arabia.

Contact No: +966509016584

Email: m.habibullah@qu.edu.sa

Received: November, 2023

Accepted: December, 2023

Printed: January, 2024

This leads to a more holistic approach that addresses the broader spectrum of child's health.⁴

The review aims to shed light on the challenges encountered in the collaboration between pediatric dentistry and other healthcare disciplines. Understanding these challenges is essential for devising strategies to overcome them and for promoting effective teamwork. The review aims to provide insights that can guide policymakers, educators, and healthcare professionals in fostering a collaborative environment optimising patient outcomes. The ultimate goal is promoting a model of care that considers the unique needs of pediatric patient.

METHODS

A comprehensive search was executed in October 2023 across multiple electronic databases, including PubMed, SCOPUS, EMBASE, COCHRANE Library, and Science Direct. The search strategy employed MeSH terms and keywords as "Interdisciplinary," "Collaboration," "Pediatric Dentistry", "Challenges" and "Opportunities". Beyond electronic searches, additional relevant articles were identified through manual searches of cross-references and textbooks. The inclusion criteria were set to encompass articles published in English from November 2000 to October 2023 that aligned with the study's objectives. The process of article selection involved a thorough assessment of the inclusion and exclusion criteria, coupled with a quality evaluation of the studies. Initially, 345 articles were identified. After a preliminary review based on titles and abstracts, 67 articles were selected. Subsequent to a full-text evaluation, 21 articles were ultimately selected for inclusion in the review, fulfilling the study's specified criteria (Figure 1).




Figure No.1: Flowchart showing the step-by-step identification of the studies via databases

A brief overview of the selected articles included in this review is provided in Table 1.

Importance of Interdisciplinary Collaboration in Pediatric Dentistry
Holistic Patient Care

Addressing Oral Health in the Context of Overall Health: Pediatric patients often present with conditions that require collaboration between dental and medical professionals to ensure comprehensive understanding of their health status. Congenital disorders, developmental delays, or chronic illnesses may have oral manifestations that necessitate coordinated care. By

addressing oral health within the broader context of a child's well-being, interdisciplinary collaboration contributes to early detection and intervention, promoting optimal health outcomes.⁵

Impact of Oral Health on Systemic Health in Pediatric Patients: Research increasingly demonstrates the intricate relationship between oral health and systemic health, especially in pediatric populations. Conditions such as periodontal disease have been linked to systemic issues like diabetes, cardiovascular and respiratory problems. Collaborative efforts ensure the treatment of oral conditions considers their broader impact on child's health, promoting not only a healthy smile but also overall well-being.⁶

Comprehensive Treatment Planning
Integration of Dental and Medical Care: Interdisciplinary collaboration allows for the integration of dentistry with medical care in the treatment planning process. Pediatric patients often require care beyond traditional dental procedures. Children with complex medical histories or special healthcare needs benefit from collaborative efforts to create cohesive and patient-centered treatment plans.⁷ In pediatric dentistry this integration leads to early intervention and coordinated care significantly impacting a child's health trajectory.

Improved Outcomes for Pediatric Patients: Synergies between dental and medical expertise contribute to improved outcomes for pediatric patients. Collaborative treatment planning lead to better preventive measures, timely interventions, and overall enhanced patient care experiences. Leveraging the expertise of multiple healthcare professionals, pediatric dentistry can achieve more holistic and patient-focused outcomes as illustrated below (Figure 2), ultimately benefiting the well-being of the child.⁸

Table No.1: Overview of selected articles

Authors	Year/ Country	Area of interest	Conclusion
1.Ferraz NK et al. ²	2014/Brazil	Clinical outcomes of untreated caries in preschool children	Early intervention/specialized care fundamental for positive dental experience.
2.Taha A et al ³	2022/India	Interdisciplinary collaboration of Pediatricians & pedodontists for children's oral health	Emphasized position of pediatricians to deliver preventive oral care.
3.Cozza P et al ⁴	2007/Italy	Mouth breathing, atypical swallow and otitis media.	Demonstrated reduction in nasal resistance and improved cognitive hearing after treatment with Rapid Maxillary Expansion.
4.Jones ML et al ⁵	2012/USA	Interdisciplinary approach on pediatric feeding team for	Elaborated dental hygienist's role in interdisciplinary team.

		CWSHN	
5.Casamassimo PS et al ⁶	2000/USA	The relation between systemic and oral health.	Need for physicians to be competent to identify oral abnormalities of concern affecting health, growth and development of children.
6.Powell V et al ⁷	2012	Integration of Medical and Dental Care and Patient Data	Health information technology (HIT) aid isolated components of healthcare delivery to improve patient safety/quality of care.
7. Crall JJ. Et al ⁸	2005/USA	Development and integration of oral health services for preschool-age children	Early establishment of dental homes, risk assessment, integration of dental/medical care components.
8.Gauger TL et al ⁹	2018/USA	Integrative models between pediatric oral health and primary care providers	Collaboration offers varying services and levels of integration.
9.Winter J et al ¹⁰	2019/Germany	Interdisciplinary prevention model for Early Childhood Caries (ECC)	Interdisciplinary team involving gynecologists, midwives, pediatricians, dentists, social services and health office for ECC prevention.
10.Cote CJ et al ¹¹	2006/USA	Monitoring and management of pediatric patients during/after sedation	Standardizing treatment protocols essential from both dental and medical perspectives
11.Fisher-Owens et al ¹²	2007/USA	Conceptual model derived from population health and social epidemiology designed to analyze children's health.	Multilevel model correlating influence of individual, family and community on oral health outcomes.
12.Townsend et al ¹³	2017/USA	Interdisciplinary approach to prepare general dentists to manage dental trauma	Combined case discussions and hospital trauma call shadowing for dental trauma management training.
13.Edelstein B et al ¹⁵	2006 /USA	Policy implications of children presenting with dental emergencies to US Pediatric Dentistry programs	Demonstrated need for public policies to ensure timely, comprehensive dental care for vulnerable children.
14.Shah S. et al ¹⁶	2018/Pakistan	Pediatric dentistry- Novel evolvments	Dentist's collaboration with other specialists to deliver oral care customized to child's specific needs.
15.Van Malsen J et al ¹⁷	2017/Canada	Early establishment of Dental Homes (DH)	Substantiated the DH mode in improving children's oral health
16.Olson CA et al ¹⁸	2018/USA	Growth/Evolution of pediatric telehealth	Proposed databases to aid development/facilitation of multicenter studies to establish value of telehealth in pediatric care.
17.Vertel N et al ¹⁹	2017/Canada	Dental Services access for Children with Special Health Care Needs (CSHCN)	Importance of early referral to tertiary-care centres for CSHCN when beyond the skill/comfort level of dentist.
18.Olayiwola JN et al ²⁰	2014/USA	Care integration in Community Health Centers	Proposed practical guide for care integration, providing framework for strategy planning.

19.Hlongwa P et al ²¹	2021/South Africa	Inter-professional collaboration (IPC) for cleft lip/palate management.	Strategies for IPC among Cleft Lip/Palate teams for patient-centered approach.
20.Mikołajewska et al ²²	2013/Poland	Interdisciplinary Therapy in Cornelia-de-Lange Syndrome	Interdisciplinary approach to manage Cornelia-de-Lange Syndrome. Need for more clinical research/guidelines.
21.Clark M ²³	2017/USA	Interdisciplinary oral health education curriculum	Discussed Smiles for Life curriculum designed to educate healthcare providers about oral disease. Support integration of oral health and primary care.



Figure No.2: Interdisciplinary Pediatric Dentistry Framework




Figure No.3: Seven Steps of Interdisciplinary Treatment Approach

An interdisciplinary treatment approach requires discrete yet definite approach to achieve successful outcomes (Figure 3)

Challenges in Interdisciplinary Collaboration Communication Barriers

Between Dental and Medical Professionals: One significant challenge in interdisciplinary collaboration is communication barriers between dental and medical professionals. Differences in terminology, communication styles, and professional cultures can impede effective information exchange. This lack of seamless communication may lead to misunderstandings, treatment delays, or overlooking crucial aspects of a patient's health. Bridging this communication gap is essential for fostering a collaborative environment where information flows seamlessly, ensuring that all aspects of child's health, oral and systemic, are considered in the decision-making process.⁹

Amongst Interdisciplinary Team Members: Interdisciplinary care often involves professionals from various fields working together. However, effective collaboration requires effective communication amongst them. Coordinating efforts, sharing insights, and maintaining open lines of communication can be challenging, especially when team members come from diverse backgrounds with differing perspectives on patient care.¹⁰ Establishing clear communication channels and promoting a culture of collaborative openness are key strategies to address this challenge.

Lack of Standardized Protocols

Variation in Treatment Approaches: In pediatric dentistry, the lack of standardised protocols and variation in treatment approaches among professionals can hinder collaborative efforts. Differing training backgrounds /philosophies may lead to divergent approaches to similar cases, potentially causing confusion and inconsistency in patient care. Standardising treatment protocols is essential for creating a unified approach that considers the best practices from both dental and medical perspectives.¹¹ This enhances efficiency, reduce the risk of errors, and

contribute to a cohesive and streamlined patient care process.

Need for Cohesive Guidelines in Pediatric Dental Care
The absence of cohesive guidelines specifically tailored to interdisciplinary pediatric dental care contributes to the challenge of ensuring consistent and evidence-based practices. Developing and implementing comprehensive guidelines and establishing standardised protocols promote a shared understanding of best practices and facilitate smoother collaboration.¹²

Professional Silos

Limited Understanding of Each Discipline's Role: Professional silos, where individuals have limited understanding of roles and expertise of professionals from other disciplines, poses a significant challenge. This lack of awareness results in missed opportunities for collaboration and a fragmented approach to patient care.¹³ Educational initiatives and awareness campaigns that promote a better understanding of each discipline's contributions and capabilities help in fostering a collaborative mindset.

Overcoming Turf Issues for Effective Collaboration: Turf issues, such as professional territorialism or a reluctance to cede responsibilities, can hinder effective collaboration. Overcoming these challenges requires a cultural shift within healthcare settings, emphasising the shared goal of improving patient outcomes over individual professional interests.¹⁴ A collaborative culture involves fostering mutual respect, recognising the value of each discipline's expertise, and encouraging a team-based approach.

Opportunities for Interdisciplinary Collaboration Education and Training

Joint Training Programs for Dental and Medical Professionals: One significant opportunity for enhancing interdisciplinary collaboration lies in the development of joint training sessions. These programs facilitate a shared learning environment allowing professionals insights into each other's expertise, practices, and perspectives.

They can include interdisciplinary coursework, collaborative case studies, and shared clinical experiences, promoting a cohesive approach to patient care from both dental and medical perspective.¹⁵

Promoting Cross-disciplinary Understanding: Beyond formal education, ongoing initiatives can promote cross-disciplinary understanding. Workshops, seminars, and interdisciplinary conferences provide platforms for dental and medical professionals to engage in dialogue, share experiences, and develop a mutual appreciation for each other's roles.¹⁶

Integrated Patient Care Models

Establishing Pediatric Dental Homes: Pediatric dental homes provide a central hub for a child's oral health, where dental professionals coordinate and

integrate care with other healthcare providers. They provide a focal point for preventive care, early intervention, and ongoing management, with dentists collaborating closely with other specialists to address the unique healthcare requirements of pediatric patients.¹⁷

Coordinated Care Delivery through Collaborative Clinics: Collaborative clinics bring together dental and medical professionals in a shared physical space provide a practical opportunity for coordinated care delivery. By offering integrated services, joint consultations, shared treatment planning, and seamless referrals between dental/medical specialists, healthcare professionals enhance communication, streamline workflows improving the overall patient experience.¹¹

Technology Integration

Electronic Health Records for Seamless Information Sharing: Leveraging technology, particularly electronic health records (EHRs), offers a significant opportunity to overcome communication barriers and enhance information sharing. Integrated EHR systems allow dental and medical professionals to access and update patient information in real-time, ensuring all healthcare providers have a comprehensive view of a child's health history.¹⁸

Telehealth and Virtual Platforms for Interdisciplinary Consultations: The integration of telehealth and virtual platforms presents an innovative opportunity for interdisciplinary collaboration. These platforms enable remote consultations, collaborative case discussions, and real-time communication among healthcare professionals, regardless of their physical location.¹⁸ By embracing technology, professionals can overcome geographical barriers, improve accessibility to expertise, and foster interdisciplinary collaboration for routine/specialised care.

CASE STUDIES

Successful Examples of Interdisciplinary Collaboration in Pediatric Dentistry

Specific Programs or Initiatives

a. Children's Hospital Dental Center:

These are renowned for their commitment to providing an interdisciplinary approach to patient care, particularly for children with complex medical conditions or special needs. These centres understand the importance of collaboration between pediatric dentists, paediatricians, as well as other healthcare professionals, to ensure comprehensive and holistic care. Pediatric dentists work closely with paediatricians, nurses, speech therapists, nutritionists, and other specialists to develop a personalised care plan that considers the child's unique medical history and requirements. This collaborative approach ensures any potential interactions between dental treatments and the child's medical conditions or medications are carefully

considered. Additionally, the centres with specialised equipment and facilities, accommodate children with physical disabilities or sensory sensitivities, creating a welcoming and safe environment. They provide the highest quality of care, emphasising both oral health and overall well-being.¹⁹

b. Community Health Centers (CHC)

CHCs play a vital role in providing an interdisciplinary approach to patient care, serving individuals and families in underserved communities. These centres prioritise comprehensive healthcare by integrating medical and dental services under one roof. Patients, particularly children, benefit from this approach as they receive coordinated care to address not only immediate health concerns but also preventive measures and health education.²⁰ By offering a one-stop solution for medical/dental care, CHCs improve access to healthcare services, promote continuity of care, and enhance overall health outcomes for their patients, especially those facing socioeconomic disparities.

c. Collaborative Care for Cleft Lip/Palate Patients

In pediatric dentistry, the interdisciplinary approach to managing cleft lip/palate patients exemplifies collaborative care. Teams comprising pediatric dentists, surgeons, speech therapists, and orthodontists work together to ensure improved surgical, dental, and speech outcomes. This collaboration underscores the necessity of early intervention and continuous, integrated care strategies, tailored to each patient's unique needs.²¹

d. Interdisciplinary Approach for Children with Syndromes

Similarly, in treating children with syndromes such as Down's or Autism Spectrum Disorders, an interdisciplinary team approach is vital. This method not only addresses dental needs but also significantly impacts the overall development and quality of life of these children. This patient-centered, collaborative approach demonstrates the profound benefits of a holistic healthcare model.²²

Enhancing Pediatric Dental Care through Outcome Evaluation and Continuous Improvement

Evaluating Interdisciplinary Collaboration

Outcomes: Outcome evaluation is crucial in assessing the effectiveness of multidisciplinary collaboration in pediatric dental care. Key metrics include patient health outcomes, patient/family satisfaction, and efficiency of care delivery. These metrics, along with feedback from stakeholders, are essential in measuring the success and impact of collaborative efforts.²³

Continuous Improvement in Multidisciplinary Settings:

Continuous improvement is essential for the

advancement of pediatric dental care. Quality improvement initiatives stemming from outcome assessments such as new training programs, process optimisations, and technology integration enable dental care to evolve continually. This approach ensures that pediatric dentistry remains up-to-date with latest best practices and research, leading to enhanced care quality and better patient outcomes.⁸

Future Directions: Research is essential to comprehend the long-term impact of interdisciplinary pediatric dental care on both oral and systemic health outcomes. Knowledge gaps persist regarding the effectiveness of collaborative interventions and their influence on overall patient well-being. Further investigation is needed to identify best practices for communication within interdisciplinary teams and explore innovative technologies facilitating seamless communication and information sharing between dental and medical professionals. Closing these gaps is crucial for advancing evidence-based practices and optimising collaborative care in pediatric dentistry.

POLICY IMPLICATIONS

Advocacy for Integrated Healthcare Policies:

Promoting interdisciplinary collaboration in pediatric dentistry necessitates strategic policy advocacy and support initiatives. Firstly, policies should endorse and incentivise joint training programs, fostering a collaborative mindset from the early stages of education. This approach aims for a shared understanding and appreciation of each discipline's role. Additionally, supporting policies that recognise and integrate oral health into broader healthcare frameworks ensures that oral health is seamlessly woven into the overall fabric of healthcare delivery. Moreover, financial incentives for collaborative clinics are crucial, encouraging the establishment and maintenance of integrated dental and medical services. Furthermore, advocating for research initiatives examining the economic benefits of interdisciplinary collaboration in pediatric dentistry reinforces the importance of collaborative models in not only improving patient outcomes but also overall efficiency and cost-effectiveness of healthcare systems.

CONCLUSION

The review has identified challenges like communication barriers and lack of standardised protocols in interdisciplinary collaboration in pediatric dentistry, while emphasising opportunities in education, integrated care models, and technology. Stressing the vital role of collaboration in providing optimal care for pediatric patients, it calls for continued research, policy development, and initiatives to support interdisciplinary approaches. The potential benefits, including enhanced overall health outcomes in children, underscore the

urgency of prioritising and fostering interdisciplinary collaboration in pediatric dentistry.

Author's Contribution:

Concept & Design of Study: Mohammed Ali
Habibullah
Drafting: Mohammed Ali
Habibullah
Data Analysis: Mohammed Ali
Habibullah
Revisiting Critically: Mohammed Ali
Habibullah
Final Approval of version: Mohammed Ali
Habibullah

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

REFERENCES

- Casamassimo PS, Fields H, McTigue DJ, Nowak AJ. Pediatric dentistry: infancy through adolescence, 5th ed. Elsevier India; 2012 Dec 20.
- Ferraz NK, Nogueira LC, Pinheiro ML, Marques LS, Ramos-Jorge ML, Ramos-Jorge J. Clinical consequences of untreated dental caries and toothache in preschool children. *Pediatr Dentist* 2014;36(5):389-92.
- Taha A, Chakrabarty D. Optimal Oral Health for Children: An Interdisciplinary Approach between Paediatricians and Paedodontists. *J Pediatr* 2022; 8(4.1000155).
- Cozza P, Di Girolamo S, Ballanti F, Panfilio F. Orthodontist-otorhinolaryngologist: an interdisciplinary approach to solve otitis media. *Eur J Paediatr Dent* 2007;8(2):83-8.
- Jones ML, Boyd LD. Interdisciplinary approach to care: the role of the dental hygienist on a pediatric feeding team. *J Allied Health* 2012;41(4):190-7.
- Casamassimo PS. Relationships between oral and systemic health. *Pediatr Clin North Am* 2000; 47(5):1149-57.
- Powell V, Din FM, Acharya A, Torres-Urquidy MH, editors. Integration of medical and dental care and patient data. Springer Science Business Media; 2012 Jan 18.
- Crall JJ. Development and integration of oral health services for preschool-age children. *Pediatr Dentist* 2005;27(4):323-30.
- Gauger TL, Prosser LA, Fontana M, Polverini PJ. Integrative and collaborative care models between pediatric oral health and primary care providers: a scoping review of the literature. *J Public Health Dentist* 2018;78(3):246-56.
- Winter J, Bartsch B, Schütz C, Jablonski-Momeni A, Pieper K. Implementation and evaluation of an interdisciplinary preventive program to prevent early childhood caries. *Clin Oral Investigations* 2019;23:187-97.
- Cote CJ, Wilson S CJ C. Guidelines for monitoring and management of pediatric patients during and after sedation for diagnostic and therapeutic procedures: an update. *Pediatr* 2006; 118: 2587-602.
- Fisher-Owens SA, Gansky SA, Platt LJ, Weintraub JA, Soobader MJ, Bramlett MD, et al. Influences on children's oral health: a conceptual model. *Pediatr* 2007;120(3):e510-20.
- Townsend J, King B, Ballard R, Armbruster P, Sabey K. Interdisciplinary approach to education: preparing general dentists to manage dental trauma. *Dental Traumatol* 2017;33(2):143-8.
- Muthu MS, Sivakumar N. Pediatric dentistry. Elsevier Health Sciences; 2011 Sep 20.
- Edelstein B, Vargas CM, Candelaria D, Vemuri M. Experience and policy implications of children presenting with dental emergencies to US pediatric dentistry training programs. *Pediatr Dentist* 2006; 28(5):431-7.
- Shah S. Paediatric dentistry-novel evolvement. *Annals Med Surg* 2018;25:21-9.
- VanMalsen J, Compton SM. Effectiveness of early pediatric dental homes: A scoping review. *Canadian J Dental Hygiene* 2017;51(1).
- Olson CA, McSwain SD, Curfman AL, Chuo J. The current pediatric telehealth landscape. *Pediatr* 2018;141(3).
- Vertel N, Harrison RL, Campbell KM. Access to dental services for children with special health care needs: A pilot study at the dental department of BC Children's Hospital. *J Can Dent Assoc* 2017;83 (h6):1488-2159.
- Olayiwola JN, Bodenheimer T, Dubé K, Willard-Grace R, Grumbach K. Facilitating care integration in community health centers. UCSF Center for Excellence in Primary Care March 2014.
- Hlongwa P, Rispel LC. Interprofessional collaboration among health professionals in cleft lip and palate treatment and care in the public health sector of South Africa. *Human Resources For Health* 2021;19(1):1-9.
- Mikołajewska E. Interdisciplinary therapy in Cornelia de Lange syndrome—review of the literature. *Adv Clin Exp Med* 2013;22(4):571-7.
- Clark M, Quinonez R, Bowser J, Silk H. Curriculum influence on interdisciplinary oral health education and practice. *J Public Health Dentistry* 2017;77(3):272-82.

C1Q Nephropathy, An Unusual Occurrence in a Middle-Aged South Asian Woman

Zohaib Ramzan¹, Shahid Anwar¹, Syed Ali Raza² and Zoha Majeed³

ABSTRACT

We present an interesting but rare occurrence of C1q nephropathy in a middle-aged South Asian woman with a history of edema and fatigue who responded well to steroids. C1q nephropathy is a glomerulopathy characterized by large amounts of C1q deposits in mesangium and is a diagnosis of exclusion after ruling out SLE, affecting a predominately pediatric population. Our case highlights the importance of lateral thinking while dealing with management and treatment outcomes in C1q nephropathy.

Citation of article: Ramzan Z, Anwar S, Raza SA, Majeed Z. C1Q Nephropathy, An Unusual Occurrence in a Middle-Aged South Asian Woman. Med Forum 2024;35(1):116-118. doi:10.60110/medforum.350126.

BACKGROUND

C1q nephropathy was described by Jennette and Hipp in 1985. It is characterized by large amounts of mesangial Ig and complement deposition with the predominant appearance of C1q after the exclusion of systemic lupus erythematosus and mesangial-proliferative disease.^{1,2} The pathogenesis is unclear. Incidence of C1q nephropathy varies in reports ranging from 0.2 to 16% with no gender differences and appears to be higher in children. Clinical presentation ranges from asymptomatic hematuria or proteinuria to frank nephrotic or nephritic syndrome in children and adults.³ The disease pattern on biopsy may vary, but the core component of diagnosis remains C1q deposition with no features of SLE. Biopsies may range from no lesion in the kidney besides C1q deposition in the mesangium; those with features of FSGS may have associated mesangial proliferation, and mesangial hypercellularity may be seen with those presenting with proliferative glomerulonephritis.⁴

For the most part, though, it is considered steroid-resistant. Those with minimal change disease-like patterns may have greater remission rates, whereas those with FSGS patterns may be more prone to the development of end-stage renal disease.

¹. Department of Nephrology / Medicine², Fatima Jinnah Medical University, Lahore.

³. Department of Medicine, Sir Ganga Ram Hospital, Lahore.

Correspondence: Dr. Zohaib Ramzan, Department of Nephrology, Fatima Jinnah Medical University, Lahore
Contact No: 03330160858
Email: zohaib321@live.com

Received: November, 2023
Accepted: December, 2023
Printed: January, 2024

A large meta-analysis found that partial remission rates were 28% and complete was 49%. Some data shows complete and partial remission with steroids. However, cyclosporine and Cytoxan have also been used with steroids for remission. Relapses may also be common⁵.

CASE PRESENTATION AND DISCUSSION

A 30-year-old female from Lahore came to the Nephrology clinic in May 2020, complaining of generalized body swelling for two weeks associated with fatigue, exertional dyspnea, and epigastric pain. Swelling is more pronounced in her lower limbs; her epigastric pain was mild, radiating, and not associated with nausea and vomiting. She was dyspneic about taking 10 to 20 steps but could carry out her daily activities without discomfort. All the symptoms were concordant in time. Her past medical and surgical history was unremarkable. Her family history and personal history are also unremarkable.

Her bp was 140/90 at the presentation, and the rest of the vitals were normal. A general physical examination showed periorbital puffiness and 2+ bilateral pitting edema up to the knees. On Respiratory examination, there was bilateral decreased air entry with normal vesicular sounds and a respiratory rate of 22 after minimal exertion.


The rest of her systemic examination was normal. Ultrasound of the abdomen showed splenomegaly with a spleen size of 12.0cm. The splenic index was 60.

ECG at presentation was normal with no acute changes, and chest x-ray showed bilateral basal infiltrates, most likely pulmonary edema. (Figure 1)

At this stage, our differential diagnoses included the following:

- Membranous glomerulopathy
- Lupus nephritis
- Ascending urinary tract infection

Symptomatic treatment was begun at this stage with further directed investigations.



The following results of investigations done in the first week of June 2020 were found:

Investigation	Result	Normal Range
Complete Urine Examination		
a) Color	Yellow	
b) Specific gravity	1.030	
c) pH	5.0	
d) glucose	nil	
e) Ketones	nil	
f) Proteins	++	
g) Hemoglobin	+++	
h) Bilirubin	nil	
i) Nitrite	negative	
j) Leucocyte esterase	nil	
k) Pus cells	8-10	
l) RBCs	25-30	
m) Epithelial cells	12-15	
n) Casts	nil	
o) Organisms	nil	
p) Yeast	nil	
Spot urinary to creatinine ratio	3.18 mg/mg	<0.20 mg/mg
Spot Urinary Protein	99.7 mg/dL	<14.0 mg/dL
Spot urinary creatinine	31.4 mg/dL	
Serum urea	22mg/dL	10-50mg/dL
Serum Creatinine	1.2mg/dL	0.6-1.4mg/dL
Serum Sodium	129 mmol/dL	135-150mmol/dL
Serum Potassium	3.2 mmol/dL	3.5-4.5mmol/dL
Serum uric acid	4.4 mg/dL	2.6-6.0mg/dL
Lipid profile		
Cholesterol	157 mg/dL	<200mg/dL
Triglycerides	93mg/dL	<150mg/dL
ESR	23mm	<20mm
Hepatitis B Screening	Negative	
Hepatitis C screening	Negative	
Autoimmune profile		Pattern: Fine cytoplasmic speckled appearance associated with Anti synthetase syndrome,
a) ANA	Positive	
b) Pattern	Fine cytoplasmic Speckled	
c) Estimated endpoint titer	1/160	

d) ASMA	Negative	polymyositis, dermatomyositis, limited systemic sclerosis DsDNA (<20.0 IU/ml)
e) AMA	Negative	
f) dsDNA	5.6 IU/ml	
g) RA Factor	Negative	
Serum C3 Levels	0.4 g/L	0.8-1.6 g/L
Serum C4 Levels	0.24 g/L	0.1-0.4 g/L

Ultrasound KUB: Right and left kidney size normal. The bladder is normal. Bilateral pleural effusion, right-sided 40ml, left-sided 20ml.

Based on these investigations and her clinical picture, lupus nephritis was suspected. A renal biopsy was planned. A renal biopsy was scheduled for the 15th of June 2020.

In the interval week till the biopsy, she developed a generalized morbilliform rash associated with itching, initially thought to be perhaps a drug eruption. She was prescribed Kestine 10mg HS, and the rash settled in two days. No inciting factors were recognized. A CBC done showed platelets of 93×10^9 . However, the rash was not purpuric or petechial; thus, urticaria was suspected due to the sudden nature and prompt response to antihistamines.

A renal biopsy was done, and the histopathology report is as follows:

Sections revealed the core of renal tissue consisting of cortical regions containing up to 10 glomeruli in a serial section. The glomeruli show increased cellularity with mild mesangial proliferation. Occasional segments show increased endothelial cells with few polymorphs. The glomerular capillary wall appears unremarkable. Few tubules show focal mild tubular atrophy. Occasionally, they have amorphous casts in lumina. Interstitium reveals a patchy sprinkling of mononuclear cells and eosinophils. Blood vessels appear unremarkable.

Her next follow-up was a month later. Clinically, her pedal edema and hypertension had resolved. Her dyspnea had improved, as had her fatigue. Based on her lab investigations, she had responded to the steroid therapy, and her lab parameters and clinical parameters had normalized to a great extent. The leucocytosis was secondary to the steroid therapy, and she had no signs or symptoms of any ongoing infection. A repeat chest x-ray done at this time also showed no pulmonary edema, and her furosemide was stopped.

Her next follow-up was in August 2020, during which she was clinically doing well with no complaints. Her lab parameters that were routinely done came back within the normal range, including her protein-to-creatinine ratio, which had further decreased significantly from 0.21 mg/mg to 0.11 mg/mg, which was now well within the normal range. She has received steroids now for 6 weeks, and it was tapered by 10mg per week. And she will be maintained on 10mg prednisolone daily once tapering is complete over

4 months. Urinary protein to creatinine ratio monitoring will continue. As we saw, though the most common presentation in the series published from Pakistan was minimal change disease, our patient, though presenting with nephrotic syndrome, had a pattern of proliferative glomerulonephritis. Beyond patterns, patterns of age groups affected by the disease in the South Asian population may be different and more centered around adults, and possibly responsiveness to therapy might also be different. In conclusion, larger reviews and studies from the communities' native countries would prove extremely beneficial in piecing together the jigsaw.

Author's Contribution:

Concept & Design of Study:	Zohaib Ramzan
Drafting:	Shahid Anwar
Data Analysis:	Syed Ali Raza, Zoha Majeed
Revisiting Critically:	Zohaib Ramzan, Shahid Anwar
Final Approval of version:	Zohaib Ramzan

Conflict of Interest: The study has no conflict of interest to declare by any author.

Source of Funding: None

REFERENCES

1. Miller R, Unda SR, Holland R, Altschul DJ. Western moyamoya phenotype: a scoping review. *Cureus* 2021;13(11).
2. Ramponi G, Folci M, Badalamenti S, Angelini C, Brunetta E. Biomarkers and diagnostic testing for renal disease in Sjogren's syndrome. *Frontiers Immunol* 2020;11:562101.
3. Bridoux F, Ferman JP, Leung N, Ronco P, Touchard G. Monoclonal Gammopathy Of Renal/Clinical Significance. *Comprehensive Clinical Nephrology: Comprehensive Clinical Nephrology-E-Book* 2023.p.23.
4. Howard C, Sheridan J, Picca L, Munir W, Meharban N, Karthik P, et al. Autoimmunity and its correlation to inflammatory vascular diseases. In *Translational Autoimmunity*. Academic Press: 2022.p.189-228.
5. Glomerulopathy F, Glomerulopathy I, Glomerulopathy CTIF. *Diseases With Organized Deposits. Diagnostic Pathology: Kidney Diseases E-Book*; 2023.p.302.

Author Index January to December 2023**Azhar Masud Bhatti**

Editor in Chief

Vol. 34, No. 1, January, 2023

Author (s)	Page No.
1. Jan MM.	1
2. Alruways MW	2
3. Nawaz E, Haq G. Deeba F, Nazir N, Rais A, Chachar S	8
4. Afzal A, Alam J, Daniyal RM, Rahat N, Samreena, Akhtar Z	12
5. Mushtaq S, Arshad S, Mushtaq L, Masood H	16
6. Shoaib M, Ch TS, Kamran M, Nazir M, Mehmood K, Mahmood S	22
7. Akhtar N, Zainab S, Sualeh Z, Arif H, Sualeh MA, Abeer SJ	25
8. Abrar SK, Siddiqui S, Aslam K, Hafeez N, Sukhia HR, Ahmed M	29
9. Munir A, Ali K, Khattak MB, Khan MA, Hashmat F	33
10. Omer Z, Kahloon OI, Khan MH, Muhammad AI, Arshad F, Iqbal MJ	37
11. Zahid MM, Farouk K, Ahmad K, Ali L, Javed HM, Ch MA	43
12. Ashraf N, Rana I, Saeedullah M, Yaqoob B	47
13. Zia D, Rehman SS, Hussain S, Mateen A, Khan MI, Munir MW	52
14. Rehman SS, Khan SH, Zia D, Mateen A, Basharat A, Munir MW	56
15. Janjua HB, Farooq A, Haroon S, Sheikh TK, Hashmi H, Rehman SS	60
16. Siddiqui DK, Tahir M, Ali M, Altaf R, Mustafa K, Siddiqui KK	64
17. Rana MM, Akhtar MS, Awan ZL, Gani D, Malik A, Bilal M	69
18. Rasheed S, Fazlani R, Shaikh S, Irshad S, Khan MP, Memon S	73
19. Khanam S, Abdullahi KO, Sarwar H, Altaf M	78
20. Mirza D, Mazahar S, Marath M, Tariq U, Sohail A, Tanveer F	83
21. Ashraf NN, Siyal NA, Ashraf ES	87
22. Iqbal M, Khan S, Danish Z, Salim I, Ashfaq HM, Raza M	91
23. Rasool G, Yaqoob A, Sawar H	96

Vol. 34, No. 2, February, 2023

Author (s)	Page No.
1. Jan MM.	1
2. Ashraf N, Saeedullah M, Yaqoob B	2
3. Memon ZR, Shaikh FD, Bijarani SA, Memon KN, Soomro F, Memon TF	7

4. Ali S, Khan MA, Munir A, Afridi MA, Siddiqi WA, Ali K	11
5. Sattar S, Sultana S, Shadab W, Afzal S, Salma U, Mobeen A	15
6. Kashif S, Khoso BK, Zeeshan F, Khalid M, Hussain W, Fakhur Uddin	20
7. Shahzad M, Jamil MN, Islam E, Ashraf H	25
8. Khan MS, Rahman E, Raza M, Azim A, Atta Ur Rahman	30
9. Rehman SS, Majeed MA, Mumtaz Y, Khan SH, Hussain S, Maqsood J	35
10. Darraj AQ,	40
11. Azhar MS, Soomra AB, Jamal MU, Naveed S, Kumari S, Khursheed R	46
12. Saddique A, Masood R, Javed R	50
13. Masood R, Saddique A, Javed R	54
14. Islam N, Khan MZH, Khan MZ, Khan HSR, Shaukat I	58
15. Shah MH, Kanval N, Naeem N, Mahmood RK, Raza AA, Naz A	62
16. Kumari S, Alam J, Naveed S, Nazir S, Fatima K, Alam S	66
17. Hussain I, Khan L, Hussain S, Riaz, Shahid AL, Naseem Ullah	71
18. Aslam A, Rehman SS, Maqsood J, Ghafoor A, Samina, Bacha F	76
19. Aghani N, Khaleeq N, Siddiqui SJ, Shaikh H, Memon M, Memon M	80
20. Habib KJ, Mansoori MS, Azim A	84
21. Mansoori MS, Habib KJ, Azim A, Haider S	89
22. kanhar IA, Ahsan S, Nazir S, Tariq S, Akbar T	94
23. Tabassum Z, Masih S, Afzal M, Khan S	98

Vol. 34, No. 3, March, 2023

Author (s)	Page No.
1. Jan MM.	1
2. Darraj AQ	2
3. Fatima M, Ain QT, Naz A, Mahmood RK, Ibrahim M, Shah MH	8
4. Rahim S, Nasir K, Qureshi R, Dhrolia MF, Ahmad A	12
5. Himayat Ullah, Mustafa G	17
6. Khoso BK, Samdani AJ	22
7. Madiha, Kalhor FA, Waqas, Tariq, Memon L, Zaighum M	27
8. Enver N, Hamid W, Ilyas M, Azeem M, Rehman A	32
9. Waheed A, Talpur GA, Qadri NA, Qayyum SA, Ram H, Khan IA	35

10. Qazi AF, Singha SP, Kazi M, Afzal K, Masood SH, Atif A	40
11. Jamil MN, Asif M, Islam E, Ashraf H, Haq F, Shahzad M	45
12. Khattak MB, Shad H, Munir A, Khan MA, Afridi MA, Ali K	49
13. Khan A, Faisal D, Khan N	53
14. Rana MM, Akhtar MS, Haq MI, Gani D, Khalid N, Malik A	58
15. Rehman SS, Hamdani SHH, Hussain S, Munir MW, Mumtaz Y, Sarwar R	63
16. Aftab U, Zaighum M, Darshana, Khan TA, Khan R, Ramzan S	67
17. Mangi AH, Shahzad G, Jatoi A	71
18. Gul U, Hashmi SFA, Kadri S, Anwari M	75
19. Khan A, Quratulain, Pervez H, Akhtar S, Inayatullah, Shah F	80
20. Ahmed SN, Qanash S, Softah A, Alsolami H, Bakhsh T, Alzahrani O, et al	85
21. Gemnani VK, Juseja AK, Nagdev DR, Abro K, Jokhio SA, Jamro FS	92
22. Saleem I, Afzal M, Sarwar H, Khan S	98
23. Rafique A, Sarwer H, Ali A	102

Vol. 34, No. 4, April, 2023

Author (s)	Page No.
1. Jan MM.	1
2. Thangam MMN	2
3. Choudhary SA, Rehman SS, Sarwar R, Zia D, Khan SH, Fatima N	7
4. Akhtar MS, Arooj A, Abbas MR	12
5. Ghias Z, Dildar S, Jaffri SA, Afzal N	15
6. Ahmed A, Tamkeen N, Khan MR, Wahid G, Fahim S, Badshah U	19
7. Dildar S, Khan AH, Jaffri SA, Sultan S, Islam N	22
8. Khalid M, Asif M, Aslam U, Hassan MH, Ahmed B, Ali E	25
9. Haq QM, Aaqil B, Nazneen Z, Siddiqi A, Faisal Z, Salam Z	30
10. Ashraf NN, Khokar AR, Ashraf ES	33
11. Mustafa G	36
12. Ramzan M, Shehzad A, Razzaq A, Javed K, Jahanzaib, Saeed A	41
13. Sami Ullah M, Ahmed S, Naeem N, Shah MH, Naz A, Mahmood RK	46
14. Afshan Z, Musharraf H, Aslam K, Sukhia HR, Siddiqui SD, Abrar SK	50
15. Bhojwani SL, Soomro N, Shahani FA, Gemnani VK, Abro K, Jokhio SA	54
16. Enver N, Shaheen A, Rashid F, Azeem M, Rehman A, Qadir S	58
17. Sunil RK, Siddique T, Kumari S, Khan T	62
18. Anwar U, Jamil B, Khan NW, Raza M, Sajjad M, Jamil J	67
19. Zeb F, Pervez S, Kaleemullah M, Wahab A, Khan B, Salih N	70

20. Pervez S, Zeb F, Khan A, Kaleemullah M, Wahab A, Khan B	
21. Mangi AH, Shahzad G, Ahmed A	75
22. Ahmad S, Iqbal J, Zafar S, Aslam J, Zahid H, Rahman W	80
23. Shamim S, Arif MS, Iftikhar N, Asif N, Jhangir U, Arif O	85
24. Wassan SM, Masood SH, Hasan SM, Memon NA, Irfan S, Manzoor W	90

Vol. 34, No. 5, May, 2023

Author (s)	Page No.
1. Jan MM.	1
2. Rana MM, Nagra SR, Awan ZL, Akhtar MS, Rauf Q, Afzal MW	2
3. Zahoor S, Majeed M, Ch. AN, Rashid N, Fareed S, Mustafa N	6
4. Asad H, Mehmood H, Hussain S, Hussain R, Arshed M, Ali I	7
5. Rafiq S, Bashir H, Akhtar A, Karim S, Mumtaz A, Anjum F	11
6. Jawaid W, Shafee SM, Singh J, Irfan M, Sheikh H, Nisa Q	15
7. Naeem U, Farooq F, Awan MN, Salim S, Iftikhar J, Mirza M	20
8. Hussain S, Afridi KK, Aslam V, Nasir M, Ahmad F	24
9. Khan SA, Rasheeq T, Khan MSN, Raza A, Mirbahar AM, Tahir M	28
10. Hammad AR, Ejaz H, Faruqi MS	33
11. Bashir N, Yasmeen S, Aftab A, Zafar JDA, Shaukat S	36
12. Anum M, Khan MS, Butt AI, Lone MA, Atif M, Hanif S	40
13. Mehtab B, Jabeen B, Aslam K, Bajwa WN, Kumar B, Lone MA	44
14. Dildar S, Asif M, Jaffri SA, Imam M, Shamsi TS	48
15. Essa MA, Gul y, Inamullah, Farooq MZ, Shafaat Ullah	51
16. Farooq MZ, Gul Y, Essa MA, Inamullah, Shafaat Ullah	56
17. Inamullah, Ahmad H, Essa MA, Farooq MZ, Shafaat Ullah, Gul Y	63
18. Anwar ul Haq, Nawab K, Khan N, Shoaib I, Gul H, Khattak MA	68
19. Afridi F, Nawab K, Haq A, Gul H, Khan N, Shoaib I	72
20. Nawab K, Khan N, Shoaib I, Haq A, Khattak MA, Gul H	76
21. Ahmad N, Saeed M, Amjad S, Rasheed A, Iftikhar U, Ahmad M	79
22. Zakir H, Manzoor W, Wassan SM, Hasan SM, Ali HA, Zehra AH	84
23. Raza SMM, Wassan SM, Ahmed MA, Kumari K, Kumar H, Naqvi AM	88

24. Korai SM, Kazi M, Qazi AF, Sughra S, Zahid S, Theba FK	93	19. Rehman M, Mian FA, Ali A, Raza MA, Niaz MA, Humayun S	78
25. Shafee SM, Jawaid W, Barry SJ, Singh J, Zaheer M, Shahbaz NN	97	20. Saleem MW, Hashmat S, Fauzia T, Zulfiqar M, Ali BA, Memon MA	82
26. Raza U, Khanam A, Iftikhar A, Rizwan S, Gauhar TM, Zehra AH	101	21. Fauzia T, Saleem MW, Zulfiqar M, Hashmat S, Memon MA, Ali BA	86
27. Mustafa G, Himayat Ullah, Banday AH, Alwadai AA, Alotaibi MS, Aljameeli KS	105	22. Akhter J, Abbas MR, Akhter MS	90
28. Mirza U, Naheed A, Naveed T, Ahmed A, Zehra T, Nisar N	111	23. Munawar S, Zafar M, Azhar M, Ali M, Sadia, Ali H	94
29. Mahar Y, Mehwish A, Sarwat S, Khokhar SK, Qamar A, Khan M	115	24. Afridi RAK, Asif M, Khan KN, Anwar K, Rehman S	98
30. Siraj F, Rahat N, Shahzad H, Anwar M, Jalbani A, Momin Z	119	25. Rana MM, Gani D, Khalid N, Afzal A, Rauf Q, Afzal MW	102
31. Ahmed A, Sultan A, Jamil B, Luxmi S, Khan T	124	26. Mangi AH, Memon MM, Shahzad G, Khan F, Khan S, Shahid Ullah	107
32. Shahid A, Shahid A, Jaleel R, Siddiqui SA, Shamaun S, Aslam M	129	27. Bhutta MR, Sheikh AH, Siddiq M, Mustafa S	112
33. Uqaili AAM, Uqaili R, Shaikh SN, Siddiqui K, Khan HS, Tagar M	135	28. Sheikh AH, Bhutta MR, Mustafa S	116
34. Kahloon OI, Omer Z, Khan MH, Riaz HB, Muhammad AI	139	29. Bilal M, Aslam V, Karim T, Zaidi A, Jan WA, Ayaz M	120
		30. Siddiqui RA, Alam J, Ram H, Talpur MGA, Khan FA, Muzzamil A	123
		31. Aslam V, Bilal M, Jan WA, Ayaz M, Karim T, Zaidi A	127
		32. Naheed A, Naveed T, Ahmed A, Nisar N, Bibi A, Hafeez J	130
		33. Memon MI, Ahmed SZ, Kumar M, Bai R, Ashfaq A, Rizvi SFA	134
		34. Khan N, Ahmed Z, Waqar F, Saeed W, Batool F, Zafar M	139

Vol. 34, No. 6, June, 2023

Author (s)	Page No.
1. Jan MM.	1
2. Qadir A, Zareen Ullah, Khalil MD, Saqib MN, Khan A, Khan A	2
3. Yaqub K, Zaigham AM, Talat H, Sahaf HA, Dar DN, Afzal M	6
4. Batool U, Ain Q, William GP, Sohail S, Umer H, Jawad F	10
5. Amir E, Anwar M, Latif AQ, Zufishan S, Arshad J, Amir D	14
6. Mustafa G, Himayat Ullah	19
7. Mushtaq S, Aziz R, Sharif A, Masood H	24
8. Khan S, Rafiq S, Sami Ullah M	28
9. Manzar S, Sohail A, Munir MF, Qureshi MM, Chishti FDA, Manzoor HMJ	32
10. Khan KR, Rasool A, Rashid M, Gillani HR, Umair M, Murtaza G	36
11. Arshad T, Chand R, Khan MA, Khokhar RA, Kumari V, Shaikh AS	40
12. Khan MF, Pathan IH, Nek N, Parkash J, Somroo AA, Muhammad H	44
13. Kausar S, Mehkari S, Qader A, Ali Z, Waqar MF, Naveed S	49
14. Kumari P, Syed SA, Atif S, Aslam K, Syed FA, Bibi T	54
15. Tunio R, Baloch D, Gemnani VK, Baloch B, Shaikh RA, Jamro S	59
16. Kumar A, Juseja AK, Jat MI, Gemnani VK, Wadhvani AK, Shaikh MZ	64
17. Mushtaq S, Saeed A, Yaseen A, Khalid S, Kamran R, Arsalan S	68
18. Asghar MR, Akhter T, Asghar A	73

Vol. 34, No. 7, July, 2023

Author (s)	Page No.
1. Jan MM.	1
2. Zafar M, Munawar S, Azhar M, Ali M, Sadia, Ali H	2
3. Shoukat S, Jaddon M, Faryal U, Saqib J	6
4. Azeem Y, Naeem U, Saleem A, Awan MN, Farooq F, Gondal SS	10
5. Tariq A, Mehmood S, Zahid MM, Arshad I	14
6. Zahid MM, Mehmood S, Tariq A, Ahmed K, Arshad I, Naz T	17
7. Amir E, Latif AQ, Fahmi S, Fahmi MS, Zufishan S, Khan E	21
8. Riaz A, Mehwish R, Shafaq M, Hameed S, Raziq H, Rauf A	26
9. Sarwat S, Rehman A, Jabeen S, Mubeen S, Pario A, Marvi N	30
10. Zia Q, Ghous M, Sarwat S, Rehman A, Ahmed MA, Ahmed MR	35
11. Khan N, Haque M, Raja IM, Jamal SA, Zufishan S, Minhas F	40
12. Siddiqui SA, Hasnain FS, Hasan A, Habib A, Jaleel R	45
13. Naeem N, Taqi I, Shamsi FB, Saeed A, Alam A	50
14. Mehwish R, Riaz A, Hameed S, Raziq H, Asghar B, Abid T	55

15. Fatima P, Nadeem MS, Atta S, Khalid FH, Afzal M, Ali R	59	45. Manzoor A, Batool ST, Ahmed T, Khalid J	188
16. Shafique MA, Waqeel N, Qayyum F, Haq MI, Balouch MOK, Kashif M	63	46. Gul S, Sultan S, Fareed A	193
17. Habib F, Sheikh AK, Iltaf S, Khadim S	68	47. Fatima SS, Sultan S, Fareed A	197
18. Nawaz A, Hasham MA, Arshad R, Moinudin IA, Kashif M, Nazir M	72	48. Sultan S, Fatima SS, Fareed A	202
19. Rasheed M, Talib M, Rasheed S, Naz S, Orakzai GS, Naeem N	77	49. Shoaib M, Langrial RZ, Hussain W, Siddique HMAB, Zafar I, Khan S	207
20. Anis R, Shafqat A, Mahmood R, Ahmed S, Khan W, Khan RT	82	50. Alam W, Azam F, Haq N	212
21. Yousaf M, Naz I, Khattak SA, Khan A, Khattak JA	86	51. Azam F, Alam W, Haq N	216
22. Rasool A, Khan KR, Rashid M, Gillani HR, Sattar S, Lodhi TA	91	Vol. 34, No. 8, August, 2023	
23. Achakzai H, Fahim Ullah, Kundi R, Khan H, Ahmed N, Imran S	95	Author (s)	Page No.
24. Haseeb M, Ahsen MI, Azeem M, Khan ZA, Javed S, Qureshi F	99	1. Adil MI.	1
25. Azeem M, Haseeb M, Khan ZA, Qureshi F, Ahsen MI, Dayar J	103	2. Munawar H, 2. Zahid MM, Ahmad K, Ahmed R, Javed HM, Mahmood S	2
26. Cheema SS, Tauseef A, Fatima M, Iqbal MD, Khurshid S, Cheema SR	107	3. Mahmood S, Zahid MM, Tariq A, Ahmed R, Ali I, Ahmad K	7
27. Cheema SR, Tauseef A, Fatima M, Iqbal MD, Rehman R, Cheema SS	112	4. Raza G, Jaffri MSA, Zehra S	11
28. Khan A, Faisal D, Khan B	116	5. Khan M, Khan M, Afridi M, Riaz I, Askar S, Sikander M	15
29. Zareen Ullah, Qadir A, Saqib MN, Khan A, Mufti ND, Sohail MW	120	6. Khalid A	18
30. Imran S, Haq U, Haq I, Ali M, Azam MF, Mahjabeen	124	7. Afzal Z, Khan A, Zeb H, Malik WM, Nasir ZA, Rahman U	21
31. Haq I, Imran S, Haq U, Azam MF, Sabih V, Ali M	128	8. Ramzan M, Sheikh AH, Mustafa S	24
32. Haq N, Akram Ullah, Haq MI, Khan WN	133	9. Sultana T, Akbar J, Khan AH, Saeed T, Mehdi SZ, Jalal MS	28
33. Manzoor R, Ali QA, Shafique MD, Mushtaq B	137	10. Hussain S, Afridi KK, Aslam V, Nasir M, Ahmad F	33
34. Haq E, Hameed A, Riaz G, Ahmad H, Khan S, Bibi R	142	11. Sameed QA, Ali F, Farooq L, Mujahid S, Khalid SS, Zafar U	38
35. Ali QA, Manzoor R, Mushtaq B, Shafique MK	146	12. Shaikh M, Kainat R, Rehman M, Anjum M, Azhar S, Johar SQ	42
36. Rahman A, Zia M, Jadoon AH, Basit A, Ali SB, Shafiq M	150	13. Kainat R, Aslam K, Johar SQ, Sohail R, Rehman M, Fatima SM	47
37. Jalbani A, Rahat N, Shazad H, Bashir P, Siraj F	154	14. Muqet MA, Jabeen W, Muddassar M, Rafiq A, Hassan F, Siddique S	52
38. Sheikh MA, Khosa GK, Rubab S, Hameed Z	158	15. Malik S, Memon AI, Kumar H, Bhatti AM, Zeb M, Gul S	57
39. Raja IM, Khan MA, Haque M, Ahmed T, Rehman A, Khan N	162	16. Noor N, Baseer A, Khan MH	62
40. Javed M, Riaz L, Tariq A, Faryad N, Saleem R, Rizwan S	166	17. Baseer A, Noor N, Badshah Y	67
41. Wajahatullah M, Khan IA, Sohail F, Soomro E, Aakash, Sikandar M	170	18. Afzal S, Ghous MH, Arooj M	72
42. Bhutto AR, Abbasi A, Rafi S, Rehman K, Akhtar ST, Haroon MH	174	19. Ghous MH, Afzal S, Arooj M	77
43. Shoaib M, Batool SM, Munir R, Langrial RZ, Naz S, Sajid M	179	20. Khan D, Haq G, Nazir N, Farhan A, Tariq S, Baqai H	81
44. Cheema HMKN, Zahra SH, Amin I, Cheema UE, Zafar S, Haider HRD	184	21. Ch SR, Iqbal K, Saeed M, Ilyas J, Ashraf A, Rehman F	86
		22. Siddique S, Muddassar M, Haroon A, Rafiq A, Rabbani M, Jabeen T	90
		23. Ashraf S, Ilyas H, Akhter Z	94
		24. Khan S, Jabbar MN, Nawaz A, Saif A, Waqas A	98
		25. Hamna S, Jahan M, Hussain W	102
		26. Naqvi MA, Hashmi A, Fatima S, Naz A, Kousar R, Ibrahim M	106
		27. Hussain A, Ahmed B, Ahmed A, Bhutto SR, Khoso M, Bashir N	110

28. Tariq A, Rashid Z, Yasir S, Narjis ST, Bhatti ZI, Akram M	114	2. Jami A, Ahmed MK, Baig MR, Lodhi MS, Khaliq SA, Kashif SM	2
29. Haq N, Rizwan, Khan M, Khan BR	119	3. Ghafoor MB, Khan SM, Sarwar F, Sultan H, Shafique M, Hassan H	7
30. Ali S, Rahim Z, Mannan R, Farooq A, Babbar S, Khattak K	123	4. Mahmood S, Ismail MO, Ali A, Ahmed J, Afridi FI, Arif H	11
31. Jan N, Imran Ullah M, Anjum S, Islam F, Marwat AG, Hayat I	128	5. Khan MS, Ubaid B, Hashim L, Alam A, Raza M	16
32. Kashif M, Ahmad W, Islam F, Marwat AG, Ahmad I, Saeed L	132	6. Khan A, Khan B, Jameel A	20
33. Ali L, Dayan M, Shahzad M, Hayat F, Qudratullah, Ihsan Ullah	136	7. Rehman S, Gul H, Khizar A, Inayat Ullah	24
34. Aslam V, Bilal M, Ayaz M, Zaidi A, Gul L, Hasan S	140	8. Babar H, Mughal Z, Ashraf S, Ayub N, Arif H, Akhtar R	28
35. Raja AA, Amjad A, Lodhi S, Inam T, Mehmood M, Mumal S	144	9. Fatima A, Gohar H, Siddiqui HZ, Mehdi RF, Naseem S, Khan FZ	32
36. Arooj M, Irshad G, Mukhtar K, Leghari S, Mastoor K, Malik KA	148	10. Shah MA, Nizamuddin, Subhan Z, Khawaja TM, Mehmood I, Sami Ullah	37
37. Mehtab B, Aslam K, Siddiqui SD, Abrar SK	153	11. Mehtab K, Naseem R, Ali SR, Anees M, Haider A, Shah M	41
38. Nisa S, Sundas K, Anwar N, Aziz R	158	12. Liaquat A, Ismail K, Usman S, Gul J, Shah R, Ismail MO	46
39. Rehman W, Nasreen S, Rehman I, Khan MAR, Qayyum F, Aslam J	163	13. Ahmed SN, Alansari B, Alkurishy AR, Algethami A, Alahmari N, Sulami H, Nadeem ZA	51
40. Yusuf A, Sultana N, Sarfaraz T, Fatima A, Ahmed S, Qureshi AM	167	14. Ali S, Aslam M, Nazir MI	56
41. Sarfaraz T, Yusuf A, Sajid S, Fatima A, Ahmed S, Qureshi AM	172	15. Naheed A, Rafay A, Sultan M, Zainab H, Manzoor N, Shahab SA	61
42. Kulsoom U, Nasrullah FD, Jaleel R, Hasnain FS, Shaikh S, Aziz S	176	16. Ahmad H, Siddiqi H, Naheed A, Yousaf MH, Nazeer MM, Iftikhar A	66
43. Khaghan BS, Akbar I, Maaz Ullah, Khan A, Khan O, Hussain SA	181	17. Ahmad F, Alshahrani AM, Mustafa G, Hussain AMB	70
44. Hussain SA, Khaghan BS, Iqbal T, Maaz Ullah, Khan A, Khan O	184	18. Bashir N, Permanand, Hussain I, Arshad S, Khoso M, Hussain A	75
45. Rahman S, Usman M, Syed A	188	19. Ghous MH, Afzal S, Arooj M	79
46. Roghani AS, Khan JUA, Alam I, Wahab A, Iffat S, Younas S	193	20. Baseer A, Badshah Y, Khan MH	84
47. Riaz S, Khan AR, Faizullah F, Alam Z, Manan A	199	21. Ali F, Malik WR, Rehman A, Hashmi MU	88
48. Khan JUA, Iftikhar A, Khalil SK, Fayaz H, Razaq S, Aziz Q	204	22. Taj A, Gul A, Jawed O, Ejaz S, Ahmed SSI, Shaikh N	92
49. Saad M, Javed H, Kiani SN, Khan NB, Shahid F, Sheikh S	208	23. Ali F, Shabbir S, Murad T, Sarfaraz A, Shafay M, Yasmin N	96
50. Shah SN, Haq N, Khan WN, Hassan N	212	24. Asif S, Navaid S, Gul M, Butt A, Aslam A, Ahmad S	100
51. Ali H, Mushtaq M, Alam W, Haider A	216	25. Hammad M, Ali F, Moeen F, Shabbir S, Murad T	103
52. Usman M, Rahman S, Syed A	221	26. Navid S, Butt A, Aslam A, Asif S, Gul M, Ahmad S	106
53. Mahmood R, Safdar A, Khan M, Javaid F, Khan IM, Khattak MT	225	27. Gul M, Ahmad A, Butt A, Navid S, Asif S, Aslam A	110
54. Farman W, Kamran M, Khan AN, Summaya	229	28. Shabbir S, Shafay M, Murad T, Ali F, Yasmin N	115
55. Basit A, Zeb S, Umar M, Khan MM, Ullah I, Mehmood Z	233	29. Hussain A, Jadoon S, Javed S, Imtiaz H, Shaheen R, Hayat NQ	119
56. Khan Z, Haziq, Zia A, Ali G	237	30. Nasir A, Afridi A, Ahmed Z, Sadiq M, Khalily W, Fayyaz A	123
57. Zia A, Khan HD, Khan Z	241	31. Hafeez A, Iqbal J, Hassan A, Bhatti R, Kadir B, Sadik M	127
58. Kumar S, Khan MT, Kumari P, Latif S, Zehravi F	245		

Vol. 34, No. 9, September, 2023

Author (s)	Page No.
1. Jan MM.	1

32. Ali A, Latif H, Imtiaz J, Rehman A, Raziq S	131	10. Malik QA, Zeb S, Cheema UN, Haqdad M, Sajida S, Munir K	41
33. Shaikh S, Ismail K, Usman S, Ismail MO, Gul J, Shah R	136	11. Aqeel R, Mohsin B, Goraya A, Maqbool S, Andrabi WI, Khan A	45
34. Akram M, Ilyas J, Ashraf A, Naeem S, Ali N, Saeed M	141	12. Sangrasi MS, Shaikh FD, Ashraf A, Memon KN, Memon SR, Nizamani WM	49
35. Raza SZ, Ahmed S, Sanauallah Z, Memon ZA, Rais Z, Ahmed S	145	13. Aftab R, Butt AF, Leghari QA, Naeem N, Mukhtar S, Nasreen M	53
36. Gul H, Rehman S, Inayat Ullah, Khizar A	150	14. Shafique S, Khan MK, Jadoon SA, Khan SQ, Ahmad F, Hussain A	59
37. Azhar T, Aslam K, Malik KA, Nasreen M, Mukhtar S, Fatima ST	154	15. Zareef U, Ahmed T, Khero R, Arifa, Qamar A, Ali I	63
38. Nawaz A, Khan MI, Khan Q, Khan MA, Iqbal M, Jan Z	159	16. Ahmed N, Khan SM, Abidin Z, Malik AA, Abbasi T, Naeem M	68
39. Khan Q, Nawaz A, Khan MI, Khan MA, Jan Z, Iqbal M	164	17. Ahmad M, Imran M, Shafi N, Khan I, Rauf HA, Khan FA	72
40. Khaghan BS, Junaid F, Hussain SA, Noor AI, Iqbal T, Maham M	169	18. Khan FA, Rauf HA, Khan I, Shafi N, Imran M, Ahmad M	76
41. Masood MS, Hussain M, Rashid I	173	19. Amir A, Amir R, Amir A, Anwar Z	81
42. Bashir N, Razwan A, Tariq N, Saeed A, Toor S, Ahmad Z	178	20. Butt AF, Aftab R, Hafeez Z, Nasreen M, Mukhtar S, Faraz A	85
43. Arfin SMQ, Hussain S, Begum I, Raza SA, Ahmed AE	182	21. Mangrio SA, Naeem M, Jaffar S, Khadija, Zia MK, Shah MT	90
44. Khan MI, Nawaz A, Khan Q, Khan MA, Iqbal M, Jan Z	187	22. Kamran QA, Munir U, Cheema TN, Saleem K, Abaid T	95
45. Siddique HMAB, Khan S, Ghaffar MA, Shoaib M, Manzoor U, Irfan N	191	23. Masood MS, Hussain M, Rashid I	99
46. Rashid Z, Khawaja AF, Syed U, Andrabi WI, Munir SB, Sajid I	195	24. Latif U, Shaheen F, Daud M, Khan A, Haider A, Khan T	104
47. Zafar B, Mansoor A, Anis R, Anwar K, Aslam S	200	25. Mahmood R, Safdar A, Khan M, Sajjad A, Khattak MT, Khan IM	109
48. Iftikhar T, Parveen S, Khursheed I, Qamar S, Qazi N, Kamal N	204	26. Hidayat R, Raiz U, Ahmad R, Rahim M, Naseer A, Khan H	114
49. Haq N, Khan WN, Shah SN, Khan M	208	27. Liaqat S, Zada J, Khan SA	120
50. Nawaz B, Ilyas K, Mushtaq H, Jamal N, Baksh K, Maheshwary N	213	28. Haque E, Abbas A, Naeem- Ullah, Hanif MS, Khan H, Khan MH	125
51. Siddiqui SJ, Sutan R, Md Isa Z, Siddiqui SJ, Gemnani VK, Abr K	217	29. Alasmari D	129
		30. Alotaibi KZ	134

Vol. 34, No. 10, October, 2023

Author (s)	Page No.
1. Jan MM.	1
2. Abid MB, Afzal N, Zia W, Shiekh MA, Mahmood F, Ghous M	3
3. Rafe A, Munawar MT, Babar ZD, Talat K	9
4. Abbas M, Aftab R, Rahim F, Ansari T, Ali A, Ahmed M	13
5. Qaiser MA, Butt N, Anjum F, Fatima M, Fiaz A, Hafeez I	18
6. Rahim A, Nawaz T, Sajjad W, Rehman S, Jan DA, Bilal A	23
7. Malik M, Idrees RB	27
8. Jahan S, Khan M, Khan M, Zahir S, Syed R, Ashraf A	31
9. Hussain S, Sahrish, Din S, Habib SD	36

Vol. 34, No. 11, November, 2023

Author (s)	Page No.
1. Jan MM.	1
2. Arif H, Ali A, Ismail MO, Afridi FI, Abbas T, Mahmood S	3
3. Hayat K, Razzaq A, Saleem S, Ahmad E, Ahmed SH	8
4. Afzal A, Alam J, Rizwana, Saddique A, Ahmed N, Rehman A	13
5. Din N, Ahmed I, Riaz G, Bhutto AL, Matani K, Khan N	17
6. Gaffar I, Awaisi ZH	21
7. Qureshi T, Ali S, Fahad T, Yousaf M, Rashid A, Fatima N	25
8. Alotaibi Y, Khan N	28
9. Lakhani M, Sadiq M, Mukhtar S, Raza I, Azwar S, Hassan N	33
10. Ahmad F, Sadiq MS, Farooq A, Athar Y, Alam MA, Anwaar A	37

11. Shah MA, Ayaz S, Nizamuddin, Mehmood I, Shah M, Samiullah	42	2. Panezai MMK, Haq Z, Gul S, Rehman O, Shah S, Tahir S	3
12. Sultana N, Zafar K, Sikandar MZ, Khalid S, Safdar S, Sikandar S	46	3. Hameed J, Ali A, Ashraf MS, Khattak AH, Ali A, Naeem H	7
13. Sikandar MZ, Sultana N, Rehman A, Aashiq S, Safdar S, Zafar K	52	4. Ali A, Hameed J, Khattak AH, Ashraf MS, Naeem H, Ali A	12
14. Khan MAW, Sikandar MZ, Javed M, Khan ZA, Amjad H, Sami M	56	5. Shafi MK, Khan GQ, Ishfaq M, Hussain M, Hadi A	16
15. Hasan SJ, Riaz A, Bukhari SN, Munir T, Javed U, Talal T	60	6. Gaffar I, Awaisi ZH	20
16. Nawaz T, Amin M, Khan S, Ali SMN	66	7. Saad Ullah M, Shaheen KH, Riaz M	24
17. Daud M, Shaheen F, Nauman, Jawad	71	8. Ahmed S, Asnad	28
18. Khan S, Nawaz KH, Jadoon S, Saleem I, Jalil M, Ahmad I	75	9. Aleem A, Aziz K, Khalid M, Nazir S, Hassan MH	31
19. Rashid H, Bano R	80	10. Khan MAW, Khan AA, Sikandar MZ, Javed M, Wahab MMS, Mehmood A	35
20. Khan A, Akbar F, Zeb H, Afzal Z, Kakakhel M	85	11. Khan KK, Zahir S, Jahan S, Saleem S, Kakakhel S, Zeb S	39
21. Naheed, Haroon, Rehman N, Manan A, Asghar M, Zia Ullah	89	12. Javed N, Pervaiz S, Zahra F, Ali G, Muazzam A	44
22. Nawaz A, Manan F, Maindad J, Tahir R	93	13. Fiaz B, Riaz U, Akbar H, Noor A, Ahmad S, Khan H	49
23. Khan SA, Zada J, Liaqat S	97	14. Uzma, Ali S, Usman M, Rahman S	53
24. Khan Z, Sulaiman, Ali S, Rahaman A, Hussain SA, Khan A	102	15. Erfan M, Shahzadi A, Sagheer A, Khan I, Jabbar A, Akram S	58
25. Javaid F, Javed N, Ali G, Pervaiz S, Fatima, Tariq N	106	16. Hunain, Uzma, Rahman S	62
26. Arshad O, Anwar P, Ahtasham H, Chaudhary MAG, Riaz H, Ishaq W	110	17. Saleem S, Erfan M, Bacha MF, Khan I, Jabbar A, Akram S	66
27. Agwan MAS	115	18. Rehmat S, Khan S, Khan I	70
28. Iqbal S, Syed NK	121	19. Orakzai AN, Wazir BG, Muhammad N	74
		20. Ahmad A, Asif M, Khattak JJ, Zahoor I, Khan MS, Anwar K	79
		21. Sahitia S, Idris IB, Safian N, Shamsuddin K, Hod R	84
		22. Mohammed NKS	90
		23. Javed MQ, Srivastava S	95
		24. Mustafa G	99
Vol. 34, No. 12, December, 2023			
Author (s)	Page No.		
1. Jan MM.	1		

Subject Index January to December 2023**Azhar Masud Bhatti**

Editor in Chief

Vol. 34, No. 1, January, 2023

Subject	Page No.
<u>CARBAPENEMASE-PRODUCING GRAM-NEGATIVE FROM WOUND INFECTIONS</u>	
• Phenotypic and Molecular Characterization of Carbapenemase-Producing Gram-Negative Isolates Detected from Wound Infections (Alruways MW)	2
<u>FACTORS OF SEVERE OBSTETRIC MORBIDITY</u>	
• Frequency of Risk Factors of Severe Obstetric Morbidity (Nawaz E, et al)	8
<u>HISTOPATHOLOGICAL GRADING OF SQUAMOUS CELL CARCINOMA</u>	
• Histopathological Grading of Squamous Cell Carcinoma at Time of Presentation in Tertiary Care Hospital (Afzal A, et al)	12
<u>FEMUR LENGTH, BI-PARIETAL DIAMETER AND ABDOMINAL CIRCUMFERENCE IN 2ND AND 3RD TRIMESTER</u>	
• Assessment of Fetal Biparietal Diameter, Femur Length and Abdominal Circumference in Local Population on Ultrasonography in Second and Third Trimester of Pregnancy (Mushtaq S, et al)	16
<u>HESITANCE FROM GETTING COVID-19 VACCINATION</u>	
• Evaluation of Hesitance from Getting COVID-19 Vaccination Among People of Mirpur AJK (ShoaiB M, et al)	22
<u>GROWTH PATTERNS OF SCHOOL CHILDREN</u>	
• Prevalence and Factors Associated with Growth Patterns of School Children in Karachi (Akhtar N, et al)	25
<u>TOOTH MOBILITY IN PERIODONTITIS PATIENTS</u>	
• Prevalence of Tooth Mobility in Periodontitis Patients (Abrar SK, et al)	29
<u>EFFICACY OF RACECADOTRIL IN ACUTE WATERY DIARRHEA</u>	
• Efficacy of Racecadotril in Acute Watery Diarrhea as Compared to Placebo (Munir A et al)	33
<u>INTERNAL DISASTER PREPAREDNESS OF TERTIARY CARE HOSPITALS</u>	
• Level of Hospital Preparedness for Internal Disasters in Tertiary Care Hospitals in Pakistan (Rawalpindi City) (Omer Z, et al)	37

Subject	Page No.
<u>COMPARE THE EFFICACY OF HOLEP VS TURP</u>	
• Comparison of Blood Loss in Holmium Laser Enucleation of the Prostate (HoLEP) Versus Transurethral Resection of Prostate (TURP) (Zahid MM, et al)	43
<u>EVALUATION OF DRUG RELEASE FROM AMOXICILLIN IN PERIODONTAL MEMBRANES</u>	
• Synthesis & Evaluation of Drug Release from Amoxicillin Loaded Chitosan Nanoparticles Incorporated in the Periodontal Membranes (Ashraf N, et al)	47
<u>HEPATITIS C WITH DIRECTLY ACTING ORAL ANTIVIRALS</u>	
• End Treatment Response of Chronic Hepatitis C with Directly Acting Oral Antivirals at 12 Weeks (Zia D, et al)	52
<u>ISCHEMIC HEART DISEASE AND DYSLIPIDEMIAS IN HYPOTHYROIDISM</u>	
• To Determine the Prevalence of Ischemic Heart Disease and Dyslipidemias in Patients of Subclinical Hypothyroidism (Rehman SS, et al)	56
<u>SELF MEDICATION OF ANTIBIOTICS IN CHILDREN</u>	
• The Factors Leading to Parental Self Medication of Antibiotics in Children (Janjua HB, et al)	60
<u>DIALYSIS WITH SUPRACLAVICULAR TUNNEL LOCATION OF DIALYSIS CATHETER</u>	
• Assessing the Comfort Level of Dialysis Patients with Supraclavicular Tunnel Location of Tunneled Dialysis Catheter: A Tertiary Care Center Experience In Karachi, Pakistan (Siddiqui DK, et al)	64
<u>IMPLICATIONS OF SUB-CATEGORIZATION OF CHRONIC HEADACHE</u>	
• Importance and Implications of Sub-Categorization of Chronic Headache and the Incidence in General Medical OPD of a Teaching Hospital (Rana MM, et al)	69
<u>BRACKA'S AND SNODGRASS SURGICAL TECHNIQUES FOR HYPOSPADIAS</u>	
• Bracka's and Snodgrass Surgical Techniques for Hypospadias Correction: A Comparative Study (Rasheed S, et al)	73

Subject	Page No.	Subject	Page No.
<u>EFFECT OF ASSERTIVENESS-BASED PROGRAMME ON PSYCHOLOGICAL WELLBEING AMONG NOVICE NURSES</u>		<u>SAFETY AND EFFICACY OF FERRIC CARBOXYMALTOSE WITH IRON</u>	
• The Effect of Assertiveness-Based Programme On Psychological Wellbeing and Work Engagement Among Novice Nurses in Psychiatric Department (Khanam S, et al) 78	78	• Comparison of Safety and Efficacy of Ferric Carboxymaltose with Iron Sucrose for the Treatment of Iron Deficiency Anemia in Pregnancy (Sattar S, et al) 15	15
<u>DESQUAMATIVE GINGIVITIS AMONG DENTAL PATIENTS</u>		<u>ANTIFUNGAL SUSCEPTIBILITY OF THE CANDIDA IN NAIL INFECTIONS</u>	
• Frequency of Desquamative Gingivitis Among Dental Patients: A Cross Sectional Multi-Center Study (Mirza D, et al) 83	83	• Antifungal Susceptibility Pattern of the Candida Species in Nail Infections (Kashif S, et al) 20	20
<u>COVID-19 ASSOCIATED EYE PROBLEMS</u>		<u>EARLY CATHETER REMOVAL AFTER TRANSURETHRAL RESECTION OF PROSTATE</u>	
• Covid-19 Associated Eye Problems in the Pakistani Population (Ashraf NN, et al) 87	87	• Impact of Early Catheter Removal After Transurethral Resection of Prostate: A Comparative Study of Post-Op Complications in Patients with Early VS Delayed Catheter Removal (Shahzad M, et al) 25	25
<u>KNOWLEDGE AND PERCEPTION OF PRACTITIONERS FOR DENTAL IMPLANT TREATMENT</u>		<u>PREFERENCES AND PRACTICES OF FIXED PROSTHODONTICS TREATMENT MODALITIES</u>	
• Knowledge and Perception of Dental Practitioners towards Dental Implant Treatment in Routine Dental Practice (Iqbal M, et al) 91	91	• Determination of Preferences and Practices of Fixed Prosthodontics Treatment Modalities by Dental Practitioners (Khan MS, et al) 30	30
<u>TRACHEOSTOMY CARE INTERVENTION ON SELF-CARE KNOWLEDGE</u>		<u>EFFECT OF OXYGEN THERAPY VIA FACE MASK</u>	
• Effect of Tracheostomy Care Intervention on Self-Care Knowledge and Level of Anxiety among Adult Patients with Permanent Tracheostomy (Rasool G, et al) 96	96	• To Compare the Effect of Oxygen Therapy Via Face Mask with Bi-Level Positive Airway Pressure Ventilation (BIPAP) in Managing Patients with Acute Cardiogenic Pulmonary Edema in Al-Nafees Medical College and Hospital, Islamabad (Rehman SS, et al) 35	35

Vol. 34, No. 2, February, 2023

Subject	Page No.	Subject	Page No.
<u>EVALUATE THE STRUCTURAL & ANTIBACTERIAL PROPERTIES FOR PERIODONTAL TISSUE REGENERATION</u>		<u>ATTITUDE AND PERCEPTION OF HEALTH PROFESSIONAL</u>	
• Fabrication & Analysis of the Nanohydroxy Apatite / Chitosan Membranes for Rejuvenation of Periodontal Tissues (Ashraf N, et al) 2	2	• Attitude and Perception of Health Professional Students Toward Inter-Professional Education in Riyadh: Cross-Sectional Study (Darraj AQ, et al) 40	40
<u>POSTGRADUATE RESIDENTS TOWARDS PROMOTION BY PHARMACEUTICAL COMPANIES</u>		<u>EVALUATION OF ANXIETY AND DEPRESSION IN SQUAMOUS CELL CARCINOMA OF ORAL CAVITY</u>	
• Perceptions of Postgraduate Residents towards Promotion by Pharmaceutical Companies at a Teaching Hospital (Memon ZR, et al) 7	7	• Evaluation of Anxiety and Depression Level in Newly Diagnosed Cases of Squamous Cell Carcinoma of Oral Cavity (Azhar MS, et al) 46	46
<u>EFFECTIVE PHOTOTHERAPY AND IV IMMUNOGLOBULIN IN KERNICTERUS</u>		<u>AWARENESS AMONG MOTHERS REGARDING HAND HYGIENE</u>	
• Frequency of Kernicterus in Rh Incompatibility after Receiving Effective Phototherapy and Intravenous Immunoglobulin Therapy (Ali S, et al) 11	11	• Assessment of Awareness Among Mothers Regarding Hand Hygiene and Safe Water in Urban Slum Areas of Southern Punjab (Saddique A, et al) 50	50
		<u>UNMET NEEDS OF FAMILY PLANNING</u>	
		• Unmet Needs of Family Planning and Associated Factors in Married Women of Reproductive Age (Masood R, et al) 54	54

Subject	Page No.
<u>ACCURACY OF ENDOSCOPIC FINDINGS OF CELIAC DISEASE</u>	
• Diagnostic Accuracy of Endoscopic Findings of Patients with Suspected Celiac Disease Taking Histological Features as Gold Standard at Tertiary Care Hospital (Islam N, et al)	58
<u>BONE MARROW INFILTRATION IN HOGKIN'S LYMPHOMA</u>	
• Pattern of Bone Marrow Infiltration in Hogkin's Lymphoma (Shah MH, et al)	62
<u>POST-OPERATIVE COMPLICATIONS OF MANDIBLE</u>	
• Comparing the Post-operative Complications of Mandible Angle Fractures Treated by Extra Oral versus Intra Oral Reduction Approach (Kumari S, et al)	66
<u>COMPARISON OF METAISEAU TECHNIQUE WITH A PERCUTANEOUS K-WIRE FIXATION FOR RADIAL NECK FRACTURES</u>	
• Comparison of Metaiseau Technique with a Percutaneous K-Wire Fixation for Displaced Radial Neck Fractures in Pediatric Age Group (Hussain I, et al)	71
<u>HYPERTENSIVE DISORDERS OF PREGNANCY</u>	
• Prevalence of Risk Factors of Hypertensive Disorders of Pregnancy in a Tertiary Care Hospital of Rawalpindi (Aslam A, et al)	76
<u>ESTROGEN AND LIPOPROTEINS IN PRE AND POSTMENOPAUSAL HYPERTENSIVE</u>	
• A Comparative Study Between Estrogen and Lipoproteins in Pre and Postmenopausal Hypertensive Women (Aghani N, et al)	80
<u>LOOP STOMA OVER DIVIDED STOMA IN CASE OF HIRSCHSPRUNG</u>	
• Success of Loop Stoma Over Divided Stoma in Case of Hirschsprung Disease (Habib KJ, et al)	84
<u>POST-OPERATIVE WOUND INFECTION AND ITS MANAGEMENT IN PEDIATRIC</u>	
• Post-Operative Wound Infection and its Management in Pediatric Surgical patients: A Study of 560 Infants and Children (Mansoori MS, et al)	89
<u>MORTALITY PATTERN OF PATIENTS IN NEONATAL ICU</u>	
• Mortality Pattern of Patients in Neonatal Intensive Care Unit at Indus Hospital, Karachi (Kanhar IA, et al)	94
<u>NURSES KNOWLEDGE AND PRACTICES REGARDING ASEPTIC TECHNIQUES</u>	
• Effect of Intervention Based Centers for Disease Control and Prevention Guidelines on Nurses Knowledge and Practices Regarding Aseptic Techniques in Operating Room (Tabassum Z, et al)	98

Subject	Page No.
<u>PSYCHOMETRIC APPRAISAL OF ARABIC INTERPROFESSIONAL ATTITUDES</u>	
• The Psychometric Appraisal of Arabic Interprofessional Attitudes Scale (Darraj AQ)	2
<u>BLOOD DONOR DEFERRAL CAUSES</u>	
• A Study of Blood Donor Deferral Causes: Pre versus Post Donation and Transfusion Transmissible Infections (Fatima M, et al)	8
<u>GROSS HEMATURIA AND/OR DROP IN HEMOGLOBIN IN POST-RENAL BIOPSIES</u>	
• Evaluation of Gross Hematuria and/or Drop in Hemoglobin in Post-Renal Biopsies within and after 8 Hours (Rahim S, et al)	12
<u>ASCITIC FLUID PROTEINS WITH SPONTANEOUS BACTERIAL PERITONITIS</u>	
• Relation of Low Ascitic Fluid Proteins with Spontaneous Bacterial Peritonitis (SBP) in Patients with Portosystemic Encephalopathy – A Comparative Study (Himayat Ullah, et al)	17
<u>SHORT CONTACT 1.5% DITHRANOL IN ALOPECIA AREATA</u>	
• Efficacy of Short Contact 1.5% Dithranol Therapy in Mild to Moderate Alopecia Areata in Patients Reporting to Tertiary Care Hospital Karachi (Khosro BK, et al)	22
<u>H3PO4 & HCL IN MICROABRASION TECHNIQUE FOR DENTAL FLUOROSIS</u>	
• Comparison Between Phosphoric Acid and Hydrochloric Acid in Microabrasion Technique for the Treatment of Dental Fluorosis (Madiha, et al)	27
<u>CVM STAGES AND MAXILLARY CANINE CALCIFICATION STAGES</u>	
• Comparison of CVM Stages and Maxillary Canine Calcification Stages for Determination of Skeletal Maturity (Enver N, et al)	32
<u>ESTIMATING STATURE USING HANDPRINT DIMENSIONS</u>	
• Estimating Stature Using Handprint Dimensions: A Medico-Legal Investigative Approach (Waheed A, et al)	35
<u>ANTI-DIABETIC EFFECTS OF OMEGA-3-FATTY ACIDS</u>	
• Miraculous Therapy for Diabetes: Use of Omega-3-Fatty Acids (Qazi AF, et al)	40
<u>SUPINE PCNL WITH RENAL CALCULI</u>	
• Outcome of Supine Percutaneous Nephrolithotomy as a Secondary Procedure for Renal Calculi (Jamil MN, et al)	45
<u>HYPOPARATHYROIDISM WITH PATIENTS OF BETA THALASSEMIA MAJOR</u>	
• Relationship of Hypoparathyroidism with Patients of Beta Thalassemia Major in a Tertiary Care Hospital (Khattak MB, et al)	49

Subject	Page No.	Subject	Page No.
<u>HYPERURICEMIA WITH LOW DOSE ASPIRIN USE IN MIDDLE AGED AND ELDERLY</u>		<u>FLIPPED CLASSROOM APPROACH ON HEALTH ASSESSMENT KNOWLEDGE AND SKILLS</u>	
<ul style="list-style-type: none"> Association of Hyperuricemia with Low Dose Aspirin Use in Middle Aged and Elderly Population (Khan A, et al) 53 		<ul style="list-style-type: none"> Effects of Flipped Classroom Approach on Health Assessment Knowledge and Skills Among Undergraduate Nursing Students (Rafique A, et al) 102 	
<u>PERIMENOPAUSAL FEMALES AS HYPERTENSION DURING VASOMOTOR SYMPTOMS</u>		Vol. 34, No. 4, April, 2023	
<ul style="list-style-type: none"> Inadvertent Labelling of Perimenopausal Females as Hypertension During Vasomotor Symptoms, Experience at a Tertiary Care Hospital (Raza MM, et al) 58 		Subject	Page No.
<u>EFFECT OF METFORMIN ON FASTING INSULIN LEVELS IN NON-DIABETIC OBESE</u>		<u>SATISFACTION AND CHALLENGES OF ONLINE LEARNING</u>	
<ul style="list-style-type: none"> Effect of Metformin on Fasting Insulin Levels in Non-Diabetic Obese Patients Over 8 Weeks of Time (Rehman SS, et al) 63 		<ul style="list-style-type: none"> Satisfaction and Challenges of Online Learning Among Healthcare Students: Towards Digital Education in the Healthcare Profession in Saudi Arabia (Thangam MMN) 2 	
<u>ANTIMICROBIAL EFFICACY OF MTAD AND EDTA</u>		<u>HYPER-PROLACTINEMIA AND HYPOTHYROIDISM IN INFERTILE WOMEN</u>	
<ul style="list-style-type: none"> Comparison of the Antimicrobial Efficacy of Mtad and Edta Against Enterococcus Faecalis in Vitro (Aftab U, et al) 67 		<ul style="list-style-type: none"> Frequency of Hyperprolactinemia and Hypothyroidism in Infertile Women (Choudhary SA, et al) 7 	
<u>ASSESSMENT OF VITAMIN D DURING COVID-19</u>		<u>PLASMA D - DIMER LEVELS WITH CTPA IN PATIENTS PULMONARY EMBOLISM</u>	
<ul style="list-style-type: none"> Assessment of Vitamin D among University Students During Covid-19 Period (Mangi AH, et al) 71 		<ul style="list-style-type: none"> Comparison of Plasma D - Dimer levels with Computed Tomography Pulmonary Angiography (CTPA) Findings in Patients Suspected of Pulmonary Embolism Clinically (Akhtar MS, et al) 12 	
<u>TREATMENT RELATED TOXICITY IN GYNECOLOGICAL MALIGNANCIES</u>		<u>TYPE 1 DIABETES MELLITUS WITH BODY MASS INDEX</u>	
<ul style="list-style-type: none"> Treatment Related Toxicity in Gynaecological Malignancies after Conventional Radiotherapy (Gul U, et al) 75 		<ul style="list-style-type: none"> Association of Type 1 Diabetes Mellitus with Body Mass Index (Ghias Z, et al) 15 	
<u>EFFICACY OF ONDANSETRON AND METOCLOPRAMIDE IN CHILDREN WITH GASTROENTERITIS WITH VOMITING</u>		<u>MRI IN EVALUATION OF PERIANAL FISTULA AS GOLD STANDARD</u>	
<ul style="list-style-type: none"> Efficacy of Single Dose Intravenous Ondansetron VS Metoclopramide in Children with Acute Gastroenteritis with Vomiting (Khan A, et al) 80 		<ul style="list-style-type: none"> Diagnostic Accuracy of Magnetic Resonance Imaging in Evaluation of Perianal Fistula Keeping Surgical Finding as Gold Standard (Ahmed A, et al) 19 	
<u>VIRTUAL TEACHING DURING THE COVID-19</u>		<u>SERUM CORTISOL VALUES IN COSYNTROPIN STIMULATION TEST</u>	
<ul style="list-style-type: none"> Virtual Teaching During the COVID-19 Pandemic and the Psychosocial Well-Being of Medical Students at College of Medicine, Jeddah (Ahmed SN, et al) 85 		<ul style="list-style-type: none"> Clinical Utility of 30- and 60-min Serum Cortisol Values in Cosyntropin Stimulation Test for Diagnosis of Adrenal Insufficiency (Dildar S, et al) 22 	
<u>SMARTPHONE ADDICTION AMONG UNIVERSITY STUDENTS</u>		<u>FOREIGN BODIES IN THE URINARY BLADDER AND MANAGEMENT</u>	
<ul style="list-style-type: none"> Smartphone Addiction among University Students: A Cross- Sectional Study (Gemnani VK, et al) 92 		<ul style="list-style-type: none"> Foreign Bodies in the Urinary Bladder! A Single-Center Experience in Endoscopic Management (Khalid M, et al) 25 	
<u>GUIDELINES ON NURSE'S KNOWLEDGE AND PRACTICES REGARDING HEMODIALYSIS</u>			
<ul style="list-style-type: none"> The Effect of Standardized Guidelines on Nurse's Knowledge and Practices Regarding Hemodialysis at Tertiary Care Hospital Lahore (Saleem I, et al) 98 			

Subject	Page No.	Subject	Page No.
<u>0.5% TIMOLOL MALEATE AND 0.2% BRIMONIDINE TARTRATE IN CONTROL OF RAISED INTRA OCULAR PRESSURE</u>		<u>HYPERURICEMIA IN PATIENTS WITH HEART DISEASES</u>	
• Efficacy of 0.5% Timolol Maleate and 0.2% Brimonidine Tartrate in Control of Raised Intra Ocular Pressure after Neodymium Yttrium Aluminium Garnet Laser Capsulotomy (Haq QM, et al) 30		• Frequency of Hyperuricemia in Patients with Heart Diseases (Anwar U, et al) 67	
<u>SPECULAR MICROSCOPY IN EMMETROPIC AND ASTIGMATIC PATIENTS</u>		<u>OBESITY IN TYPE II DIABETES MELLITUS PATIENTS</u>	
• Outcome of Specular Microscopy in Emmetropic and Astigmatic Patients (Ashraf NNA, et al) 33		• Frequency of Obesity in Type II Diabetes Mellitus Patients (Zeb F, et al) 70	
<u>KNOWLEDGE AND PRACTICE OF OXYGEN THERAPY</u>		<u>HELICOBACTER PYLORI GASTRITIS IN DYSPEPTIC PATIENTS</u>	
• Effect of Educational Program on Doctors' Knowledge and Practice of Oxygen Therapy (Mustafa G) 36		• Prevalence of Helicobacter Pylori Gastritis in Dyspeptic Patients (Pervez S, et al) 73	
<u>EVALUATION OF MULTI VESSEL DISEASE</u>		<u>ASSESSMENT OF COVID-19 AWARENESS DURING LOCKDOWN</u>	
• Evaluation of Multi Vessel Disease and its Impact on Reperfusion Success in Patients Undergoing Primary Percutaneous Coronary Intervention for Acute Myocardial Infarction (Ramzan M, et al) 41		• Assessment of COVID-19 Awareness among University Students During Lockdown Period (Mangi AH, et al) 75	
<u>HISTOPATHOLOGICAL DIAGNOSIS OF INCIDENTAL GALL BLADDER CARCINOMA</u>		<u>GENEXPERT IN ACTIVE CASE DETECTION AMONG DRUG RESISTANT TB</u>	
• Histopathological Diagnosis of Incidental Gall Bladder Carcinoma in Patients Undergoing Routine Laparoscopic Cholecystectomy (Sami Ullah M, et al) 46		• Efficacy of GeneXpert in Active Case Detection Among Contacts of Drug Resistant TB Patients (Ahmad S, et al) 80	
<u>MALOCCLUSION WITH CENTRIC RELATION AND MAXIMUM INTERCUSPATION</u>		<u>POSITIVE FLUID BALANCE AS A RISK FACTOR FOR ACUTE KIDNEY INJURY</u>	
• Malocclusion in Relation to Centric Relation and Maximum Intercuspation Coincidence and Gender (Afshan Z, et al) 50		• Positive Fluid Balance as a Risk Factor for Acute Kidney Injury: Experience of a Tertiary Care Hospital (Shamim S, et al) 85	
<u>MEASLES COMPLICATIONS IN CHILDREN</u>		<u>MEASURES OF HEPATITIS B AND THE ATTITUDE OF NURSING & DENTAL STUDENTS</u>	
• Frequencies of Different Measles Complications in Children Presenting at a Tertiary Care Hospital, Larkana (Bhojwani SL, et al) 54		• Preventative Measures of Hepatitis B and the Attitude of Nursing & Dental Students: A Comparative Study (Wassan SM, et al) 90	
<u>SKELETAL MATURITY ASSESSMENT USING MAXILLARY CANINE</u>		Vol. 34, No. 5, May, 2023	
• Skeletal Maturity Assessment Using Maxillary Canine Calcification Stages (Enver N, et al) 58		Subject	Page No.
<u>PET/CT AND NLR BEFORE AND AFTER TREATMENT WITH DIFFERENT MALIGNANCY</u>		<u>PERIMENOPAUSAL SYMPTOMS SIGNALS THE NEED TO EVALUATE</u>	
• Correlation between PET/CT Maximum Standard Uptake Value and NLR in Malignancy Patients Pre- and Post-Treatment (Sunil RK, et al) 62		• The Higher Severity Index of Vasomotor Perimenopausal Symptoms Signals the Need to Evaluate Coexisting Metabolic Syndrome for CVD Risk Factors Stratification Experience (Rana MM, et al) 2	
		<u>MAXILLOFACIAL INJURIES OF MOTOR BIKE ACCIDENTS</u>	
		• Pattern of Maxillofacial Injuries in Patients of Motor Bike Accidents with Helmet and without Helmet (Zahoor S, et al) 6	

Subject	Page No.	Subject	Page No.
<u>CARDIOPULMONARY BYPASS AND CONTINUOUS LTV</u>		<u>ASSESSMENT OF THE PERSISTENCE OF ANTI SARS-COV2 IGG ANTIBODY IN INDUSTRIAL WORKERS</u>	
• Cardiopulmonary Bypass and Continuous Low Tidal Volume (Asad H, et al)	7	• Assessment of the Persistence of Anti SARS-Cov2 Igg Antibody Over a Period of Six Months Among Industrial Workers (Dildar S, et al)	48
<u>MATERNAL AND FETAL COMPLICATION IN OBESE PREGNANT</u>		<u>OUTCOME OF DISTAL HUMERUS FRACTURES TREATED WITH ORIF</u>	
• Frequency of Maternal and Fetal Complication in Obese Pregnant Women (Rafiq S, et al)	11	• Functional Outcome of Distal Humerus Fractures Treated with Open Reduction and Internal Fixation Through Bi-Columnar Plating (Essa MA, et al)	51
<u>ETIOLOGICAL PROFILE OF NON-COMPRESSIVE MYELOPATHIES</u>		<u>DYNAMIC CONDYLAR SCREW AND LOCKING PLATE FIXATION IN FEMORAL FRACTURE</u>	
• Clinical and Etiological Profile of Non-compressive Myelopathies: Experience from a Tertiary Care Hospital in Pakistan (Jawaid W, et al)	15	• Comparative Study Between Dynamic Condylar Screw and Locking Plate Fixation in Fixation of Distal Femoral Fracture in Adults (Farooq MZ, et al)	56
<u>HPLC: COST-EFFECTIVE AND ACCURATE METHOD FOR PROPOFOL PLASMA</u>		<u>RELIEF IN ARTHOPLASTY</u>	
• High-Performance Liquid Chromatography (HPLC): A Cost-Effective and Accurate Method for Propofol Plasma Level Monitoring (Naem U, et al)	20	• Effect of Duloxetine on Pain Relief in Patients Undergoing Arthroplasty (Inamullah, et al)	63
<u>AXILLARY SURGERY IN BREAST CANCER</u>		<u>BRAIN ABSCESES FROM NECROTIZING BRAIN TUMORS USING SURGICAL FINDINGS</u>	
• Conservative Axillary Surgery in Management of Breast Cancer: An Emerging Surgical Approach (Hussain S, et al)	24	• Diffusion-Weighted Magnetic Resonance Imaging's Diagnostic Precision in Separating Brain Abscesses from Necrotizing Brain Tumors Using Surgical Findings as the Gold Standard (Haq A, et al)	68
<u>ANTIVIRALS (DAA) ON THROMBOCYTOPENIA IN CHRONIC HEPATITIS C</u>		<u>ULTRASOUND'S PREDICTIVE POWER FOR EARLY PREGNANCY FAILURE</u>	
• Effect of Direct Acting Antivirals (DAA) on Thrombocytopenia in Chronic Hepatitis C Patients Without Cirrhosis (Khan SA, et al)	28	• First Trimester Ultrasound's Predictive Power for Early Pregnancy Failure in Pakistan: A Single Center Study (Afridi F, et al)	72
<u>IRON STATUS IN FEMALE METABOLIC SYNDROME</u>		<u>FINDINGS WITH HISTOPATHOLOGICAL DIAGNOSIS OF BREAST LESIONS</u>	
• Assessment of Iron Status in Female Metabolic Syndrome Patients (Hammad AR, et al)	33	• Correlation of Mammographic and Ultrasonographic Findings with Histopathological Diagnosis of Breast Lesions (Nawab K, et al)	76
<u>CORTICOSTEROID NEBULISATION IN CHRONIC COUGH</u>		<u>SAFETY AND EFFICACY OF BIPOLAR ENERGY FOR TRANSURETHRAL RESECTION OF BLADDER TUMOR</u>	
• Rational of Corticosteroid Nebulisation in Chronic Cough (Bashir N, et al)	36	• A Prospective Study on Safety and Efficacy of Bipolar Energy for Transurethral Resection of Bladder Tumor (Ahmad N, et al)	79
<u>ORAL HEALTH KNOWLEDGE AND PRACTICE OF PREGNANT</u>		<u>GASTROESOPHAGEAL REFLUX DISEASE</u>	
• Oral Health Knowledge and Practice of Pregnant Women at Chiniot General Hospital Korangi Karachi (Anum M, et al)	40	• Gastroesophageal Reflux Disease: Population-Based Study in Tertiary Care Hospital, Karachi (Zakir H, et al)	84
<u>POST-INSERTION COMPLAINTS OF REMOVABLE PARTIAL DENTURES</u>			
• Post-Insertion Complaints of Removable Partial Dentures and Their Association with Age and Gender (Mehtab B, et al)	44		

Subject	Page No.
<u>OCCUPATIONAL HAZARD AMONG DENTIST</u>	
• Knowledge and Awareness About the Occupational Hazard Among Dentist in Tertiary Care Hospital in Karachi (Raza SMM, et al)	88
<u>ALBINO RATS AFTER CONSUMPTION OF A HIGH CARBOHYDRATE DIET</u>	
• Assessment of Gross Parameters in Maternal Albino Rats after Consumption of a High Carbohydrate Diet: A Comparative Study (Korai SM, et al)	93
<u>CARPAL TUNNEL SYNDROME AMONG OBESE</u>	
• Evidence of Carpal Tunnel Syndrome Among Obese Patients (Shafee SM, et al)	97
<u>BREAST CANCER WITH RADIOTHERAPY AND HORMONE THERAPY</u>	
• Relapse Free Survival Time of Breast Cancer Patients Treated With Adjuvant Radiotherapy and Hormone Therapy (Raza U, et al)	101
<u>LUNG FUNCTION OF HEALTHY, NON-SMOKING YOUNG PEOPLE</u>	
• Lung Function Parameters of Healthy, Non-Smoking Young People and Employees at Shaqra University, Shaqra (Mustafa G, et al)	105
<u>ASSOCIATION OF ACNE VULGARIS AND BMI</u>	
• Frequency and Association of Acne Vulgaris and BMI: A Case Control Study (Mirza U, et al)	111
<u>L-ARGININE ON STZ-INDUCED ADRENAL CORTICAL DAMAGE</u>	
• Protective Effect of L-Arginine on Streptozotocin (STZ)-Induced Adrenal Cortical Damage (Mahar Y, et al)	115
<u>HISTOPATHOLOGICAL SPECTRUM OF UPPER GASTROINTESTINAL BIOPSIES</u>	
• Histopathological Spectrum of Upper Gastrointestinal Endoscopic Biopsies in a Tertiary Care Hospital (Siraj F, et al)	119
<u>HEPA FILTERS ON THE INCIDENCE OF INVASIVE FUNGAL INFECTION IN COVID 19</u>	
• The Impact of Portable High Efficiency Hepa Filters on the Incidence of Invasive Fungal Infection in COVID 19 Inpatients in a Tertiary Care Hospital: A Retrospective Analysis From Pakistan (Ahmed A, et al)	124
<u>MATERNAL SERUM VITAMIN D LEVEL WITH PREECLAMPSIA</u>	
• Association of Maternal Serum Vitamin D Level with Preeclampsia (Shahid A, et al)	129

Subject	Page No.
<u>ROLE OF EPIGENETICS THERAPIES FOR ALZHEIMER'S DISEASE</u>	
• Investigating the Role of Epigenetics in the Development of Neurological Disorders Developing Novel Therapies for Alzheimer's Disease (Uqaili AA, et al)	135
<u>EVALUATION OF PSYCHO-SOCIAL ENVIRONMENT OF DIFFERENT SCHOOLS</u>	
• Evaluation of Psycho-Social Environment of Army Public Schools and Private Schools in Rawalpindi – A Comparative Study (Kahloon OI, et al)	139

Vol. 34, No. 6, June, 2023

Subject	Page No.
<u>HYPERKALAEMIA IN NON-DIALYSIS OF CKD</u>	
• Frequency of Hyperkalaemia in Non-Dialysis Dependent Chronic Kidney Disease (CKD) Patients (Qadir A, et al)	2
<u>DEFECTS IN SURFACE DETAIL OF 2 PHASE ONE STEP VS 2 PHASE TWO STEP</u>	
• Evaluation of Defects in Surface Detail of 2 Phase One Step Versus 2 Phase Two Step Impression Techniques: An In Vitro Study (Yaqub K, et al)	6
<u>PUMPKIN SEED EXTRACT ON BISPHENOL A INDUCED TESTICULAR TOXICITY</u>	
• Protective Effect of Pumpkin Seed Extract on Bisphenol A Induced Testicular Toxicity in Adult Albino Rats (Batool U, et al)	10
<u>HEMATOLOGICAL AND BIOCHEMICAL PARAMETERS OF DENGUE SEVERITY</u>	
• Predicting the Unpredictable: Hematological and Biochemical Parameters as Predictive Markers of Dengue Severity in Patients Without Warning Sign (Amir E, et al)	14
<u>BMI ON PULMONARY FUNCTION TEST VALUES AMONG EMPLOYEES & YOUNG</u>	
• Influence of Body Mass Index (BMI) on Pulmonary Function Test Values Among Employees & Young People of Shaqra University (Mustafa G, et al)	19
<u>PLACENTAL GRADING IN NORMAL AND PREGNANCY INDUCED HYPERTENSIVE</u>	
• Assessment of Placental Grading in Normal and Pregnancy Induced Hypertensive Mothers on Ultrasonography and Fetal Outcomes (Mushtq S, et al)	24

Subject	Page No.	Subject	Page No.
<u>CONCEPT OF PLATINUM SENSITIVITY IN CERVICAL CANCER WOMEN</u>		<u>MIDDLE CEREBRAL TO UMBILICAL ARTERY DOPPLER RATIO IN NEONATAL OUTCOMES</u>	
<ul style="list-style-type: none"> • Pattern of Recurrence in Locally Advanced Carcinoma Cervix Based on Platinum Sensitivity (Khan S, et al) 28 		<ul style="list-style-type: none"> • The Predictive Significance of the Middle Cerebral to Umbilical Artery Doppler Ratio in Determining Neonatal Outcomes in Patients with Preeclampsia and Gestational Hypertension (Asghar MR, et al) 73 	
<u>ECTROPION IN SUBCILIARY VERSUS SUBTARSAL INCISIONS</u>		<u>MYOCARDIAL BRIDGING AND CLINICAL INDICATIONS FOR CORONARY ANGIOGRAPHY</u>	
<ul style="list-style-type: none"> • Incidence of Ectropion in Subciliary Versus Subtarsal Incisions (Manzar S, et al) 32 		<ul style="list-style-type: none"> • Myocardial Bridging, Frequency, Severity, Classification and Clinical Indications for Coronary Angiography (Rehman M, et al) 78 	
<u>LOCAL STEROID INJECTION AND CARPAL TUNNEL RELEASE</u>		<u>HISTOPATHOLOGICAL VARIETIES OF PROSTATIC DISEASES</u>	
<ul style="list-style-type: none"> • Comparison of Local Steroid Injection and Carpal Tunnel Release in Carpal Tunnel Syndrome (Khan KR, et al) 36 		<ul style="list-style-type: none"> • Frequencies of Several Histopathological Varieties of Prostatic Diseases in a Tertiary Care Hospital at Larkana (Saleem MW, et al) 82 	
<u>PROPHYLACTIC AMIODARONE VS DEXMEDETOMIDINE ON REDUCING OF POSTOPERATIVE TACHYCARDIA</u>		<u>BENIGN AND MALIGNANT BREAST GROWTH AMONG WOMEN</u>	
<ul style="list-style-type: none"> • Effect of Prophylactic Amiodarone Versus Dexmedetomidine on Reducing the Frequency of Postoperative Junctional Ectopic Tachycardia after Pediatric Open Heart Surgery (Arshad T, et al) 40 		<ul style="list-style-type: none"> • Frequency of Benign and Malignant Breast Growth Among Women of Different Age Groups in Larkana Population (Fauzia T, et al) 86 	
<u>TROPONIN I AND COMPLICATION AFTER CABG</u>		<u>ELASTOGRAPHY IN DIFFERENTIATION OF MALIGNANT AND BENIGN BREAST LESIONS</u>	
<ul style="list-style-type: none"> • Relationship of Post-Operative Troponin I with complications after CABG (Khan MF, et al) 44 		<ul style="list-style-type: none"> • Diagnostic Accuracy of Strain Elastography in Differentiation of Malignant and Benign Breast Lesions Taking Histopathology as Gold Standard (Akhtar J, et al) 90 	
<u>SOFA SCORE THE REAL PREDICTOR OF MORTALITY</u>		<u>USE OF METRONIDAZOLE AFTER HEMORRHOIDECTOMY</u>	
<ul style="list-style-type: none"> • Is Sequential Organ Failure Assessment (SOFA) Score the Real Predictor of Mortality in Resource Constrained Critical Care? (Kausar S, et al) 49 		<ul style="list-style-type: none"> • The Efficacy of Oral Metronidazole in Reducing Pain Post-Hemorrhoidectomy: A Randomized Controlled Trial (Munawar S, et al) 94 	
<u>CAUSES OF EXTRACTIONS OF PERMANENT TEETH</u>		<u>TWO-STAGE (BRACKA) REPAIR IN ADULT HYOSPADIAS</u>	
<ul style="list-style-type: none"> • Pattern and Causes of Extractions of Permanent Teeth in Tertiary Care Hospitals in Karachi, Pakistan (Kumari P, et al) 54 		<ul style="list-style-type: none"> • Outcome of Two-Stage (Bracka) Repair in Adult Hypospadias: Our Experience of Sixty Cases (Afridi RAK, et al) 98 	
<u>IMNCI BASED MORBIDITY PATTERNS OF SICK CHILDREN</u>		<u>PERIMENOPAUSAL FEMALES WITH HIGH BP</u>	
<ul style="list-style-type: none"> • IMNCI Based Morbidity Patterns of Sick Children at Children Hospital CMC Hospital Larkana (Tunio R, et al) 59 		<ul style="list-style-type: none"> • Perimenopausal Females Continue to Have High Blood Pressure Spikes In spite of Taking Regular Antihypertensive Use – Need to Manage the Symptoms Not the Cause (Rana MM, et al) 102 	
<u>DEPRESSIVE DISORDER AMONG TYPE-I DIABETES</u>		<u>VITAMIN C STATUS AMONG STUDENTS DURING COVID-19</u>	
<ul style="list-style-type: none"> • Frequency of Depressive Disorder Among Patients with Type-I Diabetes Mellitus (Kumar A, et al) 64 		<ul style="list-style-type: none"> • Estimation of Vitamin C Status among University Students during COVID-19 Pandemic (Mangi AH, et al) 107 	
<u>ENHANCING CLINICAL SKILLS THROUGH VIDEO</u>			
<ul style="list-style-type: none"> • Enhancing Clinical Skills through Video: An Innovative Health Professions Educational Tool (Mushtaq S, et al) 68 			

Subject	Page No.	Subject	Page No.
<u>PEDIATRIC CHOLEDOCHAL CYST, MANAGEMENT AND OUTCOME</u>		<u>PHYSICAL ACTIVITY WITH BLOOD GLUCOSE LEVELS IN DIABETICS</u>	
<ul style="list-style-type: none"> Pediatric Choledochal Cyst, Symptoms/Signs, Management and Outcome, Our Experience (Bhutta MR, et al) 112 		<ul style="list-style-type: none"> Association of Physical Activity with Blood Glucose Levels in Type 2 Diabetes Mellitus (Shoikat S, et al) 6 	
<u>ACUTE PANCREATITIS IN CHILDREN</u>		<u>INCIDENCE OF SACROILIAC JOINT DYSFUNCTION AS A CAUSE OF LOW BACKACHE</u>	
<ul style="list-style-type: none"> Incidence of Acute Pancreatitis Among Paediatric Patients (Sheikh AH, et al) 116 		<ul style="list-style-type: none"> Incidence of Sacroiliac Joint Dysfunction as a Cause of Low Backache: An Under-Appreciated Pain Generator (Azeem Y, et al) 10 	
<u>SURGICAL SITE INFECTIONS AT SURGICAL WARDS OF LRH HOSPITAL, PESHAWAR</u>		<u>ANEMIA IN DELAYED AND EARLY UMBILICAL CORD CLAMPING</u>	
<ul style="list-style-type: none"> A Retrospective Cross-Sectional Study of Surgical Site Infections at the General Surgical Wards of the LRH Hospital in Peshawar (Bilal M, et al) 120 		<ul style="list-style-type: none"> Frequency of Anemia in Delayed and Early Umbilical Cord Clamping after Birth in Newborn Babies (Tariq A, et al) 14 	
<u>RADIOGRAPHIC ESTIMATION OF AGE USING EPIPHYSEAL FUSION</u>		<u>PRESENTATIONS IN NEWBORNS WITH ABDOMINAL DISTENTION IN EARLY DAYS</u>	
<ul style="list-style-type: none"> Radiographic Estimation of Age Using Epiphyseal Fusion at Elbow Joint in Boys & Girls of 13-17 Years in Gadap Town (Siddiqui RA, et al) 123 		<ul style="list-style-type: none"> Causes and Clinical Presentations in Newborns with Abdominal Distention in Early Days of Life (Zahid MM, et al) 17 	
<u>OUTCOME IN PATIENTS WITH ABDOMINAL TRAUMA</u>		<u>ROLE OF MICRONUTRIENTS IN DENGUE FEVER</u>	
<ul style="list-style-type: none"> Determinants of Outcome in Patients with Abdominal Trauma A Retrospective Cohort Study (Aslam V, et al) 127 		<ul style="list-style-type: none"> From Dengue Fever to Dengue Hemorrhagic Fever: Unveiling the Role of Micronutrients and Immune Modulators (Aamir E, et al) 21 	
<u>INTRADERMAL TRANSAXAMIC ACID IN THE TREATMENT OF MELASMA</u>		<u>CATALASE AND SALIVARY ALPHA AMYLASE WITH DIABETIC AND NON DIABETIC</u>	
<ul style="list-style-type: none"> Efficacy of Intra-dermal Transaxamic Acid in the Treatment of Melasma: A Retrospective Study of Clinical Cases (Naheed A, et al) 130 		<ul style="list-style-type: none"> A Comparison between Catalase and Salivary Alpha Amylase level in Patients with Diabetic and Non Diabetic (Riaz A, et al) 26 	
<u>OUTCOMES OF PECTORALIS MAJOR MYOCUTANEOUS FLAP FOR HEAD AND NECK SURGERIES</u>		<u>ANALYSIS AND VISUALIZATION OF 3 DIMENSIONAL DISSECTION TABLE</u>	
<ul style="list-style-type: none"> Functional Outcomes of Pectoralis Major Myocutaneous Flap for Reconstruction in Head and Neck Surgeries (Memon MI, et al) 134 		<ul style="list-style-type: none"> The Analysis and Visualization of 3 Dimensional Dissection Table in Anatomy Teaching in Cross-Section Images as per Student's Perspective (Sarwat S, et al) 30 	
<u>GYNECOLOGICAL PATHOLOGIES IDENTIFIED DURING DIAGNOSTIC LAPAROSCOPY</u>		<u>WHOLE BODY VIBRATION WITH EXERCISE ON BALANCE AND MUSCULAR STRENGTH IN DIABETIC</u>	
<ul style="list-style-type: none"> Prevalence and Characteristics of Gynecological Pathologies Identified During Diagnostic Laparoscopy: A Cross-Sectional Study In A Diverse Patient Population (Khan N, et al) 139 		<ul style="list-style-type: none"> The Whole Body Vibration comparison with Traditional Tai Chi Exercise on Balance and Muscular Strength in Diabetic Patient with Peripheral Neuropathy (Zia Q, et al) 35 	
Vol. 34, No. 7, July, 2023		<u>EFFECTS OF TARTRAZINE ON ORAL MUCOSAL TISSUES OF WISTAR ALBINO RATS</u>	
Subject	Page No.	<ul style="list-style-type: none"> Detrimental Effects of Tartrazine on Oral Mucosal Tissues of Wistar Albino Rats: An Experimental Study (Khan N, et al) 40 	
<u>VACUUM-ASSISTED WOUND CLOSURE AND ALGINATE DRESSINGS IN SITE INFECTIONS</u>			
<ul style="list-style-type: none"> A Randomized Trial of Vacuum-Assisted Wound Closure and Alginate Dressings in Healing Surgical Site Infections (Zafar M, et al) 2 			

Subject	Page No.	Subject	Page No.
<u>COVID VACCINATION STATUS AND REASONS FOR NON-VACCINATION IN REPRODUCTIVE AGE</u>		<u>EFFECTS OF LEAD (PB) AND CADMIUM (CD) EXPOSURE ON INFERTILITY ISSUES</u>	
<ul style="list-style-type: none"> • Covid Vaccination Status and Reasons for Non-Vaccination in Reproductive Age Women at Tertiary Care Centres (Siddiqui SA, et al) 45 	45	<ul style="list-style-type: none"> • Effects of Lead (Pb) and Cadmium (Cd) Exposure on Infertility Issues in Targeted Male Population in Industrial area of Hayatabad Peshawar Khyber Pakhtunkhwa (Yousaf M, et al) 86 	86
<u>TERTIARY LYMPHOID STRUCTURES IN BREAST DUCTAL CARCINOMA</u>		<u>CLOSED REDUCTION AND PERCUTANEOUS SCREW FIXATION FOR MANAGEMENT OF FRACTURE OF TIBIA</u>	
<ul style="list-style-type: none"> • Correlation of Tertiary Lymphoid Structures in Breast Ductal Carcinoma in Situ and Adverse Pathological Parameters (Naeem N, et al) 50 	50	<ul style="list-style-type: none"> • Functional Outcome of Closed Reduction and Percutaneous Screw Fixation for Management of Schatzker Type I Fracture of Tibia in a Tertiary Care Hospital (Rasool A, et al) 91 	91
<u>VITAMIN D LEVELS WITH SIMULTANEOUS OCCURRENCE OF PERIODONTAL AND CARDIAC DISEASES</u>		<u>C-REACTIVE PROTEIN (CRP) AND PROCALCITONIN IN NON-ALCOHOLIC FATTY LIVER</u>	
<ul style="list-style-type: none"> • Relationship of Vitamin D Levels with Simultaneous Occurrence of Periodontal and Cardiac Diseases (Mehwish R, et al) 55 	55	<ul style="list-style-type: none"> • Level of Serum C-Reactive Protein (CRP) and Procalcitonin in Non-Alcoholic Fatty Liver Disease Patients (Achakzai H, et al) 95 	95
<u>EVALUATION OF PTERYGOID MUSCLES IN TEMPOROMANDIBULAR DISORDERS</u>		<u>SATISFACTION OF DENTAL PROCEDURES IN COVID-19</u>	
<ul style="list-style-type: none"> • Evaluation of Pterygoid Muscles in Patients with Temporomandibular Disorders (Fatima P, et al) 59 	59	<ul style="list-style-type: none"> • Patient Satisfaction of Dental Procedures in Covid-19 Pandemic: A Mixed Method Survey in a Tertiary Care Dental Hospital (Haseeb M, et al) 99 	99
<u>MUCORMYCOSIS DURING THE WAVES OF COVID 19</u>		<u>PERIODONTAL PROBLEMS IN ADOLESCENTS</u>	
<ul style="list-style-type: none"> • Mucormycosis During the Waves of Covid 19 Pandemic; A Cross Sectional Study at Bahawal Victoria Teaching Hospital (Shafique MA, et al) 63 	63	<ul style="list-style-type: none"> • Frequency of Periodontal Problems in Adolescents with On-going Fixed Orthodontic Therapy (Azeem M, et al) 103 	103
<u>DIAGNOSTIC ACCURACY OF E-CADHERIN EXPRESSION IN DIAGNOSIS OF ENDOMETRIOID CARCINOMA</u>		<u>MORTALITY & REINFECTION RATE OF HEPATITIS C AMONG PATIENTS ON HEMODIALYSIS</u>	
<ul style="list-style-type: none"> • Diagnostic Accuracy of E-Cadherin Expression in Diagnosis of Endometrioid Carcinoma in Females (Habib F, et al) 68 	68	<ul style="list-style-type: none"> • One Year Mortality & Reinfection Rate of Hepatitis C Among Patients on Hemodialysis after Successful Direct-Acting Antiviral Treatment (Cheema SS, et al) 107 	107
<u>UTILIZATION AND ACCEPTABILITY OF TELEMEDICINE DURING COVID-19</u>		<u>HYPONATREMIA WITH PREGABALIN</u>	
<ul style="list-style-type: none"> • Utilization and Acceptability of Telemedicine during COVID-19 Pandemic for the Diabetes Care as an Alternative to Physical Examination (Nawaz A, et al) 72 	72	<ul style="list-style-type: none"> • Hyponatremia Associated with Pregabalin (Cheema SR, et al) 112 	112
<u>ALOE VERA GEL ACCELERATED WOUND HEALING IN ORAL MUCOSA</u>		<u>ASYMPTOMATIC HYPERURICEMIA WITH BLOOD PRESSURE IN DIFFERENT AGE</u>	
<ul style="list-style-type: none"> • Aloe Vera Gel Accelerated Wound Healing in Oral Mucosa by Reducing Inflammatory Phase (Rasheed M, et al) 77 	77	<ul style="list-style-type: none"> • Association of Asymptomatic Hyperuricemia with Blood Pressure in Different Age Groups in Semiurban Population of Karachi (Khan A, et al) 116 	116
<u>URIC ACID AND LIPID PROFILE IN ADULT</u>		<u>INTRADIALYTIC HYPERTENSION IN END STAGE RENAL DISEASE</u>	
<ul style="list-style-type: none"> • Association Between Serum Uric Acid and Lipid Profile in Adult Individuals (Anis R, et al) 82 	82	<ul style="list-style-type: none"> • Frequency of Intradialytic Hypertension in End Stage Renal Disease Patients on Maintenance Hemodialysis (Zareen Ullah, et al) 120 	120

Subject	Page No.	Subject	Page No.
<u>CONTINUOUS LOW-DOSE INFUSION OF DEXMEDETOMIDINE IN ICU-INDUCED DELIRIUM</u>		<u>PROBLEMS ENCOUNTERED BY DENTURE-WEARING PATIENTS</u>	
<ul style="list-style-type: none"> To Compare the Efficacy of Continuous Low-Dose Infusion of Dexmedetomidine & Intermittent Boluses of Dexmedetomidine in Preventing ICU-Induced Delirium (Imran S, et al) 124 	124	<ul style="list-style-type: none"> Problems Encountered by Complete-Denture-Wearing Patients (Raja IM, et al) 162 	162
<u>CLOSED-LOOP VENTILATION WITH CONVENTIONAL VENTILATION MODES IN ICU</u>		<u>VALIDITY OF MENTZER INDEX FOR IRON DEFICIENCY ANEMIA IN CHILDREN</u>	
<ul style="list-style-type: none"> Comparative Analysis of Closed-Loop Ventilation Mode with Conventional Ventilation Modes in Intensive Care Unit patients (Haq I, et al) 128 	128	<ul style="list-style-type: none"> Validity of Mentzer Index as a Screening Tool for Iron Deficiency Anemia in Children 6-12 Years of Age (Javed M, et al) 166 	166
<u>CRANIECTOMY FOR SEVERE DIFFUSE TRAUMATIC BRAIN INJURY</u>		<u>DIFFERENT REGIMES IN TREATMENT OF DIABETIC KETOACIDOSIS</u>	
<ul style="list-style-type: none"> Decompressive Craniectomy for Severe Diffuse Traumatic Brain Injury. Study of 25 Cases (Haq N, et al) 133 	133	<ul style="list-style-type: none"> Impact of Different Regimes in Treatment of Diabetic Ketoacidosis and Outcome of Liver Function Test (Wajahatullah M, et al) 170 	170
<u>HYPOGLYCEMIA IN NEONATES</u>		<u>RISK FACTORS OF COVID-19 INFECTION</u>	
<ul style="list-style-type: none"> Frequency of Hypoglycemia in Neonates a Cross-Sectional Study (Manzoor R, et al) 137 	137	<ul style="list-style-type: none"> Risk Factors of COVID-19 Infection: Certainly Uncertainty (Bhutto AR, et al) 174 	174
<u>HYPERURICEMIA IN HYPERTENSIVE PATIENTS</u>		<u>THIRD MOLAR SURGICAL EXTRACTION BY USING ENVELOP AND MODIFIED TRIANGULAR FLAP</u>	
<ul style="list-style-type: none"> Frequency of Hyperuricemia in Hypertensive Patients: A Descriptive Cross-Sectional Study at Karachi (Haq E, et al) 142 	142	<ul style="list-style-type: none"> Comparison of Post-Operative Periodontal Pocket Depth After Mandibular Third Molar Surgical Extraction by Using Envelop Flap and Modified Triangular Flap (Shoib M, et al) 179 	179
<u>IRON DEFICIENCY ANEMIA AMONG 6 TO 60 MONTHS</u>		<u>LAPAROSCOPIC PERITONEAL LAVAGE AND SIGMOIDECTOMY FOR PERFORATED DIVERTICULITIS</u>	
<ul style="list-style-type: none"> Frequency of Iron Deficiency Anemia Among Anaemic Children from Age 6 Months to 60 Months (Ali QA, et al) 146 	146	<ul style="list-style-type: none"> Comparative Analysis of Laparoscopic Peritoneal Lavage and Sigmoidectomy for Treating Perforated Diverticulitis (Cheema HMKN, et al) 184 	184
<u>EDENTULISM PATTERNS IN PAKISTAN BY AGE AND GENDER</u>		<u>ANTIBIOTIC SENSITIVITY IN BLOOD CULTURE POSITIVE TYPHOID</u>	
<ul style="list-style-type: none"> Partial Edentulism Patterns in Pakistan-Stratified by Age and Gender (Rahman A, et al) 150 	150	<ul style="list-style-type: none"> Antibiotic Sensitivity Pattern in Blood Culture Positive Typhoid Fever (Manzoor A, et al) 188 	188
<u>ENDOMETRIAL BIOPSIES PERFORMED FOR ABNORMAL UTERINE BLEEDING</u>		<u>USE OF INJECTABLE DMPA FOR CONTRACEPTION</u>	
<ul style="list-style-type: none"> Clinico-Pathological Study of Endometrial Biopsies Performed for Abnormal Uterine Bleeding: An Audit in a Tertiary Care Centre in Karachi, Sindh, Pakistan (Jalbani A, et al) 154 	154	<ul style="list-style-type: none"> Short-Term Use of Injectable Depot medroxyprogesterone Acetate (DMPA) for Contraception in Immediate Postpartum Women (Gul S, et al) 193 	193
<u>SAFETY OF SOFOSBUVIR AND RIBAVIRIN IN TREATMENT OF CHRONIC HEPATITIS C</u>		<u>ULTRASONOGRAPHICALLY MEASURED LOWER UTERINE SCAR AND OBSTETRIC OUTCOME WITH PREVIOUS C SECTION</u>	
<ul style="list-style-type: none"> Efficacy and Safety of Sofosbuvir and Ribavirin in Treatment of Chronic Hepatitis C Infection G3 of Multi-Transfused Children Aged 4 To 12 Years; A Single Center Study (Sheikh MA, et al) 158 	158	<ul style="list-style-type: none"> Association Between Ultrasonographically Measured Lower Uterine Segment Scar Thickness and Obstetric Outcome in Women with Previous Caesarean (Fatgima SS, et al) 197 	197

Subject	Page No.	Subject	Page No.
<u>MORBIDLY ADHERENT PLACENTA IN PREVIOUS SCAR</u>		<u>KLINTRUP MAKINEN SCORING SYSTEM IN BREAST CARCINOMA</u>	
• Frequency of Morbidly Adherent Placenta in Previous Scar: An Experience in a Tertiary Care Hospital (Sultan S, et al)	202	• Inter Observer Agreement on Klintrup Makinen Scoring System in Breast Carcinoma (Khalid A)	18
<u>PARENTAL ORAL HEALTH LITERACY AND IMPACT ON CHILD'S ORAL HEALTH</u>		<u>PRE-OPERATIVE CARDIAC TROPONIN I LEVELS WITH POST OPERATIVE AFTER CABG</u>	
• Parental Oral Health Literacy and Its Associated Impact On Child's Oral Health Profile Among 15-Years-Old School Children in the Population of Multan, South Punjab, Pakistan (Shoaib M, et al)	207	• Association of Elevated Pre-Operative Cardiac Troponin I Levels with Post Operative Outcome After Coronary Artery Bypass Grafting at Peshawar Institute of Cardiology (Afzal Z, et al)	21
<u>PERIPHERAL NEURECTOMY FOR TRIGEMINAL NEURALGIA</u>		<u>SACROCOCYGEAL TERATOMA IN INFANTS</u>	
• Peripheral Neurectomy: Minimally Invasive Surgical Modality for Trigeminal Neuralgia: A Retrospective Study (Alam W, et al)	212	• Management of Sacrococcygeal Teratoma in Infants and Children (Ramzan M, et al)	24
<u>Timing of Shunt Placement and Infection in Hydrocephalus</u>		<u>VITAMIN D3 LEVEL IN CHILDREN WITH RTI</u>	
• A Retrospective Study on the Association Between the Timing of Shunt Placement and Shunt Infection in Hydrocephalus Associated with Myelomeningocele (Azam F, et al)	216	• Vitamin D3 Level in Children with Respiratory Tract Infection; A Case Control Study (Sultana T, et al)	28
Vol. 34, No. 8, August, 2023		<u>LYMPHEDEMA AFTER TREATMENT OF BREAST CANCER</u>	
Subject	Page No.	• Lymphedema after Treatment of Breast Cancer A Comprehensive Study (Hussain S, et al)	33
<u>HEMOSTASIS SECURING DURING LAPAROSCOPIC APPENDECTOMY</u>		<u>ANALGESIC ACTIVITY OF THE ETHANOLIC EXTRACT OF CUSCUTA REFLEXA STEMS IN MICE</u>	
• Outcome of Hemostasis Securing During Laparoscopic Appendectomy by Using Ligature Versus Vessel Sealing Devices (Munawar H, et al)	2	• Evaluation of Analgesic Activity of the Ethanolic Extract of Cuscuta Reflexa Stems in Albino Wistar Mice (Sameed QA, et al)	38
<u>CORONA ON MENTAL HEALTH AND BEHAVIOR IN CHILDREN</u>		<u>BIOCHEMICAL MARKERS OF CARDIO-VASCULAR DISEASES</u>	
• Effect of Corona Pandemic (Covid-19) on Mental Health And Behavior in Children (Cross Sectional Study) (Mahmood S, et al)	7	• Biochemical Markers of Cardiovascular Diseases : Diagnostic and Prognostic Applications (Shaikh M, et al)	42
<u>THERAPIES TO REDUCE CHOLESTEROL AND GLUCOSE IN DIABETICS</u>		<u>MISSING TEETH AND PROSTHODONTIC TREATMENT MODALITIES</u>	
• Comparison of Therapies to Reduce Total Cholesterol And Glucose Levels in Type II Diabetics (Raza G, et al)	11	• Awareness to Consequences of Missing Teeth and Prosthodontic Treatment Modalities in Partially Dentate Patients Reporting for Dental Extraction (Kainat R, et al)	47
<u>C-REACTIVE PROTEIN IN AMI</u>		<u>CBCT VS DPR FOR ALVEOLAR BONE HEIGHT</u>	
• To Find Out the Role of C-Reactive Protein in Acute Myocardial Infarction as a Marker of Prognosis (Khan M, et al)	15	• Cone Beam Computed Tomography versus Digital Panoramic Radiography for Alveolar Bone Height Assessment: A Prospective Comparative Study (Muqet MA, et al)	52
		<u>CLOSURE OF LOOP ILEOSTOMIES IN TYPHOID PERFORATION</u>	
		• Early Closure of Loop Ileostomies in Typhoid Perforation (Malik S, et al)	57

Subject	Page No.	Subject	Page No.
<u>MACHINE LEARNING AND CT IN THORACIC RADIOLOGY</u>		<u>PEER PRESSURE AND TOBACCO USE AMONG ADOLESCENTS</u>	
• Role of Machine Learning and Computed Tomography in Thoracic Radiology (Noor N, et al) 62	62	• Peer Pressure and Tobacco Use Among Adolescents in Urban Slums (Hamna S, et al) 102	102
<u>CT IN EVALUATING POSTOPERATIVE COMPLICATIONS AFTER THORACIC SURGERY</u>		<u>AMOXICILLIN, DOXYCYCLINE AND CEPHALOSPORIN ACTIVITY AGAINST HUMAN LOWER RESPIRATORY TRACT PATHOGEN</u>	
• Role of Computed Tomography in Evaluating Postoperative Complications After Thoracic Surgery (Baseer A, et al) 67	67	• Comparison of Amoxicillin, Doxycycline and Cephalosporin Activity Against Human Lower Respiratory Tract Pathogen in Cold Season at South Punjab, Pakistan (Naqvi MA, et al) 106	106
<u>URINARY PROSTAGLANDINS WITH URIC ACID IN HYPERURICEMIA</u>		<u>LYCOPENE IN THE MANAGEMENT OF ORAL SUBMUCOUS FIBROSIS</u>	
• Association of Urinary Prostaglandins with Uric Acid in Hyperuricemia Patients (Afzal S, et al) 72	72	• Role of Lycopene in the Management of Oral Submucous Fibrosis (Hussain A, et al) 110	110
<u>THERAPIES FOR EARLY DIAGNOSIS OF PROSTATE CANCER</u>		<u>CAUSES, PREVENTION STRATEGIES AND TREATMENT FOR CHILDHOOD OBESITY</u>	
• Alternative and Complementary Therapies for Early Diagnosis of Prostate Cancer (Ghous MH, et al) 77	77	• Causes, Prevention Strategies and Treatment for Childhood Obesity for Growing Epidemic and its Health Risks (Tariq A, et al) 114	114
<u>BODY MASS INDEX AND INSULIN RESISTANCE IN POLYCYSTIC OVARIAN SYNDROME</u>		<u>EFFECTIVENESS OF LEVATERACETAM IN BRAIN TUMORS</u>	
• Relationship Between Body Mass Index and Insulin Resistance in Patients of Polycystic Ovarian Syndrome (Khan D, et al) 81	81	• Effectiveness of Levateracetam in Patients with Brain Tumors (Haq N, et al) 119	119
<u>HB LEVEL WITH BMI IN ADOLESCENT GIRLS</u>		<u>ROLE OF PLATELET-RICH PLASMA (PRP) IN WOUND HEALING</u>	
• Association of Hemoglobin Level with BMI in Adolescent Girls of Rural Area of Pakistan (Ch SR, et al) 86	86	• Role of Platelet-Rich Plasma (PRP) in Wound Healing after Surgical Extraction Mandibular Third Molar (Ali S, et al) 123	123
<u>PERIODONTAL DISEASES IN DIFFERENT TRIMESTERS OF PREGNANCY</u>		<u>EFFECTIVENESS OF TRIPLE THERAPY IN THE ERADICATION OF H PYLORI</u>	
• Prevalence of Periodontal Diseases in Different Trimesters of Pregnancy Attending Ante-Natal Clinic (Siddique S, et al) 90	90	• A Prospective Study of the Effectiveness of Triple Therapy in the Eradication of H Pylori (Jan N, et al) 128	128
<u>CLOMIPHENE CITRATE ALONE AND PLUS SILDENAFIL CITRATE TO IMPROVE OVULATION</u>		<u>EPIDEMIOLOGY AND PATTERN OF COLONIC DIVERTICULOSIS IN LOWER GI ENDOSCOPY</u>	
• Comparison of Clomiphene Citrate Alone and Clomiphene Plus Sildenafil Citrate to Improve Ovulation Induction in Patients with Unexplained Subfertility (Ashraf S, et al) 94	94	• Epidemiology and Pattern of Colonic Diverticulosis in Patients Undergoing Lower Gastrointestinal Endoscopy: A Descriptive Study at Hayatabad Medical Complex Peshawar (Kashif M, et al) 132	132
<u>CALCIUM CARBONATE VS CALCIUM ACETATE ON HYPERPHOSPHATEMIA IN END STAGE KIDNEY DISEASE</u>		<u>PCNL VS SANDWICH THERAPY OF STAGHORN STONES</u>	
• Comparison Between the Effect of Calcium Carbonate versus Calcium Acetate on Hyperphosphatemia in Patients with End Stage Kidney Disease (Khan S, et al) 98	98	• Comparison of Safety and Effectiveness of Multi Tract Conventional and Mini Percutaneous Nephrolithotomy Simultaneously Versus Sandwich Therapy in Management of Staghorn Stones (Ali L, et al) 136	136

Subject	Page No.	Subject	Page No.
<u>SURGICAL SITE INFECTIONS AFTER IRRIGATION OF APPENDECTOMY WOUNDS</u>		<u>CONTINUOUS HAND SEWEN ANASTOMOSIS WITH SINGLE LAYER EXTRA MUCOSAL INTERRUPTED ANASTOMOSIS</u>	
<ul style="list-style-type: none"> The Frequency of Surgical Site Infections After Irrigation of Appendectomy Wounds with Sterile Saline Solution or Imipenem Solution is Compared. A Retrospective Study (Aslam V, et al) 140 	140	<ul style="list-style-type: none"> Comparison Between the Full Thickness Continuous Hand Sewen Anastomosis with Single Layer Extra Mucosal Interrupted Anastomosis (Hussain SA, et al) 184 	184
<u>MULTIVITAMIN AND IRON THERAPY AGAINST VITAMIN A IN IRON-DEFICIENT WOMEN</u>		<u>PRE-OPERATIVE INVESTIGATIONS ON COMPLICATIONS OF APPENDICECTOMY</u>	
<ul style="list-style-type: none"> Role of Combined Multivitamin and Iron Therapy against Vitamin A in the treatment of Iron-Deficient Anemic Women (Raja AA, et al) 144 	144	<ul style="list-style-type: none"> Impact of Delayed Pre-operative Investigations on Complications of Appendectomy: A Retrospective Study (Rahman S, et al) 188 	188
<u>FACE TO FACE VS ON-LINE TEACHING SATISFACTION OF MEDICAL STUDENTS</u>		<u>KNOWLEDGE OF MEDICAL STUDENTS TOWARDS IRRITABLE BOWEL SYNDROME</u>	
<ul style="list-style-type: none"> Comparison of Face to Face Versus On-Line Teaching Satisfaction of Medical Students (Arooj M, et al) 148 	148	<ul style="list-style-type: none"> Knowledge, Practices and Attitudes of Medical Students towards Irritable Bowel Syndrome: A Cross Sectional Study (Roghani AS, et al) 193 	193
<u>STRENGTH OF DENTURE BASE WITH ARTIFICIAL TEETH</u>		<u>MUCOCUTANEOUS MANIFESTIONS IN HIV</u>	
<ul style="list-style-type: none"> Bench Curing Affecting Bond Strength of Denture Base with Artificial Teeth (Mehtab B, et al) 153 	153	<ul style="list-style-type: none"> Frequency of Mucocutaneous Manifestions in HIV Infected Patients (Riaz S, et al) 199 	199
<u>PRE-ECLAMPSIA AND OBESITY IN PRIMIGRAVIDA</u>		<u>ENZYMES AND NAFLD</u>	
<ul style="list-style-type: none"> Association Between Pre-Eclampsia and Obesity in Primigravida (Nisa S, et al) 158 	158	<ul style="list-style-type: none"> Effect of Probiotics on Liver Enzymes and Non Alcoholic Fatty Liver Disease (Khan JUA, et al) 204 	204
<u>DIABETES MELLITUS IN PATIENTS OF BIOPSY PROVEN LIVER CIRRHOSIS</u>		<u>LOW LEVEL OF FERRITIN IN NEWBORN AND IRON DEFICIENCY ANEMIA IN MOTHERS</u>	
<ul style="list-style-type: none"> Prevalence of Diabetes Mellitus in Patients of Biopsy Proven Liver Cirrhosis (Rehman W, et al) 163 	163	<ul style="list-style-type: none"> Relationship of Low Level of Ferritin in Newborn and Iron Deficiency Anemia in Mothers: A Single Center Study (Saad M, et al) 208 	208
<u>POLYCYSTIC OVARY WITH EFFECT OF LETROZOLE VERSUS CLOMIPHENE</u>		<u>VENTRICULOPERITONEAL SHUNT PLACEMENT FOR HYDROCEPHALUS</u>	
<ul style="list-style-type: none"> Evaluation of Pregnancy Rate in Polycystic Ovary Syndrome Patient with Effect of Letrozole Versus Clomiphene Citrate, Mirpur AJK (Yusuf A, et al) 167 	167	<ul style="list-style-type: none"> Early Versus Delayed Ventriculoperitoneal Shunt Placement For Hydrocephalus Associated With Myelomeningocele: A Systematic Review (Shah SN, et al) 212 	212
<u>THYROID PROFILE AND LIVER FUNCTION TESTS IN PREGNANT WOMEN</u>		<u>NEURO-ENDOSCOPIC PROCEDURES IN THE HYDROCEPHALUS</u>	
<ul style="list-style-type: none"> Evaluation of Thyroid Profile and Liver Function Tests in Pregnant Women of Mirpur AJK (Sarfaraz T, et al) 172 	172	<ul style="list-style-type: none"> Assessing the Effectiveness of Neuro-Endoscopic Procedures in the Treatment of Hydrocephalus (Ali H, et al) 216 	216
<u>KNOWLEDGE OF CONTRACEPTION IN WOMEN OF REPRODUCTIVE AGE</u>		<u>ASSESSING AND MANAGING BREAST NIPPLE DISCHARGE</u>	
<ul style="list-style-type: none"> Knowledge, Attitude and Practice of Contraception in the Women of Reproductive Age in Civil Hospital Karachi (Kulsoom U, et al) 176 	176	<ul style="list-style-type: none"> Comprehensive Multidisciplinary Approach for Assessing and Managing Breast Nipple Discharge: Current Insights and Future Directions (Usman M, et al) 221 	221
<u>ILEOSTOMY REVERSAL WITH AND WITHOUT NG TUBE</u>			
<ul style="list-style-type: none"> Outcome of Ileostomy Reversal With and Without Nasogastric Tube (Khaghan BS, et al) 181 	181		

Subject	Page No.	Subject	Page No.
<u>FNAC OF TESTIS TO RULE OUT OBSTRUCTIVE AND NON-OBSTRUCTIVE AZOOSPERMIA</u>		<u>ANTIBACTERIAL ACTIVITY OF TINOSPORA CORDIFOLIA AND SOLANUM NIGRUM AGAINST E-COLI</u>	
<ul style="list-style-type: none"> Fine Needle Aspiration Cytology of Testis to Rule Out Obstructive and Non-Obstructive Azoospermia (Mahmood R, et al) 225 	225	<ul style="list-style-type: none"> Detection of In-Vitro Antibacterial Activity of Tinospora Cordifolia and Solanum Nigrum Leaf Extracts against Multidrug Resistant-Escherichia Coli (Mahmood S, et al) 11 	11
<u>CT GUIDED BIOPSY OF PULMONARY LESIONS</u>		<u>AETIOLOGY FOR DELAY REPLACEMENT OF MISSING TEETH WITH PROSTHESIS</u>	
<ul style="list-style-type: none"> Outcomes and Complications of Computed Tomography Guided Biopsy of Pulmonary Lesions; Our Experience - A Single Centre Experience (Farman W, et al) 229 	229	<ul style="list-style-type: none"> Determination of Aetiology for Delaying Replacement of Teeth in Partially Edentulous Arches Prosthodontically (Khan MS, et al) 16 	16
<u>VARIOUS TYPES OF TB IN DIABETICS</u>		<u>OXYGEN SATURATION DIFFERENCES BETWEEN FINGERS</u>	
<ul style="list-style-type: none"> The Frequency of Various Forms of Active Tuberculosis in Patients with Diabetes (Basit A, et al) 233 	233	<ul style="list-style-type: none"> Comparison of Oxygen Saturation in Different Fingers of Hands Measured by Portable Pulse Oximeter in Healthy Adults (Khan A, et al) 20 	20
<u>SPONTANEOUS PARTIAL OR COMPLETE ABSORPTION OF HERNIATED LUMBAR DISC</u>		<u>PREVALENCE AND TREATMENT OF MALARIA IN CHILDREN HOSPITALIZED</u>	
<ul style="list-style-type: none"> Clinical Improvement after Spontaneous Partial or Complete Absorption of Herniated Lumbar Disc (Khan Z, et al) 237 	237	<ul style="list-style-type: none"> Prevalence and Treatment of Malaria in Children Hospitalized to Tertiary Care Hospital in Swabi (Rehman S, et al) 24 	24
<u>OUTCOMES OF TOTAL JOINT REPLACEMENT AS HIP OR KNEE</u>		<u>PARAMETERS OF CHRONIC MYELOID LEUKEMIA</u>	
<ul style="list-style-type: none"> Assessing the Long Term Outcomes of Total Joint Replacement Surgeries Such as Hip or Knee Replacement (Zia A, et al) 241 	241	<ul style="list-style-type: none"> Demographic Parameters of Patients Presenting with Chronic Myeloid Leukemia in Mayo Hospital Lahore. A New Epidemiological Addition (Babar H, et al) 28 	28
<u>FAMILY PLANNING MESSAGES AND EFFECTS ON KNOWLEDGE OF CONTRACEPTION</u>		<u>DRUG RESISTANT SALMONELLA</u>	
<ul style="list-style-type: none"> Exposure of Family Planning Messages and its Association with Knowledge, Attitude and Practices of Contraception in Karachi (Kumar S, et al) 245 	245	<ul style="list-style-type: none"> Current Rise of Extensive Drug Resistant Salmonella Cases in a Tertiary Care Hospital of Karachi, Pakistan (Fatima A, et al) 32 	32
Vol. 34, No. 9, September, 2023		<u>RISK OF CARDIOVASCULAR DISEASE IN PATIENTS WITH COPD</u>	
Subject	Page No.	<ul style="list-style-type: none"> Role of Rosuvastatin in Preventing Risk of Cardiovascular Disease in Patients with Chronic Obstructive Pulmonary Disease (Shah MA, et al) 37 	37
<u>SPLENOPORTAL-INDEX DIAGNOSTIC TOOL FOR ESOPHAGEAL VARICES</u>		<u>SELF-EXAMINATION OF LUMP IN BREAST</u>	
<ul style="list-style-type: none"> Splenoportal-Index – A diagnostic Tool for the Non-Invasive Detection of Esophageal Varices (Jami A, et al) 2 	2	<ul style="list-style-type: none"> Awareness of Self-Examination of Lump in the Breast Among Married Women of Karachi (Mehtab K, et al) 41 	41
<u>BETA THALASSEMIA IN FETUSES OF BETA THALASSEMIA CARRIER MOTHERS</u>		<u>SESAMOL AND ITS LIPOSOMAL NANOCARRIER IN COLORECTAL CANCER</u>	
<ul style="list-style-type: none"> Prenatal Diagnosis of Beta-Thalassemia Disease and Fate of Fetuses Among Beta-Thalassemia Carrier Mothers: An Experience at Tertiary Care Hospital of Rahim Yar Khan (Ghafoor MB, et al) 7 	7	<ul style="list-style-type: none"> Comparison of Sesamol and its Liposomal Nanocarrier in the Quest for Anti-Metastatic Agents in Colorectal Cancer Cell Line HT-29 (Liaquat A, et al) 46 	46

Subject	Page No.	Subject	Page No.
<u>BARRIERS TO IMPROVE STUDENTS LEARNING</u>		<u>FORENSIC ANALYSIS OF GUNSHOT RESIDUE</u>	
• The Barriers to Improve Medical Students Learning of Clinical Skills (Ahmed SN, et al) 51	51	• The Forensic Analysis of Gunshot Residue Techniques and Applications (Ali F, et al) 96	96
<u>URETEROSCOPY WITH HOLMIUM LASER FOR RENAL PELVIS STONE</u>		<u>CS WITH UNENGAGED FETAL HEAD IN PRIMIGRAVIDA</u>	
• Outcome of Ureteroscopy with Holmium Laser for Renal Pelvis Stone Clearance (Ali S, et al) 56	56	• To Assess the Association of Cesarean Section with Unengaged Fetal Head in Primigravida Females Presenting in Labour for Delivery (Asif S, et al) 100	100
<u>SKIN DEEP ANXIETY</u>		<u>FORENSIC ODONTOLOGY IN DISASTER VICTIM IDENTIFICATION</u>	
• Skin Deep Anxiety: Investigating Social Appearance Anxiety and Coping Strategies in Patients with Skin Diseases (Naheed A, et al) 61	61	• The Role of Forensic Odontology in Disaster Victim Identification A Case Study Approach (Hammad M, et al) 103	103
<u>DERMATOLOGY OUTPATIENT CARE</u>		<u>STREPTOCOCCAL INFECTION WITH PRETERM PREMATURE RUPTURE OF MEMBRANES</u>	
• Dermatology Outpatient Care: Bridging the Gap Between Patient Expectations and Healthcare Delivery (Ahmad H, et al) 66	66	• Prevalence of Group B Streptococcal Infection in Patients with Preterm Premature Rapture of Membranes (Navid S, et al) 106	106
<u>NUTRITIONAL STATUS AMONG 12-18 YEARS</u>		<u>COMPARE THE POLYCYSTIC OVARIAN SYNDROME TREATED WITH MYOINOSITOL VS METFORMIN</u>	
• Assessment of Nutritional Status Among 12-18 Year Age Group Using Who Z-Scores in Post-Pandemic Era of Covid-19 (Ahmad F, et al) 70	70	• Compare the Mean Change in Homeostatic Model Assessment (HOMA) Index in Polycystic Ovarian Syndrome Patients Treated with Myoinositol Versus Metformin (Gul M, et al) 110	110
<u>POSTOPERATIVE SENSITIVITY BETWEEN AMALGAM AND COMPOSITE IN DENTAL RESTORATIVE TREATMENT</u>		<u>AUTOPSY FINDINGS IN ACUTE METHANOL TOXICITY</u>	
• Compare the Percentage of Postoperative Sensitivity Between Amalgam and Composite in Patients Undergoes Dental Restorative Treatment (Bashir N, et al) 75	75	• Acute Methanol Toxicity: Correlation with Autopsy Findings, A Descriptive Study (Shabbir S, et al) 115	115
<u>ACUTE KIDNEY INJURY TREATED WITH VANCOMYCIN WITH PIPERACILLIN-TAZOBACTAM</u>		<u>ANTERIOR LONGITUDINAL LIGAMENT OSSIFICATION IN VERTEBRAL SPINE</u>	
• A Comparative Analysis of Acute Kidney Injury Incidence in Patients Treated with Vancomycin in Combination with Piperacillin-Tazobactam or Meropenem (Ghous MH, et al) 79	79	• A Study on Anterior Longitudinal Ligament Ossification in Vertebral Spine Patient (Hussain A, et al) 119	119
<u>EMPHYEMA AFTER TUBE THORACOSTOMY IN CHEST TRAUMA</u>		<u>ENDOSCOPIC SINUS SURGERY IN CHRONIC RHINOSINUSITIS</u>	
• Frequency of Empyema After Tube Thoracostomy in Patients with Penetrating Chest Trauma (Baseer A, et al) 84	84	• Outcomes of Endoscopic Sinus Surgery in Patients with Chronic Rhinosinusitis and Immunoglobulin Deficiencies (Nasir A, et al) 123	123
<u>MICRODISCECTOMY VS CONVENTIONAL DISCECTOMY</u>		<u>RISK FACTORS AND COMPLICATIONS OF ACUTE PANCREATITIS</u>	
• Outcomes of Microdiscectomy Versus Conventional Discectomy (Ali F, et al) 88	88	• Pattern of Risk Factors and Complications of Acute Pancreatitis (Hafeez A, et al) 127	127
<u>THROMBOCYTOSIS AND ELEVATED C-REACTIVE PROTEIN LEVELS IN ESOPHAGEAL CARCINOMA</u>			
• Frequency of Thrombocytosis and Elevated C- Reactive Protein Levels in Different Stages of Esophageal Carcinoma in a Tertiary Care Hospital of Karachi (Taj A, et al) 92	92		

Subject	Page No.	Subject	Page No.
<u>IN-DEPTH OF DENTAL AGE ASSESSMENT BY KVAAL TECHNIQUE</u>		<u>KNOWLEDGE OF PARENTS REGARDING ANTIBIOTICS USE IN CHILDREN</u>	
<ul style="list-style-type: none"> An In-Depth Investigation of Dental Age Assessment in the Mardan Population: Employing the Kvaal Technique Via Digital Panoramic Radiography Analysis (Ali A, et al) 131 	131	<ul style="list-style-type: none"> Knowledge, Attitude and Practices of Parents Regarding Antibiotics Use in Children (Masood MS, et al) 173 	173
<u>EFFECTS OF ATORVASTATIN AND ITS LIPOSOMAL PREPARATION IN COLORECTAL CANCER CELL LINE HT-29</u>		<u>UMBILICAL CORD SEPARATION WITH OR WITHOUT ALCOHOL</u>	
<ul style="list-style-type: none"> Comparative Effects of Atorvastatin and its Liposomal Preparation in Impeding the Migratory Potential of Colorectal Cancer Cell Line HT-29 (Shaikh S, et al) 136 	136	<ul style="list-style-type: none"> Comparison of Umbilical Cord Separation with or without Alcohol Application in Newborn (Bashir N, et al) 178 	178
<u>SCREEN TIME ON SLEEP QUALITY AND MENTAL HEALTH IN YOUNG</u>		<u>H. PYLORI INFECTION AMONG DYSPEPTIC PATIENTS</u>	
<ul style="list-style-type: none"> Impact of Screen Time on Sleep Quality and Mental Health in Young Male Students (Akram M, et al) 141 	141	<ul style="list-style-type: none"> Frequency of H. Pylori Infection among Dyspeptic Patients and Correlation with Histopathological Evaluation (arfin SMQ, et al) 182 	182
<u>ADAPTIVE E-TBL DURING COVID-19</u>		<u>WOUND INFECTION TREATED WITH SEGMENTAL SPINAL INSTRUMENTATION</u>	
<ul style="list-style-type: none"> The Effectiveness and Experiences with Adaptive Online Team based Learning (TBL) Amid Social Distancing in Health Sciences (Raza SZ, et al) 145 	145	<ul style="list-style-type: none"> Frequency of Wound Infection in Patients Treated with Segmental Spinal Instrumentation: A Comprehensive Study (Khan MI, et al) 187 	187
<u>HYPOXIC ISCHEMIC ENCEPHALOPATHY BY MRI ASSISTANCE</u>		<u>PREVALENCE OF DENTAL CARIES AMONG SCHOOL CHILDREN</u>	
<ul style="list-style-type: none"> Prevalence and Pattern of Hypoxic Ischemic Encephalopathy in Pediatric Patients Analysis by MRI Assistance (Gul H, et al) 150 	150	<ul style="list-style-type: none"> A Survey on the Prevalence of Dental Caries Among Elementary School Girls of Multan, Pakistan: A Cross-Sectional Study (Siddiqui HMAB, et al) 191 	191
<u>STUDENTS PERCEPTION OF MANIKIN BASED TRAINING</u>		<u>LUNG ULTRASOUND OF NEONATAL RESPIRATORY DISORDERS</u>	
<ul style="list-style-type: none"> Exploring Students Perception of Manikin Based Training in a Medical Skills Lab (Azhar T, et al) 154 	154	<ul style="list-style-type: none"> Role of Lung Ultrasound for the Diagnosis of Neonatal Respiratory Disorders (Rashid Z, et al) 195 	195
<u>KNEE FLEXOR STRENGTH AFTER ANTERIOR CRUCIATE LIGAMENT RECONSTRUCTION</u>		<u>RAISED TSH LEVEL AMONG THIRD TRIMESTER</u>	
<ul style="list-style-type: none"> Influence of Medial Hamstring Tendon Harvest on Knee Flexor Strength after Anterior Cruciate Ligament Reconstruction (Nawaz A, et al) 159 	159	<ul style="list-style-type: none"> Insignificant Variation of Raised Thyroid Stimulating Hormone Level Among Third Trimester of Pregnancy in Diagnosing Subclinical Hypothyroidism (Zafar B, et al) 200 	200
<u>ANTERIOR DECOMPRESSION AND CAGE PLACEMENT REDUCE IN KYPHOTIC ANGLE IN TB SPINE</u>		<u>CERVICAL CANCER WITH COMPLAINT OF ABNORMAL VAGINAL BLEEDING</u>	
<ul style="list-style-type: none"> The Improvement in Kyphotic Angle after Anterior Decompression and Cage Placement in Tuberculosis of Thoracolumbar Spine (Khan Q, et al) 164 	164	<ul style="list-style-type: none"> Prevalence of Cervical Cancer Among Females with Complaints of Abnormal Vaginal Bleeding at Tertiary Care Hospital, Rawalpindi (Iftikhar T, et al) 204 	204
<u>ADMINISTRATION OF CALCIUM GLUCONATE AFTER TOTAL THYROIDECTOMY</u>		<u>ANALYSIS OF INTRAMEDULLARY TUMORS</u>	
<ul style="list-style-type: none"> Role of Prophylactic Administration of Calcium Gluconate in the Early Post Operative Period after Total Thyroidectomy at Surgical department Ayub Teaching Hospital Abbottabad (Khaghan BS, et al) 169 	169	<ul style="list-style-type: none"> Analysis of Intramedullary Tumors; Experience at Tertiary Care Hospital (Haq N, et al) 208 	208
		<u>SMARTPHONE ADDICTION'S INFLUENCE ON ADOLESCENT HEALTH</u>	
		<ul style="list-style-type: none"> The Mobile Paradox: Examining Smartphone Addiction's Influence on Adolescent Health and Aggression (Nawaz B, et al) 213 	213

Subject	Page No.	Subject	Page No.
<u>MISCONCEPTIONS TOWARDS HIV/AIDS AMONG UNIVERSITY STUDENTS</u>		<u>MORTALITY IN PATIENTS WITH ACUTE MI COMPLICATED WITH VENTRICULAR SEPTAL RUPTURE</u>	
<ul style="list-style-type: none"> Knowledge and Misconceptions Towards HIV/AIDS Among the University Students: A Review (Siddiqui SJ, et al) 217 		<ul style="list-style-type: none"> Status of In-Hospital Mortality in Patients with Acute ST-Elevation Myocardial Infarction Complicated with Ventricular Septal Rupture (Hussain S, et al) 36 	
Vol. 34, No. 10, October, 2023		<u>INTERNAL WITH EXTERNAL OSTEOTOMY IN PATIENTS WITH COSMETIC RHINOPLASTY</u>	
Subject	Page No.	<ul style="list-style-type: none"> Comparative Assessment of Internal Osteotomy with External Osteotomy in Patients Undergoing Cosmetic Rhinoplasty; A Randomized Controlled Trial (Malik QA, et al) 41 	
<u>CHILD DEVELOPMENT WITH OROFACIAL CLEFTS</u>		<u>ULTRASONOGRAPHY IN DIAGNOSIS OF INTUSSUSCEPTIONS IN CHILDREN</u>	
<ul style="list-style-type: none"> Identify Different Challenges throughout Child Development with Orofacial Clefts (Abid MB, et al) 3 		<ul style="list-style-type: none"> Diagnostic Accuracy of Ultrasonography in Diagnosis of Intussusceptions in Pediatric Patients (Aqeel R, et al) 45 	
<u>RELATIONSHIP OF KERATOCUNUS TO CONSANGUINITY</u>		<u>SOCIODEMOGRAPHIC PROFILE OF LHWS WITH BARRIERS IN BASIC SERVICE DELIVERY</u>	
<ul style="list-style-type: none"> Assessing the Relationship of Keratocunus to Consanguinity (Rafe A, et al) 9 		<ul style="list-style-type: none"> Sociodemographic Profile of Lady Health Workers and Its Association with Barriers in Basic Service Delivery in Pakistan (Sangrasi MS, et al) 49 	
<u>MOODLE' AS A FORMATIVE ASSESSMENT - PERCEPTION OF MEDICAL STUDENTS</u>		<u>ANTI-CANCER ACTIVITY OF GREWIA ASIATICA ETHANOLIC ON SF767 CELL LINE</u>	
<ul style="list-style-type: none"> Introducing 'MOODLE' as a Formative Assessment Tool: Perception of Medical Students (Abbas M, et al) 13 		<ul style="list-style-type: none"> Potential Anti-Cancer Activity of Grewia Asiatica Ethanolic Extract on Sf767 Cell Line (Aftab R, et al) 53 	
<u>DIAGNOSTIC ACCURACY OF THROMBOCYTOPENIA</u>		<u>EARLY DETECTION OF ALZHEIMER'S DISEASE</u>	
<ul style="list-style-type: none"> Diagnostic Accuracy of Thrombocytopenia and Fib 4 Score in Predicting Esophageal Varices Using Endoscopy as Gold Standard Among Patients with Chronic Liver Disease (Qaiser MA, et al) 18 		<ul style="list-style-type: none"> Developing New Biomarkers for the Early Detection of Alzheimer's Disease (Shafique S, et al) 59 	
<u>MYOCARDIAL INFARCTION COMPLICATED BY COMPLETE HEART BLOCK TREATED WITH PERCUTANEOUS CORONARY INTERVENTION</u>		<u>EARLY DETECTION OF ORAL POTENTIALLY MALIGNANT DISORDERS</u>	
<ul style="list-style-type: none"> Outcomes of Late Presenters with Acute Anterior Wall Myocardial Infarction Complicated by Complete Heart Block Treated with Percutaneous Coronary Intervention (Rahim A, et al) 23 		<ul style="list-style-type: none"> Evaluation of Salivary Biomarkers for Early Detection of Oral Potentially Malignant Disorders: A Prospective Study (Zareef U, et al) 63 	
<u>ANALYSIS OF DIAGNOSTIC ACCURACY WITH LOW DOSE VS STANDARD DOSE WITH CTF</u>		<u>PROPHYLACTIC PHENYLEPHRINE INFUSION IN PREVENTION OF SUBARACHNOID BLOCK</u>	
<ul style="list-style-type: none"> Analysis of Diagnostic Accuracy with Low Dose versus Standard Dose Computed Tomography Fluoroscopy Guided Biopsies (Malik M, et al) 27 		<ul style="list-style-type: none"> Effectiveness of Prophylactic Phenylephrine Infusion in Prevention of Subarachnoid Block Associated Hypotension for Elective Cesarean Section (Ahmed N, et al) 68 	
<u>COMPARATIVE STUDY OF SIMVASTATIN AND ROSUVASTATIN</u>		<u>TUBERCULOSIS AND CHRONIC KIDNEY DISEASE</u>	
<ul style="list-style-type: none"> Analyzing Hepatotoxicity: A Comparative Study of Simvastatin and Rosuvastatin, and Their Reversal Using Montelukast and Coenzyme Q10 (Jahan S, et al) 31 		<ul style="list-style-type: none"> Tuberculosis and Chronic Kidney Disease: Evaluating the Management Challenges and Treatment Strategies (Ahmad M, et al) 72 	

Subject	Page No.
<u>ANTIHYPERTENSIVE DRUGS IN HTN PATIENT WITH NON-INSULIN DEPENDENT DM</u>	
• Prescribing Pattern of Antihypertensive Drugs in Hypertensive Patient with Non-Insulin Dependent Diabetes Mellitus (Khan FA, et al)	76
<u>NEWBORN CARE PRACTICES IN PAKISTAN</u>	
• Newborn Care Practices in Pakistan – A Study in Community (Amir A, et al)	81
<u>GREWIA ASIATICA ETHANOLIC EXTRACT ON HEPG2 CELL LINE</u>	
• Ethanolic Extract of Grewia Asiatica: A Promising Anti-Cancer Agent Against HepG2 Cell Line (Butt AF, et al)	85
<u>CLINICOPATHOLOGICAL AND SURGICAL APPROACHES IN THYROID CARCINOMA WITH HASHIMOTO'S THYROIDITIS</u>	
• Exploring the Clinicopathological Attributes and Surgical Approaches in Cases of Papillary Thyroid Carcinoma Occurring Concurrently with Hashimoto's Thyroiditis (Mangrio SA, et al)	90
<u>CHEST INJURIES CAUSED BY BLUNT TRAUMA - AUTOPSY ANALYSIS</u>	
• A Study Examining Chest Injuries Caused by Blunt Trauma Through Prospective Autopsy Analysis (Kamran QA, et al)	95
<u>KNOWLEDGE OF PARENTS ABOUT BETA THALASSEMIA PATIENTS</u>	
• Knowledge, Attitude, and Practices of Parents of Beta Thalassemia Patients (Masood MS, et al)	99
<u>COMPARISON OF SOFOSBUVIR + VELPATASVIR VS SOFOSBUVIR + DACLATASVIR IN CHRONIC HCV</u>	
• Comparison of Drug Regimens (Sofosbuvir and velpatasvir versus Sofosbuvir and Daclatasvir) in Treatment of Chronic Hepatitis C Virus Patients in Terms of Efficacy and Safety (Latif U, et al)	104
<u>FREQUENCY OF DIFFUSE LARGE B CELL LYMPHOMA</u>	
• Frequency of Immune-Histochemical Markers-Based Diffuse Large B Cell Lymphoma Subgroups (Mahmood R, et al)	109
<u>HEPATORENAL SYNDROME IN CIRRHOTIC PATIENTS</u>	
• Frequency of Hepatorenal Syndrome in Cirrhotic Patients Presenting with Spontaneous Bacterial Peritonitis (Hidayat R, et al)	114

Subject	Page No.
<u>B-SCAN ULTRASONOGRAPHY FOR CAUSES OF LOW VISION IN DIABETIC RETINOPATHY</u>	
• Use of B-Scan Ultrasonography is Determining the Causes of Low Vision in Patients with Diabetic Retinopathy (Liaqat S, et al)	120
<u>URINARY AND SEXUAL OUTCOMES AFTER ANASTOMOTIC URETHROPLASTY</u>	
• Urinary and Sexual Outcomes After Anastomotic Urethroplasty Performed For Posterior Urethral Distraction Defect (Haque E, et al)	125
<u>GINGIVAL DEPIGMENTATION USING DIODE LASER WITH CONJUNCTIVE</u>	
• Gingival Depigmentation Using Diode Laser with the Conjunctive Application of Vitamin C: A Case Series (Alasmari D)	129
<u>OSSIFYING FIBROMA WITH A SECONDARY ANEURYSMAL BONE CYST IN MAXILLARY SINUS</u>	
• A Rare Presentation of Ossifying Fibroma with a Secondary Aneurysmal Bone Cyst in the Maxillary Sinus – A Case Report (Alotaibi KZ)	134

Vol. 34, No. 11, November, 2023

Subject	Page No.
<u>USE OF PLANT EXTRACTS AS ALTERNATIVE TREATMENT IN UTI</u>	
• Multidrug Resistant- Escherichia coli – Use of Plant Extracts as Alternative Treatment in Urinary Tract Infection (Arif H, et al)	3
<u>DEPRESSION AMONG PARENTS OF NEWBORNS ADMITTED TO ICU</u>	
• Burden and Determinants of Depression among Parents of Newborns Admitted To Neonatal Intensive Care Unit (Hayat K, et al)	8
<u>ORAL HEALTH IN POST TREATMENT OF ORAL SQUAMOUS CELL CARCINOMA</u>	
• Assessment of Common Oral Dysfunction and Their Impact on Oral Health Related Quality of Life in Post Treatment Patients of Oral Squamous Cell Carcinoma : A Prospective Study (Afzal A, et al)	13
<u>DIABETIC RETINOPATHY WITH THE SEVERITY OF CORONARY ARTERY DISEASE</u>	
• Evaluation of the Relationship Between Diabetic Retinopathy with the Severity of Coronary Artery Disease in Patients with Non ST Elevation Myocardial Infarction (NSTEMI) (Din N, et al)	17

Subject	Page No.	Subject	Page No.
<u>TOOTH AGENESIS IN ORTHODONTIC PATIENTS</u>		<u>KNOWLEDGE OF STUDENTS ABOUT RADIATION PROTECTION PROTOCOLS</u>	
<ul style="list-style-type: none"> Prevalence of Tooth Agenesis in Orthodontic Patients in Nishtar Institute of Dentistry (Gaffar I, et al) 21 	21	<ul style="list-style-type: none"> Assessment of Knowledge, Attitude and Practice of Undergraduate Final Year Dental Students and House Officers Regarding Radiation Protection Protocols: A Question Based Cross-Sectional Study (Hasan SJ, et al) 60 	60
<u>PSYCHOLOGICAL WELL-BEING OF MEDICAL STUDENTS</u>		<u>ANGIOPLASTY COMPARED WITH CORONARY STENTING IN NARROW CORONARY ARTERIES</u>	
<ul style="list-style-type: none"> Quantitative Analysis of Hope, Gratitude, and Satisfaction as Predictors of Psychological Well-Being in Medical Students (Qureshi T, et al) 25 	25	<ul style="list-style-type: none"> Effectiveness of Balloon Angioplasty Compared with Coronary Stenting in Narrow Coronary Arteries (Nawaz T, et al) 66 	66
<u>RELATIONSHIP BETWEEN PERIODONTAL DISEASE AND SYSTEMIC HEALTH</u>		<u>VITAMIN-D DEFICIENCY WITH CHRONIC LIVER DISEASE</u>	
<ul style="list-style-type: none"> A survey on the Awareness of Interrelationship of Periodontal Disease and Systemic Health Among Qassim Population (Alotaibi Y, et al) 28 	28	<ul style="list-style-type: none"> Vitamin-D Deficiency in Patients with Chronic Liver Disease: A Single Center Cross-Sectional Study (Daud M, et al) 71 	71
<u>OPTIC NERVE CLASSIFICATION BASED ON DELANO'S SYSTEM</u>		<u>TRANEXAMIC ACID IN THE MANAGEMENT OF INTRACEREBRAL HEMORRHAGE</u>	
<ul style="list-style-type: none"> Analyzing the Occurrence of the Various Types of Optic Nerve Classification Based on DeLano's System in Gender on Both Sides (Lakhani M, et al) 33 	33	<ul style="list-style-type: none"> Efficacy and Safety of Tranexamic Acid in the Management of Hyper Acute Spontaneous Intracerebral Hemorrhage (Khan S, et al) 75 	75
<u>POSTOPERATIVE COMPLICATIONS AND SATISFACTION IN THIRD MOLAR EXTRACTION</u>		<u>RISK MANAGEMENT STRATEGIES IN HEALTHCARE INSTITUTIONS</u>	
<ul style="list-style-type: none"> Evaluation of Postoperative Complications and Patient Satisfaction in Third Molar Surgical Extraction Techniques: A Clinical Study (Ahmad F, et al) 37 	37	<ul style="list-style-type: none"> Assessing the Efficacy of Risk Management Strategies in Healthcare Institutions for Error Reduction and Patient Safety Enhancement (Rashid H, et al) 80 	80
<u>ERADICATION OF HELICOBACTER PYLORI INFECTION</u>		<u>IMPROVEMENT IN LVF IN PATIENTS HAVING PERCUTANEOUS CORONARY INTERVENTION</u>	
<ul style="list-style-type: none"> Eradication of Helicobacter Pylori Infection; The Efficacy of High Dose Vonoprazan and Amoxicillin Dual Therapy (Shah MA, et al) 42 	42	<ul style="list-style-type: none"> Improvement in Left Ventricular Function in Patients Having Percutaneous Coronary Intervention (PCI) for Acute ST-Elevation Myocardial Infarction (STEMI) with Late Presentation (Khan A, et al) 85 	85
<u>DATING SCAN BY WOMEN OF REPRODUCTIVE AGE</u>		<u>DISEASES AND DEMOGRAPHIC CHARACTERISTICS IN DERMATOLOGY DEPARTMENT</u>	
<ul style="list-style-type: none"> Unveiling the Challenges: Why Dating Scan Remain Elusive for Many Women? (Sultana N, et al) 46 	46	<ul style="list-style-type: none"> Pattern of Diseases and Demographic Characteristics of Admitted Patients in the Dermatology Unit, Hayatabad Medical Complex, Peshawar (Naheed, et al) 89 	89
<u>CORD BLOOD PH IN MECONIUM-COATED NEONATES</u>		<u>MINI-PERCUTANEOUS-NEPHROLITHOTOMY IN SOLITARY FUNCTIONING KIDNEYS</u>	
<ul style="list-style-type: none"> Cord Blood pH in Meconium-Coated Neonates: A Cross Sectional Study (Sikandar MZ, et al) 52 	52	<ul style="list-style-type: none"> Outcome of Mini-Percutaneous-Nephrolithotomy in Solitary Functioning Kidneys, Single Centre Experience (Nawaz A, et al) 93 	93
<u>PREVALENCE AND CAUSES OF RENAL DISORDERS</u>		<u>B-SCAN ULTRASONOGRAPHY</u>	
<ul style="list-style-type: none"> Prevalence and Causes of Renal Disorders in Patients Undergoing Ultrasound-Guided Renal Biopsy: A Cross-Sectional Study (Khan MAW, et al) 56 	56	<ul style="list-style-type: none"> Diagnostic Application of B-scan Ultrasonography (Khan SA, et al) 97 	97

Subject	Page No.
<u>PREGNANCY RELATED ACUTE KIDNEY INJURY</u>	
• Etiology and Outcome of Pregnancy Related Acute Kidney Injury (Khan Z, et al)	102
<u>HYPERURECMIA WITH HYPERTHYROIDISM</u>	
• Association of Hyperurecemia with Hyperthyroidism. A Cross Sectional Descriptive Study (Javaid F, et al)	106
<u>KNOWLEDGE OF CROSS INFECTION CONTROL AMONG DENTAL PRACTITIONERS</u>	
• Knowledge and Awareness of Cross Infection Control Among Dental Practitioners in a Tertiary Care Hospital in Islamabad (Arshad O, et al)	110
<u>MICROCOMPUTED TOMOGRAPHY IN ENDODONTICS</u>	
• Micro-Computed Tomography as a Transformative Tool in Endodontics: A Narrative Review (Agwan MAS)	115
<u>SOFT TISSUE RECONSTRUCTION WITH FREE GINGIVAL GRAFT TECHNIQUE</u>	
• Soft Tissue Reconstruction with Free Gingival Graft in the Anterior Mandibular Region Prior to Orthodontic Management - A Case Report (Iqbal S, et al)	121

Vol. 34, No. 12, December, 2023

Subject	Page No.
<u>COMPARISON OF LEF WITH MTX AND HYDROXYCHLOROQUINE FOR RA</u>	
• Comparison of Leflunomide Monotherapy with Combination of Methotrexate and Hydroxychloroquine in Active Rheumatoid Arthritis (Panzai MMK, et al)	3
<u>NON-OPIOID VERSUS OPIOID ANALGESIA IN NEURO-SURGERY</u>	
• Comparison Between Non-Opioid Versus Opioid Analgesia in Neuro-Surgery (Hameed J, et al)	7
<u>NALBUPHINE FOR MANAGING POST-ANESTHESIA SHIVERING</u>	
• Efficacy of Intravenous Nalbuphine for Managing Post-anesthesia Shivering (Ali A, et al)	12
<u>CONGENITAL TALIPES EQUINOVARUS (CTEV) TREATED WITH PONSETI TECHNIQUE</u>	
• Achilles Tenotomy in Patients with Congenital Talipes Equinovarus (CTEV) Treated with Ponseti Technique (Shafi MK, et al)	16

Subject	Page No.
<u>SHEAR BOND STRENGTH OF DIFFERENT METHODS OF ORTHODONTIC BONDING</u>	
• Comparison of Shear Bond Strength of Different Methods of Orthodontic Bonding (Gaffar I, et al)	20
<u>MRW AND RNFL IN DETECTION OF GLAUCOMA</u>	
• Comparing Diagnostic Accuracy of Minimum Rim Width (MRW) and Retinal Nerve Fibre Layer (RNFL) in Detection of Glaucoma (Saad Ullah M, et al)	24
<u>CONGENITAL HEART DISEASE FOR ECHOCARDIOGRAM BY PEDIATRICIAN</u>	
• Echocardiographic Evaluation Reveals the Prevalence and Patterns of Congenital Heart Disease in Pediatric Populations: Insights from a Peripheral Cardiac Center in Azad Jammu and Kashmir (Ahmed S, et al)	28
<u>PELVIC ORGAN PROLAPSE AND URINARY SYMPTOMS IN WOMEN</u>	
• Relationship Between Pelvic Organ Prolapse and Urinary Symptoms in Women (Aleem A, et al)	31
<u>AVF AND HEMOGLOBIN LEVELS IN HEMODIALYSIS-DEPENDENT ESRD PATIENTS</u>	
• Association Between Arteriovenous Fistula (AVF) and Hemoglobin Levels in Hemodialysis-Dependent End-Stage Renal Disease (ESRD) Patients (Khan MAW, et al)	35
<u>RISK FACTORS OF DENTAL CARIES</u>	
• Prevalence and Risk Factors of Dental Caries among Patients Seeking Care at Tertiary Hospitals (Khan K, et al)	39
<u>ESTIMATE NUMBER OF ENTERIC PATHOGENS IN RAW CHICKEN</u>	
• Microbiological Profile and Susceptibility Pattern of Enteric Organisms in Raw Broiler Chicken Meat From Abattoirs of Lahore, Pakistan (Javed N, et al)	44
<u>SPONTANEOUS BACTERIAL PERITONITIS IN WITH CIRRHOTIC ASCITES</u>	
• Frequency of Spontaneous Bacterial Peritonitis in Asymptomatic Outpatients with Cirrhotic Ascites (Fiaz B, et al)	49
<u>MANAGEMENT OF BLUNT LIVER TRAUMA</u>	
• Etiology and Management of Blunt Liver Trauma in a Tertiary Care Hospital (Uzma, et al)	53
<u>DEPRESSION AND ANXIETY AMONG MELASMA PATIENTS</u>	
• Frequency of Depression and Anxiety Among Melasma Patients Presented at Tertiary Care Hospital (Erfan M, et al)	58

Subject	Page No.	Subject	Page No.
<u>ACCURACY RIPA-SA SCORE IN DETECTING ACUTE APPENDICITIS</u>		<u>PO PAIN BETWEEN STANDARD VS MINI-PERCUTANEOUS NEPHROLITHOTOMY</u>	
<ul style="list-style-type: none"> Diagnostic Accuracy of RIPA-SA Score in Detecting Acute Appendicitis (Hunain, et al) 62 	62	<ul style="list-style-type: none"> Comparison of Post-Operative Pain Between Standard Versus Mini-Percutaneous Nephrolithotomy (Ahmad A, et al) 79 	79
<u>VITAMIN-D DEFICIENCY WITH ALOPECIA AREATA</u>		<u>ASSESSMENT OF CHILDHOOD VACCINE COVERAGE: EMERGING METHODOLOGIES</u>	
<ul style="list-style-type: none"> Prevalence of Vitamin-D Deficiency among Individuals Diagnosed with Alopecia Areata (Saleem S, et al) 66 	66	<ul style="list-style-type: none"> Emerging Methodologies Used in Assessing Childhood Vaccination Coverage: A Comparative Scoping Review (Sahitia S, et al) 84 	84
<u>PCI OF ASYMPTOMATIC PATIENTS WITH ISCHEMIA BETWEEN 12 AND 48 HOURS</u>		<u>AYURVEDIC ANTIOXIDANT REGIMEN</u>	
<ul style="list-style-type: none"> Improvement in LV Functions After 40 Days Following PCI of Asymptomatic Patients with Ischemia between 12 and 48 Hours (Rehmat S, et al) 70 	70	<ul style="list-style-type: none"> Nishamalaki Drug- An Ayurvedic Antioxidant Regimen for Periodontal Diseases and Diabetes Mellitus? (Mohammad NKS) 90 	90
<u>LITHOTRIPTERS FOR THE TREATMENT OF URETERIC STONES</u>		<u>DISCOLOURED NONVITAL TOOTH USING THE WALKING BLEACH TECHNIQUE</u>	
<ul style="list-style-type: none"> Comparative Efficacy and Safety of Hand-Held and Conventional Intra-Corporeal Pneumatic Lithotripsy in the Treatment of Ureteric Stones (Orakazi, AN, et al) 74 	74	<ul style="list-style-type: none"> Treatment of Discoloured Nonvital Tooth Using the Walking Bleach Technique: A Case Report (Javed MQ, et al) 95 	95
		<u>IMMOTILE CILIA SYNDROME</u>	
		<ul style="list-style-type: none"> Immotile Cilia Syndrome: A case Report (Mustafa G) 99 	99